



Silymarin effect as an antioxidant to improve damages induced by CCl₄ on some characteristics of male rats reproductive system

Measer A. Ahmed¹, Aziz K. Hamid², Hussien M. Tayawi³

¹ Department of Biology, College of Education for pure Science, Tikrit University, Tikrit, Iraq

² Department of Biology, College of Science, Tikrit University, Tikrit, Iraq

³ Sallahdin department, Ministry of Education, Iraq

DOI: <http://dx.doi.org/10.25130/tjps.23.2018.029>

ARTICLE INFO.

Article history:

-Received: 5 / 10 / 2017

-Accepted: 11 / 12 / 2017

-Available online: / / 2018

Keywords: silymarin, CCl₄, sperm, antioxidant.

Corresponding Author:

Name: Measer A. Ahmed

E-mail:

measer80@gmail.com

Tel: +9647701827668

Affiliation:

Introduction

Infertility and its related problems are a major disturbing issues in people life [1]. The common cause of infertility in males is their inability to produce a sufficient number of active sperms [2,3]. Many studies results approved that oxidative stress can affect sperm production and reduce sperm count by production of free radicals and oxidation of germ cells in the testis tissue [4,5]. Studies have shown that using antioxidants can effectively treat infertility in males through reduction of damages caused by free radicals, strengthening the blood-testis barrier and protecting and repairing of sperm DNA [5,6]. The effectiveness of herbs has been validated during the past decades through modern research. These herbs generally contain antioxidants, cell membrane-stabilizing agents, or nutrients that inhibit the depletion of endogenous antioxidants such as glutathione. Silymarin is the major active compound from milk thistle extract (*Silybum marianum*). It consists of seven flavonoglignans (silibinin, isosilibinin, silychristin, isosilychristin and silydianin) and a flavonoid (taxifolin) which is a unique type of bioflavonoid and an antioxidant that scavenges damaging free radicals. [7].

Abstract

This study was conducted to investigate if silymarin can prevent the adverse effect of oxidative stress induced by carbon tetrachloride (CCl₄) on some characters of male rat reproductive system. Twenty male albino rats were divided into 4 groups, 5 for each included, control group, CCl₄ group, silymarin group and silymarin plus CCl₄ group. Groups with silymarin received dose (150mg/kg) and group with CCl₄ received (3ml/kg) at the end of experiment. The period of the study was 35 days. CCl₄ can induce male reproductive toxicity through damage testis structure, weights, reduction in epididymis, sperm count, morphology, viability and motility. Silymarin plays its antioxidant role with scavenging free radical resulted by CCl₄ and showed significant increasing in body and testis weight, normal sperms counting and vital sperms in compared to CCl₄ group.

studies have been done till now on the therapeutic and biological properties of silymarin. Among its properties can point to anti-inflammatory, antioxidant, anti-cancer and hepatoprotective properties [8,9]. Silymarin plays antioxidant role with scavenging free radical as well as increasing the levels of glutathione peroxidase and superoxide dismutase (SOD) [10].

This study was performed to investigate if silymarin can prevent the adverse effect of oxidative stress induced by CCl₄ on some characters of male rat reproductive system.

Materials and methods


Twenty healthy male albino rats with an average weight of 200-250 g were used in this experimental study. During the experiment, the rats were kept in the condition of 12 hours of light and 12 hours of darkness. Water and food were given to them without any restrictions during the period of the experiment which was (35) days. The animals were then divided randomly into four groups 5 for each including control, CCl₄ group was given (3ml CCl₄) before 24 hours of the test end, silymarin group and silymarin plus CCl₄ group. Groups with silymarin received dose (150mg/kg) of silymarin.

At the end of the test and after weighing, the rats were anesthetized using chloroform and blood samples were directly taken from hearts using a 5 ml syringe. The blood serum was then collected using centrifuge (3000 rpm for 15 min) and kept at -20°C. serum were used to measure creatinine GOT and GPT levels by spectrophotometer and analysis kits. The testis were removed and tissue sections obtained. The sections were then stained with Hematoxylin and Eosin stain. counting and checking the cells were made using light microscopy. epididymis pounding in


physiological solution, slides made to determine viability and mortality of sperms using eosin-nicrosin stain. Haemocytometer were used to find total sperm count using eosin stain.

Data to each group were recorded and analyzed using SPSS and one-way analysis of variance (one-way ANOVA) test. P<0.05 was used to show significancy. Data in the section of Results were calculated and compared as Mean±SEM.


Results




figure(1) GOT(U/L)




figure(2) GPT(U/L)




figure(3) creatinine (mg/dl)




figure(4) weight




figure(5) weights of testis and epididymis



figure(6) normal and abnormal sperm %




figure(7) vital and dead sperm%




figure(8) sperm count * 10^6




(A) control group



(B) CCl4 group



(C) Silymarin



(D) Silymarin+CCl4

figure(9). photograph of testicular tissue (H&E.,40x)

The average level of GOT ,GPT, Creatinine, normality of sperms , epididymis weight vitality and sperm count (figure:1,2,3,5,6,7, 8) in Group received

silymarin showed no significant change at level $P \leq 0.05$ compared to the control group, While there were increasing in body and testis weight (figure:4,5).

Group received CCl₄ showed a significant increasing in GOT, GPT, creatinine, sperm abnormality and mortality (figure: 1,2,3,6,7) compared to the control group, where CCl₄ group showed a significant decreasing in body and testis weight, normal sperm, sperm vitality and sperm count (figure: 4,5,6,7,8) at level $P \leq 0.05$ compared to control group. CCl₄+silymarin group showed a significant decreasing in level of GOT, GPT, creatinine abnormality and mortality of sperms (figure:1,2,3,6,7) in compared to CCl₄ group, while there were a significant increase in body and testis weight, normal sperm, sperms counting and vital sperms (figure:4,5,6,7,8) in compared to CCl₄ group. In CCl₄ group, testis seminiferous tubules (figure 9-B) are almost empty of spermatozoa. Most cells have damage to cellular components. CCl₄+silymarin group (figure 9-D) Most of the cells in the stages of morphological transition, the tail of the sperm are thin and the sperm offspring are similar to the normal state but less dense than the normal state. Silymarin group Tissue (figure 9-C) is nearly similar to the control (figure 9-A).

Discussion

CCl₄ can impair male reproductive function by inducing oxidative stress (11). Silymarin is a potent antioxidant and could compensate the adverse effect of CCl₄ on GOT, GPT enzymes creatinine (figure 1,2,3) and some sperm characters (figure 9-B). Viability and motility are important parameters of mature sperm which indicate its structural and functional quality to move toward an egg for successful fertilization (12). CCl₄ can induce male reproductive toxicity through damage in testis structure, decrease in testosterone level (13), testis and epididymis weights, reduction in epididymis sperm count, normal morphology, viability and motility (figure 4,5,6,7,8). CCl₄ is proposed to exert its cytotoxicity by free radicals generation and the activation of oxidative sensitive signaling pathways, which lead to a hepatotoxicity in result to form intermediate radicals Trichloromethyl CCl₃ which metabolize by cytochrome p450 (13). Generating free radicals cause peroxidation of hepatic membrane and raise of liver enzymes (GOT, GPT) and increasing in creatinine level (figure 1,2,3) because of inhibiting of insulin level which induce cells to use proteins as a main source for energy production in addition to the increasing of oxidation (14).

References

1. Eskandari F and Momeni H (2016). Protective effect of silymarin on viability, motility and mitochondrial membrane potential of ram sperm treated with sodium arsenite Int J Reprod BioMed Vol. 14. No. 6. pp: 397-402.
2. Amin A and Hamza AA (2006). Effects of Roselle and Ginger on cisplatin-induced reproductive toxicity in rats. Asian J Androl. 8(5): 607-12.
3. Imai H, Suzuki K, Ishizaka K, Ichinose S, Oshima H and Okayasu I (2001). Failure of the expression of

silymarin is a polyphenolic flavonoid with a potent antioxidant property which not only acts as free radical scavenger but also increases the capacity of cell antioxidant enzymes (10,12). Silymarin has long been used as a hepatic protective remedy (15). In addition to central effects of silymarin on hypothalamus – pituitary - testis axis, increased testosterone levels in the study showed related silymarin to synthesis and metabolism of this hormone. Silymarin is counted as a potent inhibitor of aromatase enzyme. Aromatase catalyzes the conversion of testosterone to estrogen. By inhibiting this enzyme, [16]. Investigation showed improving on testis tissue under effect of silymarin (figure 9-C). Studies on silybin (one of the structural isoforms of silymarin) effects on testicular tissue of laboratory white mice showed that this flavonoid is able to improve the testicular parameters, It has been shown that silymarin increased the concentration of norepinephrine, serotonin and dopamine in certain areas of the brain of laboratory white mice [17]. increasing of gonadotropin hormones from pituitary gland is related to increased release of norepinephrine by silymarin. Norepinephrine by increasing the synthesis of nitric oxide will increase releasing of GnRH from hypothalamus and LH and FSH hormones from pituitary gland [18,19]. Testosterone is an important factor of spermatogenesis process [16]; so, an increase in sperm count and density is detected due to the enhancement of the level of the mentioned hormones by silymarin. The sperm membrane is so sensitive to damages caused by oxygen free radicals [20]. Since free radicals produced in daily reactions of the body are more effective in reducing the count and motility of sperm, one of probable mechanisms of silymarin effects on enhancement of sperm count (figure, 8) may be caused by the antioxidant properties of it. Studies results show that antioxidants in male reproductive system reduce oxidative stress in testis and increase the activity of Leydig cells and so resulting in increase of secretion of testosterone and improvement of spermatogenesis process [21,22]. The additive or synergistic effect of silymarin on the content of glutathione peroxidase has been approved in the various tissues of rats [23,24]. Glutathione peroxidase (GPX) which plays a special role in protection of sperms and ductus deferens in testis [25].

phospholipid hydroperoxide glutathione peroxidase in the spermatozoa of human infertile males. J Biol Reprod. 64: 674–683

4. Agarwa A, Virk G, Ong C, du Plessis S. (2014). Effect of Oxidative Stress on Male Reproduction World J Mens Health. 32(1): 1–17.
5. Gharagozloo P, Aitken RJ (2011). The role of sperm oxidative stress in male infertility and the significance of oral antioxidant therapy. Hum Reprod. 26:1628-40.

6. Choudhary R, Chawala VK, Soni ND, Kumar J, Vyas RK. Oxidative stress and role of antioxidants in male infertility. *Pak J Physiol* 2010;6:54-9
7. Kim, N.C.; Graf, T.N.; Sparacino, C.M.; Wani, M.C.; Wall, M.E. Complete isolation and characterization of silybins and isosilybins from milk thistle (*Silybum marianum*). *Org. Biomol. Chem.* 2003, 1, 1684–1689.3
8. Anthony KP, Saleh M. (2013). Free Radical Scavenging and Antioxidant Activities of Silymarin Components. *Antioxidants*. 2013;2(4):398-407.
9. El-Lakkany NM, Hammam OA, El-Maadawy WH, Badawy AA, Ain - Shoka AA, Ebeid FA. (2012). Antiinflammatory/ anti-fibrotic effects of the hepato protective silymarin and the schistosomicide praziquantel against *Schistosoma mansoni*-induced liver fibrosis. *Parasit Vectors*. 5(9): 21-32
10. Radko L., Cybulski W.(2007). Application of silymarin in human and animal medicine. *Journal of Pre-Clinical and Clinical Research*, 1(1): 022-026.
11. Salih A and Rahim S(2015). The protective effect of aqueous and alcoholic extracts of *Alhagi maurorum* plant against experimentally created liver damage in male rats using CCL4. Tikrit university. Iraq.
12. Farzaneh Eskandari and Hamid Reza Momeni. (2016). Protective effect of silymarin on viability, motility and mitochondrial membrane potential of ram sperm treated with sodium arsenite *Int J Reprod BioMed* Vol. 14. No. 6. pp: 397-402.
13. Huo HZ, Wang B, Liang YK, Bao YY and Gu Y(2011). Hepato protective and antioxidant effect of Licorice extract against CCL4-induced oxidative damage in rats. *Int. J. Mol. Sci.*,12,6529-6543.
14. Iwakami S, Misu H, Takeda T, Sugimori M, Matsugo S, Kaneko S. (2011). Concentration-dependent dual effects of hydrogen peroxide on insulin signal transduction in H4IIEC hepatocytes. *PLoS One*. 6(11):e27401.
15. Claus Hellerbrand, Jörn M. Schattenberg, Philipp Peterburs, Anja Lechner and Reto Brignoli.(2016). The potential of silymarin for the treatment of hepatic disorders Hellerbrand. *Clinical Phytoscience* 2:7 DOI 10.1186/s40816-016-0019-2)
16. Hassanali Abedi, Hossein Kargar Jahromi, Seyed Momammad Amin Hashemi, Hojatollah Karimi Jashni, Zahra Kargar Jahromi and Mohammad Pourahmadi.(2016). The Effect of Silymarin on Spermatogenesis Process in Rats *International Journal of Medical Research & Health Sciences*. 5, 6:146-150
17. Osuchowski MF, Johnson VJ, He Q and Sharma RP. (2004). Alteration in regional brain neurotransmitters by silymarin, a natural antioxidant flavonoid mixture, in BALB/c mice. *Pharm Biol* . 42: 384-9. 17.
18. Selvage DJ, Johnston CA. (2004). Interaction between norepinephrine, oxytocin, and nitric oxide in the stimulation of gonadotropin-releasing hormone release from proestrous rat basal hypothalamus explants. *J Neuroendocrinol*. 16(10):819-24.
19. Całka J. (2006). The role of nitric oxide in the hypothalamic control of LHRH and oxytocin release, sexual behavior and aging of the LHRH and oxytocin neurons. *Folia istochemica Etcytobiological* . 44(1):3-12.)
20. Sikka SC.(2001). Relative impact of oxidative stress on male reproductive function. *Current Medicinal Chemistry*.: 8: 851-862.
21. Glade MJ and Smith K.(2015). Oxidative Stress, Nutritional Antioxidants, and Testosterone Secretion in Men. *Ann NutrDisord&Ther*.;2(1): 1019.
22. Sheweita SA, Tilmisany AM, Al-Sawaf H. (2005). Mechanisms of male infertility: role of antioxidants. *J Curr Drug Metab*; 6(5): 495-501.
23. Valenzuela A.(1989). Selectivity of silymarin on the increase of the glutathione content in different tissues of the rat. *Planta Medica*. 55:420-422.
24. Vargas - Mendoza N, Madrigal - Santillán E, Morales-González Á, Esquivel-Soto J, Esquivel-Chirino C, García - Luna y González - Rubio M, Gayosso - de - Lucio JA, Morales - González JA. (2014). Hepatoprotective effect of silymarin. *World J Hepatol* .6(3): 144-149.
25. Drevet JR. (2006). The antioxidant glutathione peroxidase family and spermatozoa: a complex story. *J MolCell Endocrinol*. 16:70-9.

تأثير السلمارين كمضاد للاكسدة في تحسين الضرر المستحدث بوساطة رباعي كلوريد الكربون على بعض خصائص الجهاز التكاثري للجرذان البيض

ميسر عبدالله احمد¹ ، عزيز خالد حميد² ، حسين محمد طياوي³

¹قسم علوم الحياة ، كلية التربية للعلوم الصرفة ، جامعة تكريت ، تكريت ، العراق

²قسم علوم الحياة ، كلية العلوم ، جامعة تكريت ، تكريت ، العراق

³مديرية تربية صلاح الدين وزارة التربية ، العراق

الملخص

انجزت هذه الدراسة لمعرفة كفاءة السلمارين في منع التأثير الضار للاجهاد التاكسدي الذي يحثه رباعي كلوريد الكربون CCl₄ على بعض خصائص الجهاز التكاثري لذكور الجرذان البيض. عشرون ذكراً من الجرذان البيض قسمت الى اربع مجموعات، خمس حيوانات لكل منها، تضمنت المجموعات المختبرية، مجموعة السيطرة، المجموعة المعاملة بـ (CCl₄ 3m/Kg)، مجموعة السلمارين (150mg/Kg) ومجموعة السلمارين مع CCl₄. كانت مدة الدراسة 35 يوماً. اظهر CCl₄ تأثيراً سميّاً على الجهاز التكاثري من خلال تضرر نسيج الخصية وانخفاض وزن البربخ، عد النطف، الشكل الطبيعي للنطف، حيوية وحركة النطف. اظهر السلمارين فعاليته المضادة للاكسدة لازالة الجذور الحرة الناتجة عن تأثير CCl₄ ولوحظ ارتفاعاً معنوياً في وزن الجسم والخصى، النطف الطبيعية، عد النطف وحيوية النطف بالمقارنة مع المجموعة المعاملة بـ CCl₄.