



Determination of radionuclide concentrations in cigarette tobacco by using Thallium-doped NaI (TI) scintillate technique

Dalia Khawwam Kamil, Asma Ahmad Aziz

Department of Physics, Faculty of Education for Pure Science, Tikrit University, Tikrit, Iraq

DOI: <http://dx.doi.org/10.25130/tjps.25.2020.033>

ARTICLE INFO.

Article history:

-Received: 3 / 9 / 2019

-Accepted: 12 / 11 / 2019

-Available online: / / 2020

Keywords: cigarette tobacco, radioactivity, NaI(Tl) detector, radiological hazard

Corresponding Author:

Name: Asma Ahmad Aziz

E-mail:

Asmaa.ahmed1967@gmail.com

Tel:

Introduction

The environment is defined as the total external conditions that affect the life, growth and survival of the organism. It is also known that the natural environment depends on three main elements: air, water and soil which considered the basics of life. The ecosystem is characterized by a balance between its elements. Therefore, the presence of radiation activity in a certain environment above the allowable limits harmful to humans and living organisms. Generally, the environment in which radiation activity is high is classified as radioactive environment [1].

There are many developments that have occurred and produced polluted environment. Pollution, however, is defined as the introduction of pollutants into the environment that cause disruption in the ecosystem. The pollutants either be exotic substances to the environment or natural substances exceeded the permissible level.

On the other hand, smoking is a widespread phenomenon on a global scale. Moreover, the risk of smoking on humans and the environment are increasing day by day, as tobacco smoke produces large amounts of pollutants. These pollutants are more threatening to human health than that produced by the burning of diesel fuel or the exhaust of cars especially in places such as lactation [2].

ABSTRACT

This study includes the determination of natural and industrial radionuclides concentrations in different samples of cigarette tobacco. The radionuclides are U^{238} , Th^{232} series and K^{40} , as well as industrial radionuclides of Cs^{137} . Radiation risk coefficients were calculated. The results showed that the specific activity rates of Ra^{226} , Bi^{207} , Ac^{228} , K^{40} , and Cs^{137} were within the global limit. The radiological hazard radium equivalent Ra_{eq} , intake dose (D_v), annual effective dose for external and internal exposure ($AEDE_{out}$ and $AEDE_{in}$), health risk external and internal hazard indices data (H_{ex} and H_{in}) and gamma risk radiation were all lower than the allowable global limits.

When organisms, including humans, are exposed to nuclear radiation, a huge chemical changes occur in the tissues of these living organisms, resulting in a significant damage. This damage or chemical change may occur only after a period of time known as the incubation period and may not appear in the person exposed to radiation directly. Otherwise, a change may occur in the genetic composition of future generations [3].

The radioactive materials existed on the surfaces of soils and rocks, water and air, and reach into the human body by means of many ways. First, eating plants and animal's meat, which in its turn absorbed these radioactive materials from the soil. Second, by drinking water and air breathing contaminated with radon gas located in the ground, which is the main source of radioactive that reaches the human body [4].

Studies on alpha or beta as well as gamma radiation have demonstrated the presence of a lot of radioisotopes in all types of cigarettes. Smoking is a type of practice that helps swallowing and inhaling these radioactive isotopes through the mouth and nose.

The inhaled radiotherapy, consequently, reaches the lungs as well as the rest of the upper respiratory tract and the gastrointestinal tract. Smokers are expected to show radioactivity accumulation over time, which

results in increasing the risk of oral, lung and respiratory cancers. whereas, no evidences have been found for the lung cancer among nonsmokers. A team of scientists at Boston University, indicates that the smoking is a change in the structure of lung cells which exposes smokers to increased risk of developing lung cancer [5] [6].

Radon and its dissociation products can enter the human body through both the respiratory system and the digestive system. For the digestive system, it is not dangerous because the presence of food in the stomach, even with a thickness of up to a millimeter, can stop most of the alpha particles resulting from the breakdown of radon and its neonates. The in halation of neonates suspended by air into the respiratory system may adheres to the wall of the lungs. Studies have shown that the dose that the lungs receive as a result of exposure to radon is 2 to 3 times higher than that received by the stomach. While more recent studies revealed that radon is the main cause of lung cancer after smoking [7].

The study aims to measure the concentration of radionuclides for 20 samples of cigarette from different origins. Thallium-doped NaI (TI) scintillator detector is used to determine the specific radiological activity of the natural radioactive elements (uranium U^{238} , thorium Th^{232} , and potassium $K40$). and cesium Cs^{137} , and the radium equivalent and the air intake rate and the internal and external risk rates.

Materials and Methods

A total of 20 samples of cigarette of different types were collected from the most wide spread cigarette in the local markets. The samples were then dried under sunlight for two days, then grinded in an electric mill to obtain a fine and homogeneous powder. For the purpose of measurement, a quantity of 210gm of each sample was placed in a sealed can to prevent contamination. Later, the samples were transferred into measuring container of a suitable volume (not less than 250gm) to measure radionuclides concentrations. Thallium-doped NaI (TI) detector was used to detect the gamma radiation in the studied samples. The Measurement was performed at Baghdad University - Faculty of Science - Physics Department.

The size of the detector system is (3 X 3)inch and operates at 750 Volt, also has a high efficiency of up to 60%. The detector base surrounded with a thick shield of lead to reduce the risk of radiation background in the laboratory.table:1 shows the sample code and type.

Table1: shows the sample code and type

No.	Sample code	Sample type
1	D1	Rothmans
2	D2	Zumerret
3	D3	Milano
4	D4	Pleasure
5	D5	M.M
6	D6	Royale club
7	D7	Dunhil
8	D8	Winston
9	D9	Oscar
10	D10	Kent
11	D11	Miami
12	D12	Summer
13	D13	Prestige
14	D14	Pine
15	D15	Vecory
16	D16	Camil
17	D17	Classic gold
18	D18	Gauloises
19	D19	Aspin
20	D20	Gitanes

For the purpose of calibration of both energy and efficiency of the used detector, a multi-energy source (Co^{60} , Cs^{137} and Am^{242}) was placed in front of the detector for a period of 3600sec. the energy and efficiency values of the radioactive elements used in the calibration process of tobacco models. Genie 2000 program was employed to draws the relationship between calculating relative efficiency of each energy recorded.

Where the calculation of radiation hazard indicators for gama rays

Radium Equivalent Activity (Raeq)

Radiation coefficient is used to ensure uniform distribution of radionuclides, represented by radium Ra^{226} , thorium Th^{232} and potassium K^{40} and measured in units (Bq / kg) and can be calculated by the following equation [8]:

$$A=Cps/(\epsilon_{ff}I_{\gamma}t m)$$

A: activity , cps :count per second ,E:efficiency , I_{γ} : intensity % ,t: time measurement 3600sec , m: sample mass 220gm

$$Raeq = ARa + 1.43A_{Th} + 0.077A_k$$

Whereas ARa , A_{Th} and A_k are the efficiency of Ra^{226} , Th^{232} and k^{40} in Bq / kg, respectively, and the highest value of Raeq should be less than the globally allowed limit of 370Bq / kg.

Dose Rate in Air (D_{γ} Absorbed):

The rate of absorbed dose of Kama rays in the air D_γ at 1m above ground level in nGy/ h can be calculated using the specific activity of radionuclides for radium Ra^{226} for thorium Th^{232} and potassium as in the following equation [8]:

$D_\gamma = 0.462 A_{Ra} + 0.604 A_{Th} + 0.0417 A_k \leq 1$
 (0.462,0.604,0.0417):- are the conversion factors used to calculate the ratio of the air absorbed dose to natural radionuclides (uranium $^{238}_{92}U$, thorium $^{232}_{90}Th$ and potassium $^{40}_{19}k$).

Outdoor Hazard Index:

Radiation coefficient is used to determine the risk of natural Kama radiation and can be calculated from the following equation [9]:

$$Hex = ARa / 370 + ATh / 259 + Ak / 4810 \leq 1$$

Indoor Hazard Index (Hin)

It can be calculated from the following equation [9]:

$$Hin = ARa / 185 + ATh / 259 + Ak / 4810 \leq 1$$

* The external and internal risk index should be less than one [8].

Annual Effective Dose (AEDE) Equivalent:

To calculate the annual effective dose, the conversion factor from the absorbed dose to the active dose should be taken into account. The factor of occupancy is 0.7Sv / Gy as a conversion factor from the air absorbed dose to the annual received dose. The following [8]:

$$AEDE_{in} (mSv/y) = D (nG / h) \times 0.7Sv / Gy \times 0.8 \times 8760 h / y \leq 1$$

The equivalent annual effective external dose is calculated from the following equation [9]:

$$AEDE_{out} (mSv/y) = D (nG / h) \times 0.7Sv / Gy \times 0.2 \times 8760 h / y \leq 1$$

Activity Concentration Index ($I_{\gamma r}$):

Radiation coefficient from which to estimate the risk levels of gama rays associated with natural radionuclides in the samples, can be calculated from the following equation [8]:

$$I_{\gamma r} = ARa / 300 + ATh / 200 + Ak / 3000$$

Results and Discussion

Table: 2 shows the result of radionuclides concentration in tobacco, The acquired results revealed that the specific activity levels of radium (Ra^{226}) were between (33.40-3.64) Bq/kg and (18.52) Bq/kg on average. These results, however, were less than the allow able limit which is 35Bq/kg [10]. While for Bi^{207} , the specific activity levels were below detection limits (B.D.L) which is 35 Bq/kg. The specific activity levels of Ac^{228} were within B.D.L (33.45Bq/kg) and within the global limits ,and for potassium K^{40} were between (4.08 Bq/kg) to (390.31 Bq/kg) and (197.195 Bq/m³) on average [10]. The results appeared less than the global limit (400 Bq/kg). The specific activity levels for Cs^{137} were between B.L.D and (9.56 Bq/kg), where some of the results appeared close to global allowable limits(14.8 Bq/kg) , while Figure 1 illustrates the specific activity of the radionuclides Ra^{224} , Ac^{228} , and K^{40} in the studied tobacco samples.

Table 2: shows the result of radionuclides concentration in tobacco.

Sample ID	Ra-226 Bq/ kg	Bi-207 Bq/ kg	Ac-228 Bq/ kg	k-40 Bq/ kg	Cs-137 Bq/ kg
D1	3.64	10.90	19.07	289.20	B.L.D
D2	15.98	25.05	25.63	180.89	10.45
D3	27.62	2.50	22.85	5.99	14.88
D4	9.99	B.L.D	B.L.D	290.09	B.L.D
D5	19.88	B.L.D	B.L.D	220.70	6.89
D6	27.25	2.32	33.45	131.94	14.02
D7	26.22	14.7	1.09	159.69	B.L.D
D8	19.98	1.55	5.11	14.79	B.L.D
D9	1.505	B.L.D	B.L.D	167.08	12.00
D10	13.00	8.62	B.L.D	149.51	B.L.D
D11	20.40	2.70	10.73	300.45	B.L.D
D12	28.58	13.18	B.L.D	290.20	B.L.D
D13	19.00	1.69	1.0	5.01	12.80
D14	32.97	24.0	7.5	35.02	11.93
D15	18.31	1.29	19.86	4.08	5.60
D16	33.40	30.89	2.0	390.31	B.L.D
D17	18.77	2.09	11.87	25.46	13.27
D18	10.00	9.85	24.16	63.30	B.L.D
D19	15.05	3.0	5.317	162.70	9.56
D20	4.50	1.00	16.46	5.55	B.L.D
W.A	35	35	30	400	14.8

Lower permissible limit: B.L.D

International standard Value : W.A




Fig. 1: illustrates the specific activity of the radionuclides Ra^{224} , Ac^{228} , and K^{40} for the tobacco samples.

The data for radiological risk were as following: Ra_{eq} (radium equivalent) has a values between 14.38-85.25 Bq/kg and 49.81 Bq/kg on average which in its turn lower than the average global allowable limit (370 Bq/kg) [11].The values for air intake dose were between (7.66-38.35 nGyh⁻¹) and the average value was (23.01 nGyh⁻¹) which is much lower than theaverage global allowable limit (84 nGyh⁻¹) [12]. The data for the annual external and internal effective dose (outdoor) were between(0.03-0.05 mSv/y) and an average value of (0.04 mSv/y) which is also less than the allowable global limit (0.07 mSv/y). While the annual effective dose (indoor) ranged between (0.04-0.16 mSv/y) with an average value of (0.1 mSv/y) which is lower than the global limit (0.5 mSv/y) [13]. The calculated external risk index shows a results between (0.386-0.043), whereas for one of the studied samples was bigger than the global limit. The observed average value was (0.2145) lower than the global limit (1). The internal risk index revealed a values between (0.373-0.039) and an average value of (0.206) lower than the global limit (1). The gamma risk indicator presented a values between (0.27-0.07) and an average value of (0.17) which also lower than the global limit of 1Bq/m³[13]. And Table3: Indicates

values for Ra_{eq} , air intake dose, annual affective dose, risk indexes and gamma risk indicator.

Table 3: Indicates values for Ra_{eq} , air intake dose, annual affective dose, risk indexes and gamma risk indicator.

Sample ID	Ra_{eq} (Bq/kg)	$D\gamma$ (nGy/h)	$AEDE_{Outdoor}$ (mSv/y)	$AEDE_{Indoor}$ (mSv/y)	H_{in}	H_{ex}	$I\gamma$
D1	60.44	28.65	0.03	0.14	0.163	0.193	0.22
D2	75.63	34.55	0.04	0.17	0.204	0.272	0.27
D3	60.76	26.81	0.03	0.13	0.960	0.163	0.20
D4	32.33	16.71	0.02	0.08	0.090	0.114	0.13
D5	36.87	18.38	0.02	0.09	0.099	0.154	0.14
D6	85.24	38.35	0.05	0.19	0.229	0.303	0.3
D7	40.1	19.43	0.02	0.09	0.108	0.179	0.15
D8	28.43	12.95	0.02	0.06	0.076	0.130	0.11
D9	14.38	7.66	0.01	0.04	0.039	0.043	0.07
D10	24.51	12.24	0.02	0.06	0.066	0.101	0.09
D11	58.87	28.43	0.03	0.14	0.158	0.213	0.22
D12	50.93	25.3	0.03	0.12	0.137	0.215	0.19
D13	20.82	9.60	0.01	0.05	0.056	0.108	0.07
D14	46.39	21.22	0.03	0.1	0.125	0.215	0.16
D15	47.01	20.62	0.03	0.1	0.127	0.177	0.16
D16	66.31	32.92	0.04	0.16	0.179	0.269	0.25
D17	37.7	16.9	0.02	0.08	0.149	0.201	0.13
D18	49.42	21.85	0.03	0.11	0.133	0.160	0.17
D19	35.18	16.94	0.02	0.08	0.096	0.136	0.12
D20	28.47	12.25	0.02	0.06	0.373	0.386	0.10
W.A	370	84	0.07	0.5	1	1	1

References

- [1] Atared, Kh. and Shaimaa, F.(2013). Reality of Environment and Energy Statistics in Iraq, Ministry of Planning, Central Statistical Organization.
- [2] Inferniti d. Giovanni. (2002). Tobacco Control Authority. the British Health and Smoking Authority. Journal of Science and Health (Scientific Discovery and Research of Tobacco),
- [3] Abdul-Mahmoud and Abdul-Karim. (2002). Environmental and Health Effect of Radioactive Materials. Decisions of the Scientific Conference on the Impact of the Use of DU Weapons in Man and Environment in Iraq, Part I , p. 15 .
- [4] Al-Qamoudi Salem.(2007). Scientific Culture, Humanities Branch, Cairo University, Al-Mustafa Electronic Library, Cairo, Egypt , P: 33-44.
- [5] Hampson, SE. et al.(1998) . Risk-Anal, (For radon and eating cigarettes together) ,**18(3)** ,pp (50-343) .
- [6] global status report WHO. (1997).Tobacco or health .
- [7] Ali, Gh. (1999). Radon and its impact on the environment and man. Publishing House of Corn and Development, **11(3)**, p: 459.
- [8] Hadeel , Gh. (2015). Finding a quasi-experimental relationship for measurements of radon emitted from building materials inside residential houses. College of Education for Pure Sciences, Ibn Al-Haytham, University of Baghdad.
- [9] Cartwright B. And Shirk E. (1978). A nuclear Track Recording Polymer of Unique Sensitivity and Resolution, Nucl. Inset and Meth, **15(3)**, pp. (457-460).
- [10] Report of the Ministry of Environment . (2012). Center for Radiation Protection, Department of Radiation Control.
- [11] Al-Nafai, Murtada and Al-Mayahi, Basem. (2008). Measurement of Radioactivity of Surface Water and Sediment Models by gamma Rays Spectrum in Some Hilla Areas, Basrah Research Journal (Operations), **34(2)**,p:6 .
- [12] Tawfiq, N. ; Mansour, H., and Karim, M. S. (2015)." Measurement of Radon Gas Concentrations in Tap Water for Baghdad Governorate by Using Nuclear Track Detector (CR-39)", International Journal of Physics, **3(6)**, (233-238) .
- [13] Al-Ubaidi, A. M. (2015). Environmental Radioactivity of Al-Rashidiyah Site –Baghdad. Ph.D. Thesis, University of Baghdad, College of Science for Women.

أيجاد تراكيز النويدات المشعة في تبغ السكائر باستخدام تقنية الكاشف الوميضي

ايوديد الصوديوم NaI(Tl)

داليه خوام كامل ، اسماء احمد عزيز

قسم الفيزياء ، كلية التربية للعلوم الصرفة ، جامعة تكريت ، تكريت ، العراق

الملخص

تتضمن هذه الدراسة ايجاد تراكيز النويدات المشعة الطبيعية والمتمثلة بسلسلة اليورانيوم U238 وسلسلة الثوريوم Th232 والبوتاسيوم، K40 والنويدات المشعة الصناعية المتمثلة بالسيزيوم، Cs137 لعشرين عينة من تبغ السكائر المختلفة المنشأ، باستخدام تقنية مطيافيه اشعة كما كاشف ايوديد الصوديوم المطعم بالتاليوم، ((TI) NaI) وحساب معاملات الخطورة الاشعاعية في هذه النماذج وظهرت النتائج ان معدلات الفعالية النوعية للراديويم Ra226 والبزموت Bi 207 والاكثينيوم Ac228 والبوتاسيوم K40-والسيزيوم Cs137-في جميع النماذج ضمن الحدود المقبولة عالمياً. ومعدلات مؤثرات الخطورة الاشعاعية المتمثلة بفعالية الراديويم المكافئة (Raeq) ومعدل الجرعة الممتصة في الهواء (Dy) والجرعة الفعالة السنوية للتعرض الخارجي (AEDEOUT) والجرعة الفعالة السنوية للتعرض الداخلي (ADEAin) ودليل الخطورة الخارجي (Hex) ودليل الخطورة الداخلي (Hin) ودليل الخطورة لأشعة كما كانت جميعها اقل من الحدود المقبولة بها عالمياً .