

Cardiovascular reflexes in response to head down tilting in normal subjects

*Dr.Yesar M. H. Al-Shamma Prof. of Physiology M.B.Ch.B, Ph. D. (U.K), Dr. Abdulameer Abdulbary Al_Amery M.B.Ch.B, F.I.C.M.S(med), F.I.C.M.S (card) and Dr. Ahmed Bader Abd_Alwahid M.B.Ch.B.
*Head of physiology department, college of medicine, Kufa University.

انعكاسات القلب و الأوعية الدموية استجابة لميلان جسم الإنسان إلى الأسفل عند الأشخاص الطبيعيين

الخلاصة:

الهدف من هذه البحث هو دراسة انعكاسات القلب و الأوعية الدموية استجابة لميلان جسم الإنسان إلى الأسفل بدرجة ٢٠°.

أجريت هذه الدراسة على ٦٠ شخص من الأصحاء (٤٠ ذكر و ٢٠ أنثى) لغرض دراسة تأثير ميلان جسم الإنسان إلى الأسفل بدرجة ٢٠° بواسطة استخدام الطرق غير المباشرة لقياس متغيرات دايينميكية القلب والأوعية الدموية حيث استعمل جهاز الضغط الزئبقي لقياس ضغط الدم وجهاز أوكسي ميتر لقياس النبض (HR) وجهاز الايكو دوبلر لقياس كمية الدم التي يضخها القلب في النبضة الواحدة (SV)، كمية الدم التي يضخها القلب في الدقيقة الواحدة (CO)، سرعة الوقت المتكاملة (VTI).

تم تقسيم الأشخاص إلى فئتين عمريتين: الفئة العمرية الأولى (٢٠-٤٠ سنة) مكونه من ٣٢ ذكر و ٨ إناث والفئة العمرية الثانية (٤١-٦٠ سنة) مكونه من ٨ ذكور و ١٢ أنثى.

وكانت نتائج البحث تشير إلى أن التغيرات في الفئة العمرية الأولى تمثلت بزيادة معنوية في مقدار ما يضخه القلب في النبضة الواحدة (SV) وما يضخه في الدقيقة الواحدة (CO) وسرعة الوقت المتكاملة (VTI) بينما ان هناك انخفاض معنوي في عدد نبضات القلب (HR) و المقاومة المحيطية الكلية للأوعية الدموية (TPVR) وكل هذه التغيرات حدثت بسبب ميلان الرأس بدرجة ٢٠° إلى الأسفل. بينما أشارت النتائج إلى عدم وجود أي تغيير معنوي في ضغط الدم الانقباضي (SBP) و ضغط الدم الانبساطي (DBP) ومعدل ضغط الدم (MBP).

في الفئة العمرية الثانية أشارت التغيرات إلى وجود زيادة معنوية في مقدار ما يضخه القلب في النبضة الواحدة (SV) وما يضخه في الدقيقة الواحدة (CO) وسرعة الوقت المتكاملة (VTI) وضغط الدم الانقباضي (SBP) وضغط الدم الانبساطي (DBP) ومعدل ضغط الدم (MBP) استجابة إلى ميلان الرأس بدرجة ٢٠° إلى الأسفل بينما لم تحصل أي تغييرات معنوية في عدد نبضات القلب (HR) و المقاومة المحيطية الكلية للأوعية الدموية (TPVR).

Abstract

Objectives: The aim of this study is to investigate the changes in hemodynamic variables in response to head down tilting.

Methods: This study was carried out on 60 normal healthy subjects, 40 males and 20 females in order to study the cardiovascular reflexes in response to 20° head down tilting by using non-invasive techniques including mercury sphygmomanometer for measurement of blood pressure , pulse oximeter for measurement of heart rate, Doppler echocardiography for estimation of velocity time integral (VTI) ,stroke volume (SV) and cardiac output (CO).The subjects were divided into two groups: the first group includes subjects with age between 20-40 years and this group consist of 32 males and 8 females.The second group includes the subjects with age between 41-60 years and this group consist of 8 males and 12 females.

Results: In the first group there is significant increase in velocity time integral (VTI) , stroke volume (SV) and cardiac output (CO) and significant decrease in heart rate (HR) and total peripheral vascular resistance (TPVR) in response to 20°head down tilting. While there are no significant differences in systolic, diastolic and mean blood pressure.

In the second group there is significant increase in velocity time integral (VTI), stroke volume (SV), cardiac output (CO), systolic, diastolic and mean blood pressure in response to 20° head down tilting. While there are no significant differences in heart rate (HR) and total peripheral vascular resistance (TPVR).

Keywords: head down tilt, cardiac output.

Introduction

There have been many researches concerning the assessment of cardiovascular reflexes in normal subjects to test the integrity of the cardiovascular performance such as head up tilting at different angles of tilt (1,2,3), lower body negative pressure (LBNP) which simulate head up tilting but subject remain in supine position(4,5), neck suction to stimulate carotid baroreceptors(6), valsalva maneuver(7), supine and upright exercise(8) or immersion in water (9) in which there is shifting of blood from the lower part of the body to the thoracic cavity so that it will cause loading of cardiopulmonary low pressure and arterial high pressure receptors(10) but there were a very little researches concerning head down tilt for the assessment of cardiovascular reflexes. Head down tilt (HDT) has been widely accepted as an effective simulated weightlessness model in humans by inducing a fluid shift from the lower body towards the thoracic compartment and result in a loading of cardiopulmonary pressure and arterial high pressure receptors. So the aim of this study is to investigate the changes in hemodynamic variables in response to head down tilting.

Material and Method:

1. Subjects

This study was carried out on 60 normal subjects 40 men and 20 women. They are apparently normal healthy subjects. They are examined by consultant physician and they have no symptoms of any diseases with normal cardiovascular status which evaluated by physical examination, also chest X-ray, electrocardiogram and echocardiography, was consistently within normal range. They divided into two age groups the first age group (20-40 years) and it consist of 32 males and 8 females while the second group age group (41-60 years) and it consist of 8 males and 12 females.

2. Materials: The study is performed by using Transthoracic 2-D and Doppler echocardiography techniques G.E Vivid S5:10.3.0 build 114 (General Electric) with a probe frequency 1.7-3.4 MHz , Mercury sphygmomanometer, pulse oximeter and electrically motorized tilt table.

3. Method: Study protocol consists of two phases:

Phase -1 (supine control state): The subject first placed in supine position for 5-15 minutes for a steady state. The steady state mean that heart rate changing by less than 2-3 beats in consecutive minutes (1,11). Then measurements of blood pressure by sphygmomanometer and heart rate by pulse oximeter were made.

Cardiac output was measured by echocardiography. Left ventricular outlet tract (LVOT) diameter was measured at the level of the aortic annulus from the parasternal long-axis view (12) in systole (13) using 2-D echocardiography. The diameter of the aortic orifice was measured at the attachment of the valve cusps and it was determined only at baseline, and this value was assumed to be constant and used during second phase of CO calculations. LVOT area was calculated as π (LVOT diameter/2)². The pulsed Doppler sample volume was positioned in the middle of the left ventricular outflow

immediately proximal to the aortic valve leaflets from the apical 5-chamber view (14). The outline contour of the velocity curve was traced and the velocity time integral was derived automatically. Doppler-determined CO was calculated as the product of heart rate \times the Doppler time velocity integral \times cross-sectional area of the aortic annulus,

which was considered as a circle. So we use the following equations:

$$SV = 0.785 \times D2 \times VTI \quad (15,16)$$

$$CO = SV \times HR$$

Total Peripheral vascular resistance (TPVR) is calculated from the equation: $TPVR = MBP / CO$

Phase -2 :The subject was tilted by 20° head down tilting and after a steady state the measurement of all hemodynamic variables was repeated.



Tilt table (20° head down tilting).

4. Statistical Analysis:

Paired t-test was used to determine the differences in hemodynamic variables during supine position and 20 °head down tilting.

Results:

Data collected and represented in table (1).

Table (1): comparison between hemodynamic variables in age group 1 and 2 in supine position : systolic blood pressure(SBP), diastolic blood pressure(DBP) mean blood pressure (MBP), heart rate (HR) , velocity time integral (VTI) stroke volume (SV), cardiac output (CO) and total peripheral vascular resistance(TPVR).

hemodynamic variables	Group 1		Group 2	
	Supine position	20° head down tilt	Supine position	20° head down tilt
SBP(mm.Hg)	117.3 ± 8.18	118.1 ± 7.79	125 ± 9.92	132.5 ± 11
DBP(mm.Hg)	74.1 ± 6.91	74.6 ± 7.44	78.8 ± 5.97	84.75 ± 6.86
MBP(mm.Hg)	88.5 ± 5.91	89 ± 6.27	94.2 ± 6.49	100.7 ± 7.25
HR(bpm)	75.9 ± 5.99	72.4 ± 6.39	73.1 ± 6.6	72.35 ± 6.72
VTI(cm)	22 ± 2.11	26.8 ± 2.48	20.2 ± 3.98	23 ± 4.29
SV(ml)	71.2 ± 10.1	86.7 ± 12.63	65 ± 8.31	75.3 ± 10.34
CO(L/m)	5.4 ± 0.86	6.3 ± 1.03	4.7 ± 0.48	5.3 ± 0.5
TPVR	16.8 ± 2.94	14.6 ± 2.64	20.2 ± 2.53	19.1 ± 2.51

1. Cardiovascular reflexes in response to 20° head down tilting in group1 (20 - 40 years) indicated that VTI, SV and CO were significantly higher in response to 20° head down tilting than that of supine position . Whereas the HR, TPVR were significantly lower in 20° head down tilting and there were no significant changes in SBP,DBP and MBP.

2.Cardiovascular reflexes in response to 20° head down tilting in group2 (41- 60 years): indicated that the SV, CO, VTI, SBP, DBP and MAP were significantly higher in response to 20° head down tilting than that of supine position, where as there are no significant differences in HR and TPVR.

Discussion:

1. Cardiovascular reflexes in response to 20° head down tilting in group1 indicated that VTI, SV and CO were significantly higher in response to 20° head down tilting than that of supine position which is mostly because in head down tilting blood and fluid are redistributed from the caudal to the cephalic portions of the body, leading to an increased venous return. This increase in venous return induces distension of the heart and adjacent vessels. There is an intrinsic ability of the heart to adapt to increasing volumes of inflowing blood called the Frank-Starling mechanism of the heart so the greater the heart muscle is stretched the greater the quantity of blood pumped into the aorta.(17)

The HR and TPVR were significantly decreased in response to head down tilting. This result agree with other researchers who reported that the cardiopulmonary blood volume and the filling pressure of the heart play a key role in the regulation of the cardiovascular function and the body fluid volume homeostasis. Experimental studies have shown that changes in cardiac filling originate reflexes through the activation of receptors in the heart. The mechanisms probably involve a combined stimulation of cardiopulmonary low-pressure and arterial high-pressure receptors. In humans relative contribution of each type of receptors, low vs. high cannot be ascertained (18,19, 20). When activated, these receptors induce powerful depressor reflexes with bradycardia and sympathetic inhibition. In this study there was no change in SBP, DBP and MBP which is in agreement with other researchers who reported that the increase in stroke volume lead to an increase in arterial blood pressure which returned to approximately the same level for an individual within 18 sec. the initial increase in blood pressure is due to primary hydrostatic shift and blood pressure returned to approximately the same level for an individual by the secondary response (depressor reflexes)

2. Cardiovascular reflexes in response to 20° head down tilting in group 2 indicated that VTI, SV and CO were significantly higher in response to 20° head down tilting than that of supine position which is mostly due to increase venous return to the heart similar to that at group 1.The HR and TPVR were not significantly changed while SBP, DBP

and MBP were significantly higher in response to 20° head down tilting than that of supine position. Normally increase in age is associated with gradual dysfunction of regulatory function (21,22). However, the mechanism of this important phenomenon is unknown. It has been suggested that age-related dysfunction of regulatory mechanism might result from:

A. Altered autonomic reflex function:

The structure and function of autonomic neurons appear to be altered with aging. These structural changes are associated with dysfunction of body cardiovascular maintenance, blood pressure, water and electrolyte distribution, and energy metabolism. Several studies indicate that basal sympathetic activity increases in some elderly individuals, and the increase may be associated with dysregulation of the ability of the sympathetic nervous system to respond to a variety of challenges. (23)

B. Reduce vascular response.

Due to the cholesterol deposits of atherosclerosis and endothelial dysfunction Overall changes in vasomotility with aging are characterized by (i) decreased relaxation (ii) increased constriction, and the resulting imbalance of vascular tone leading to increased vasoconstriction Simultaneously , the vasoconstrictor action of adrenergic stimulation may be increased (24).

C. The reduction either in the afferent or central integrating parts of the baroreflex pathway (25).

Conclusion:

The cardiovascular regulatory mechanism facilitate the adaptation of cardiovascular system to any volume or pressure overload on the cardiovascular system. Head down tilting results in loading of the arterial high pressure and cardiopulmonary low pressure receptors leading to a depressor reflex (via the vasomotor center) in a manner opposite to that of head up tilting. There is age related dysfunction of regulatory function of the cardiovascular system.

References:

1. **Hainsworth R. and Al-Shamma Y.M.H.(1988).** Cardiovascular responses to upright tilting in healthy subjects, clinical sciences,74;pp 17-22
2. **Al-Shamma, Y. M., Khawaja S.A.M and Alabidy (2002).** Effect of upright tilting on cardiovascular reflexes using echocardiographic method for estimating cardiac output. Kufa med. J. (5).2.p85-96.
3. **Murrell C.J.,Cotter J.D., George K., Shave R., Wilson L., Thomas K. , Williams M.J. Ainslie P.N. (2011).** Cardiorespiratory and cerebrovascular responses to head-up tilt I: Influence of age and training status Experimental Gerontology 46 (2011) 9–17.
4. **Al-Shamma, Y. M. H. and Hainsworth .R. (1985).**The cardiovascular responses to upright tilting and lower body negative pressure in man.proceeding of the physiological society, J of physiology 369,188.
5. **Wallace JP, Trail GT, Franke WD.(2010).** LBNP tolerance analyzed retrospectively using a structural equation model. Aviat Space Environ Med. 2010 Apr;81(4):363-8
6. **Hainsworth R. and Al-Shamma Y.M.H.(1988).** Cardiovascular responses to stimulation of carotid baroreceptors in healthy subjects, clinical sciences,75;pp 159-165.
7. **Al-Shamma Y.M.H., ALmumin A.S. and Abood A,K (2007).** Effect of valsalva maneuver on cardiovascular reflexes. QMJ(3):3-21.
8. **Al-Shamma Y.M.H., Zwain A. M. , and Kasim A.L. (2008).** Cardiovascular reflexes in response to supine and upright exercise. Kufa Med.J. (11) :87-107.

9. **Shiraishi M, Schou M, Gybel M, Christensen N J and Norsk P.(2002)** Comparison of acute cardiovascular responses to water immersion and head-down tilt in humans . J Appl Physiol 92: 264–268.
10. **Al-Shamma Y.M.H. and Al-Naily F.H.A.(2005).**The effect of immersion in water on cardiopulmonary response. Kufa Med. J. 8:(1),(134-146).
11. **Al-Shamma, Y. M.,Khawaja S.A.M and Alabidy (2002).** Effect of upright tilting on cardiovascular reflexes using echocardiographic method for estimating cardiac output. Kufa med. J. (5).2.p85-96.
12. **Ihlen H, Myhre E, Amlie JP, Forfang K, Larsen S.(1985).** Changes in left ventricular stroke volume measured by Doppler echocardiography. I& Heart J:54:378-383.
13. **Kitabatake A, Inoue M, Asao M, Ito H, Masuyama T, Tanouchi J, Morita T, Hori M, Yoshima H, Ohnishi K, (1984).** Noninvasive evaluation of the ratio of pulmonary to systemic flow in atrial septal defect by duplex Doppler echocardiography. Circulation;69:73–79.
14. **Lewis JF, Kuo LC, Nelson JG, Limacher MC, tinones MA (1984).** Pulsed Doppler echocardiographic determination of stroke volume and cardiac output: clinical validation of two new methods using the apical window. Circulation;70: 425-431.
15. **Solomon (2007).** Echocardiographic Instrumentation and Principles of Doppler Echocardiography In Essential Echocardiography A Practical_Handbook part 1 p3-13.
16. **Adams, Ahmed A. Alsaileek, Christopher H. Cabell, John B. Chambers (2010)** The principle of flow assessment. In springer echocardiography 2010 chapter 3 page 50-51.
17. **Guyton and hall (2007).** Overview of the Circulation; Medical Physics of Pressure, Flow, and Resistance. In textbook of medical physiology ch 14 unit IV p166.
18. **Eckberg, D. L., and P. Sleight (1992).** History, reflex interactions. In: Human Baroreflexes in Health and Disease. Oxford, UK: Clarendon,1992, p. 13-15, 153–215.
19. **Rowell, L. B. (1993)** Neural-humoral adjustments to orthostasis and long-term control. In:Human Cardiovascular Control. NewYork: Oxford Univ. Press, 1993.
20. **Westerhof B.E.,Gisolf J.,Karemaker J.M., Wesseling K. H., Secher N.H. and van Lieshout J.J (2006).** Time course analysis of baroreflex sensitivity during postural stress. Am J Physiol Heart Circ Physiol 291: H2864–H2874.
21. **Gaffney F.A. , Nixon J.V., Karlsson ES, Campbell W, Dowdey A.B.C and Blomqvist C.G.(1985) .** Cardiovascular deconditioning produced by 20 hours of bedrest with head-down tilt (–5°) in middle-aged healthy men Am J Cardiology VOL 56 Issue Pages 634-638.
22. **Vargas E , Lye M, Faragher B, Goddard C, Moser B, and Davies I (1986).** cardiovascular haemodynamics and the response of vasopressin, aldosterone, plasma renin activity and plasma catecholamines to head-up tilt in young and old healthy subjects . Age ageing 15 (1): 17-28.
23. **Kuchel GA, Hof PR, (2004).** Autonomic Nervous System in Old Age. Vol. 33. New York: Karger.
24. **Timiras and Navazio(2007).** Aging-Related Changes in Cardiac Output In:Physiological Basis of Aging and Geriatrics fourth edition ch 20 p 340.
25. **O'mahony D, Green A', Bennett C, Sinclair A J (1998).** efferent brainstem autonomic function is unchanged in age- related loss of arterial elated loss of arterial baroreceptor sensitivity age_ageing-p34-a-p34.