

SURVEY OF THE MARINE FISHES IN IRAQ

Abbas J. Al-Faisal* and Falah M. Mutlak
Marine Science Centre, University of Basrah, Basrah, Iraq
*Corresponding author: abbasjsm71@yahoo.com

Received Date: 19 April 2018

Accepted Date: 28 August 2018

ABSTRACT

A survey of fish species in the Iraqi marine waters was carried out for the period from November 2014 to March 2018. The list included 214 species representing 75 families.

The family Carangidae dominated the marine fishes in Iraq, which was represented by 24 species, followed by Haemulidae with 11 species, and then Serranidae and Sparidae with nine species for each, while 34 families contained a single species only.

Key words: Checklist, fish species, Iraq, marine waters, survey.

INTRODUCTION

The Arabian Gulf is a semi-enclosed shallow sea located in the subtropical climate. The Gulf is considered biologically impoverished, due to its environmental characteristics, in addition to its young age (Randall, 1995; Jabado *et al.*, 2015). The environments of Iraqi marine waters are fairly different in comparison with other parts of the Arabian Gulf; the Shatt Al-Arab River, which is formed by the confluence of the Euphrates and Tigris rivers, is considered as a major fresh water discharge into the northwest Arabian Gulf. The potential sources of nutrients and organics in the northwestern Arabian Gulf are Shatt Al-Arab river and its associated marshes (Al-Yamani, 2008; Al-Mudaffar and Mahdi, 2014). The Shatt Al-Arab estuary plays an important role as feeding, nursery and protective grounds for many fish species (Hussain *et al.*, 1999).

The fish species recorded in the Arabian Gulf were derived from the Indian Ocean, besides demonstrating that the Indian Ocean species cover the entire gulf and reaching even its most northern range (Hussain and Naama, 1989). The fish fauna of the Arabian Gulf have been identified by several systematic works (Blegvad and Løppenthin, 1944; Kuronuma and Abe, 1986; Randall, 1995; Assadi and Dehgani, 1997; Carpenter *et al.*, 1997; Bishop, 2003; Iwatsuki *et al.*, 2013).

The previous studies in the Iraqi marine waters could be divided into two parts, the first dealing with species composition and seasonal fluctuations either in the inland Iraqi marine waters, like in Khor Al-Zubair (Ali, 1985; Hussain *et al.*, 1988; Hussain and Naama, 1989; Ali and Hussain, 1990; Hussain *et al.*, 1994) and Khor Abdullah (Younis, 1990; Hussain and Younis, 1997), and the second dealing with classification of fishes or checklist in the regional Iraqi waters (Khalaf, 1961; Mahdi, 1962; Mahdi and Georg, 1969; Al-Daham, 1982; Mohamed *et al.*, 2001).

Survey of the marine fishes

The aim of the present study is to update the list of marine fish species based on a survey in them from the Iraqi marine waters for the period from November 2014 till March 2018.

MATERIALS AND METHODS

Fish samples were collected from the Iraqi marine waters (29° 46' 50"N 48° 39' 46" E to 29° 78' 83"N 48° 75' 78"E) (Map 1), northwest of the Arabian Gulf during the period from November 2014 to March 2018, by using a fishing boat supplemented by trawl net; also the fish samples were collected from commercial fisheries, through coordination and cooperation with fishermen of the region, which used different fishing nets, benthic trawl nets, fixed gill nets, gill nets (mesh 78×78 mm, 67×67 mm, 57×57 mm, 48×48 mm, 42×42 mm, 33×33 mm, 30×30 mm and 28×28 mm), traps with different circular base (2 m, 2.5 m and 4 m) and fish hooks, as well from commercial fishery of Al-Nasr society for the fishing and marketing of marine fish in Al-Faw district, 100 km south of Basrah city.

Fish species were identified depending on Kuronuma and Abe (1986); Carpenter *et al.* (1997) and Froese and Pauly (2018). The fish images were taken by Canon SX600 HS. The arrangement of higher taxa (Classes, Orders and Families) of fishes followed Nelson (2006).



Map (1): Fishing area in the Iraqi marine waters.

Abbas J. Al-Faisal and Falah M. Mutlak

RESULTS AND DISCUSSION

A total of 214 species belonging to 75 families and 21 orders were recorded from the Iraqi marine waters (Tab. 1), (Pl. 1 and 2 showed some species). The number of cartilaginous fish species (Class: Elasmobranchii) was 16 species and the number of bony fish species (Class: Actinopterygii) was 198 species; the order Perciform dominated the ichthyofauna of the Iraqi marine waters, which is represented by 36 families and 132 species. A total of 34 families contained a single species only, 16 families contained two species and seven families contained three species, the remaining families were represented by more than three species. The most successful fish families are Carangidae (11.21%) which contains 24 species, followed by Haemulidae (5.14%) with 11 species, while Serranidae and Sparidae (4.20%) in the third place each is represented by nine species per each, then Sciaenidae and Lutjanidae (3.73%) with eight species each.

The previous studies have developed a database for the identification of marine fish species in Iraqi marine waters, but they have been based on fisheries or have been short term studies. The most prominent study about ichthyofauna of Iraqi marine waters was carried out by Mohamed *et al.* (2001) for the period 1995-1999, by using two fishing boats; they recorded 116 species belonging to 58 families. The present study is the next study of Mohamed *et al.* (2001) about ichthyofauna of the Iraqi marine waters; the number of species reached 214 which belong to 75 families; the increase in fish species in the present study could be due to using different fishing gears at different depths by the fishermen. On the other hand, this increase in the number of species may be due to modifications in Iraqi marine habitat outside its control. The quantity of freshwater discharges into the northwest Arabian Gulf varies from time to time, as the construction of dams on the Tigris and Euphrates rivers in Turkey and decreased the discharge of Karun river; the various species were distributed in the northwestern Arabian Gulf as could be expected in view of salinity variations, different topography and muddy substratum (Mohamed *et al.*, 2001).

The Perciform is considered to be the largest order within the teleost fishes (Nelson, 2006), this order dominates the ichthyofauna in northwest of the Arabian Gulf (Khalaf, 1961; Mahdi, and Georg, 1969; Al-Daham, 1982; Kuronuma, and Abe, 1986; Bishop, 2003), and which is represented by 36 families and 132 species in the present study.

The most successful fish family in the Iraqi marine waters was Carangidae, represented by 24 species, this number is likely to increase, where were occurrence 32 species in Kuwait (Bishop, 2003) and 40 species in the Arabian Gulf (Carpenter *et al.*, 1997). This applies to the rest of families as Haemulidae, Serranidae, Sparidae and Sciaenidae.

This study provided an updating list of the ichthyofauna of Iraqi marine waters as to be a base for various studies in the future.

Survey of the marine fishes

Table (1): List of fish species from the Iraqi marine waters.

Order	Family	Species
Orectolobiformes	Hemiscylliidae	<i>Chiloscyllium arabicum</i> Gubanov, 1980
	Rhincodontidae	<i>Rhincodon typus</i> Smith, 1828
Carcharhiniformes	Carcharhinidae	<i>Carcharhinus leucas</i> (Müller & Henle, 1839)
		<i>Rhizoprionodon acutus</i> (Rüppell, 1837)
	Sphyrnidae	<i>Sphyrna mokarran</i> (Rüppell, 1837)
Rhinopristiformes	Pristidae	<i>Pristis zijsron</i> Bleeker, 1851
	Rhinobatidae	<i>Glaucostegus granulatus</i> (Cuvier, 1829)
Myliobatiformes	Dasyatidae	<i>Brevitrygon imbricata</i> (Bloch & Schneider, 1801)
		<i>Brevitrygon walga</i> (Müller & Henle, 1841)
		<i>Himantura uarnak</i> (Gmelin, 1789)
		<i>Maculabatis gerrardi</i> (Gray, 1851)
		<i>Pastinachus sephen</i> (Forsskål, 1775)
		<i>Pateobatis bleekeri</i> (Blyth, 1860)
	Gymnuridae	<i>Gymnura poecilura</i> (Shaw, 1804)
	Myliobatidae	<i>Aetobatus flagellum</i> (Bloch & Schneider, 1801)
		<i>Aetobatus narinari</i> (Euphrasen, 1790)
Anguilliformes	Muraenesocidae	<i>Muraenesox cinereus</i> (Forsskål, 1775)
Clupeiformes	Chirocentridae	<i>Chirocentrus dorab</i> (Forsskål, 1775)
		<i>Chirocentrus nudus</i> Swainson, 1839
	Clupeidae	<i>Anodontostoma chacunda</i> (Hamilton, 1822)
		<i>Nematalosa nasus</i> (Bloch, 1795)
		<i>Sardinella albella</i> (Valenciennes, 1847)
		<i>Sardinella gibbosa</i> (Bleeker, 1849)
		<i>Sardinella longiceps</i> Valenciennes, 1847
		<i>Tenualosa ilisha</i> (Hamilton, 1822)
	Dussumeriidae	<i>Dussumeria acuta</i> Valenciennes, 1847
	Engraulidae	<i>Thryssa dussumieri</i> (Valenciennes, 1848)
		<i>Thryssa hamiltonii</i> Gray, 1835
		<i>Thryssa</i> sp.
		<i>Thryssa vitrirostris</i> (Gilchrist & Thompson, 1908)
		<i>Thryssa whiteheadi</i> Wongratana, 1983
	Pristigasteridae	<i>Ilisha compressa</i> Randall, 1994
		<i>Ilisha melastoma</i> (Bloch & Schneider, 1801)
Gonorynchiformes	Chanidae	<i>Chanos chanos</i> (Forsskål, 1775)
Siluriformes	Ariidae	<i>Plicofollis dussumieri</i> (Valenciennes, 1840)
		<i>Netuma bilineata</i> (Valenciennes, 1840)
		<i>Netuma thalassina</i> (Rüppell, 1837)
	Plotosidae	<i>Plotosus lineatus</i> (Thunberg, 1787)
Aulopiformes	Synodontidae	<i>Saurida tumbil</i> (Bloch, 1795)
		<i>Saurida undosquamis</i> (Richardson, 1848)
Gadiformes	Bregmacerotidae	<i>Bregmaceros neclabanus</i> Whitley, 1941
Ophidiiformes	Ophidiidae	<i>Neobythites steatiticus</i> Alcock, 1894
Batrachoidiformes	Batrachoididae	<i>Allenbatrachus grunniens</i> (Linnaeus, 1758)
Mugiliformes	Mugilidae	<i>Liza carinata</i> (Valenciennes, 1836)
		<i>Liza klunzingeri</i> (Day, 1888)
		<i>Planiliza subviridis</i> (Valenciennes, 1836)

		<i>Mugil cephalus</i> Linnaeus, 1758
		<i>Valamugil speigleri</i> (Bleeker, 1858)
Beloniformes	Belonidae	<i>Ablennes hians</i> (Valenciennes, 1846)
		<i>Strongylura strongylura</i> (van Hasselt, 1823)
		<i>Tylosurus crocodilus</i> (Péron & Lesueur, 1821)
	Hemiramphidae	<i>Hemiramphus marginatus</i> (Forsskal, 1775)
		<i>Hyporhamphus limbatus</i> (Valenciennes, 1846)
		<i>Rhynchorhamphus georgii</i> (Valenciennes, 1846)
Beryciformes	Monocentridae	<i>Monocentris japonica</i> (Houttuyn, 1782)
Gasterosteiformes	Pigasidae	<i>Pegasus volitans</i> Linnaeus, 1758
Syngnathiformes	Syngnathidae	<i>Hippocampus kuda</i> Bleeker, 1852
Scorpaeniformes	Platycephalidae	<i>Platycephalus indicus</i> (Linnaeus, 1758)
		<i>Grammolites scaber</i> (Linnaeus, 1758)
		<i>Pseudosynanceia melanostigma</i> Day, 1875
		<i>Minous monodactylus</i> (Bloch & Schneider, 1801)
		<i>Synanceia nana</i> Eschmeyer & Rama-Rao, 1973
Perciformes	Acanthuridae	<i>Acanthurus sohal</i> (Forsskal, 1775)
	Apogonidae	<i>Apogonichthyoides taeniatus</i> (Cuvier, 1828)
	Ariommatidae	<i>Ariomma indica</i> (Day, 1870)
	Carangidae	<i>Alectis ciliaris</i> (Bloch, 1787)
		<i>Alectis indica</i> (Rüppell, 1830)
		<i>Alepes djedaba</i> (Forsskal, 1775)
		<i>Alepes kleinii</i> (Bloch, 1793)
		<i>Alepes melanoptera</i> (Swainson, 1839)
		<i>Alepes vari</i> (Cuvier, 1833)
		<i>Atropus atropus</i> (Bloch & Schneider, 1801)
		<i>Atule mate</i> (Cuvier, 1833)
		<i>Carangoides armatus</i> (Rüppell, 1830)
		<i>Carangoides bajad</i> (Forsskal, 1775)
		<i>Carangoides chrysophrys</i> (Cuvier, 1833)
		<i>Carangoides malabaricus</i> (Bloch & Schneider, 1801)
		<i>Caranx sexfasciatus</i> Quoy & Gaimard, 1825
		<i>Decapterus russelli</i> (Rüppell, 1830)
		<i>Gnathanodon speciosus</i> (Forsskal, 1775)
		<i>Megalaspis cordyla</i> (Linnaeus, 1758)
		<i>Parastromateus niger</i> (Bloch, 1795)
		<i>Scomberoides commersonnianus</i> Lacepède, 1801
		<i>Scomberoides tol</i> (Cuvier, 1832)
		<i>Selar crumenophthalmus</i> (Bloch, 1793)
		<i>Seriolina nigrofasciata</i> (Rüppell, 1829)
		<i>Trachinotus baillonii</i> (Lacepède, 1801)
		<i>Trachinotus mookalee</i> Cuvier, 1832
		<i>Uraspis helvola</i> (Forster, 1801)
	Chaetodontidae	<i>Chaetodon gardineri</i> Norman, 1939
	Drepanidae	<i>Drepane longimana</i> (Bloch & Schneider, 1801)
		<i>Drepane punctata</i> (Linnaeus, 1758)
	Echeneidae	<i>Echeneis naucrates</i> Linnaeus, 1758

Survey of the marine fishes

	Ephippidae	<i>Ephippus orbis</i> (Bloch, 1787)
		<i>Platax teira</i> (Forsskål, 1775)
	Gerreidae	<i>Gerres infasciatus</i> Iwatsuki & Kimura, 1998
		<i>Gerres limbatus</i> Cuvier, 1830
		<i>Gerres longirostris</i> (Lacepède, 1801)
		<i>Gerres oyena</i> (Forsskål, 1775)
	Gobiidae	<i>Acentrogobius dayi</i> Koumans, 1941
		<i>Bathygobius fuscus</i> (Rüppell, 1830)
		<i>Boleophthalmus dussumieri</i> Valenciennes, 1837
		<i>Cryptocentrus lutheri</i> (Klausewitz, 1960)
		<i>Periophthalmus waltoni</i> Koumans, 1941
		<i>Trypauchen vagina</i> (Bloch & Schneider, 1801)
	Haemulidae	<i>Diagramma pictum</i> (Thunberg, 1792)
		<i>Plectorhinchus pictus</i> (Tortonese, 1936)
		<i>Plectorhinchus sordidus</i> (Klunzinger, 1870)
		<i>Pomadasys aheneus</i> McKay & Randall, 1995
		<i>Pomadasys argenteus</i> (Forsskål, 1775)
		<i>Pomadasys kaakan</i> (Cuvier, 1830)
		<i>Pomadasys maculatus</i> (Bloch, 1793)
		<i>Pomadasys olivaceus</i> (Day, 1875)
		<i>Pomadasys punctulatus</i> (Rüppell, 1838)
		<i>Pomadasys stridens</i> (Forsskål, 1775)
		<i>Pomadasys taeniatus</i> McKay & Randall, 1995
	Labridae	<i>Cheilinus lunulatus</i> (Forsskål, 1775)
	Lactariidae	<i>Lactarius lactarius</i> (Bloch & Schneider, 1801)
	Leiognathidae	<i>Leiognathus equulus</i> (Forsskål, 1775)
		<i>Leiognathus oblongus</i> (Valenciennes, 1835)
		<i>Nuchequula gerreoides</i> (Bleeker, 1851)
		<i>Photopectoralis bindus</i> (Valenciennes, 1835)
	Lethrinidae	<i>Lethrinus borbonicus</i> Valenciennes, 1830
		<i>Lethrinus microdon</i> Valenciennes, 1830
		<i>Lethrinus nebulosus</i> (Forsskål, 1775)
	Lutjanidae	<i>Lutjanus argentimaculatus</i> (Forsskål, 1775)
		<i>Lutjanus ehrenbergii</i> (Peters, 1869)
		<i>Lutjanus fulviflamma</i> (Forsskål, 1775)
		<i>Lutjanus lutjanus</i> Bloch, 1790
		<i>Lutjanus malabaricus</i> (Bloch & Schneider, 1801)
		<i>Lutjanus russellii</i> (Bleeker, 1849)
		<i>Lutjanus sanguineus</i> (Cuvier, 1828)
		<i>Pristipomoides filamentosus</i> (Valenciennes, 1830)
	Menidae	<i>Mene maculata</i> (Bloch & Schneider, 1801)
	Mullidae	<i>Parupeneus cyclostomus</i> (Lacepède, 1801)
		<i>Parupeneus margaritatus</i> Randall & Gueze, 1984
		<i>Upeneus doriae</i> (Günther, 1869)
		<i>Upeneus tragula</i> Richardson, 1846
		<i>Upeneus vittatus</i> (Forsskål, 1775)
	Nemipteridae	<i>Nemipterus bipunctatus</i> (Valenciennes, 1830)
		<i>Nemipterus japonicus</i> (Bloch, 1791)

		<i>Nemipterus peronii</i> (Valenciennes, 1830)
		<i>Nemipterus randalli</i> (Russell, 1986)
		<i>Scolopsis taeniata</i> (Cuvier, 1830)
	Polynemidae	<i>Eleutheronema tetradactylum</i> (Shaw, 1804)
		<i>Polydactylus sextarius</i> (Bloch & Schneider, 1801)
	Pomacanthidae	<i>Pomacanthus maculosus</i> (Forsskål, 1775)
	Pomacentridae	<i>Neopomacentrus sindensis</i> (Day, 1873)
	Priacanthidae	<i>Priacanthus hamrur</i> (Forsskål, 1775)
		<i>Priacanthus tayenus</i> Richardson, 1846
	Rachycentridae	<i>Rachycentron canadum</i> (Linnaeus, 1766)
	Scaridae	<i>Chlorurus sordidus</i> (Forsskål, 1775)
		<i>Scarus ghobban</i> Forsskål, 1775
	Scatophagidae	<i>Scatophagus argus</i> (Linnaeus, 1766)
	Sciaenidae	<i>Argyrosomus hololepidotus</i> (Lacepède, 1801)
		<i>Johnius amblycephalus</i> (Bleeker, 1855)
		<i>Johnius belangerii</i> (Cuvier, 1830)
		<i>Johnius dussumieri</i> (Cuvier, 1830)
		<i>Nibea maculata</i> (Bloch & Schneider, 1801)
		<i>Otolithes ruber</i> (Bloch & Schneider, 1801)
		<i>Pennahia anea</i> (Bloch, 1793)
		<i>Protonibea diacantha</i> (Lacepède, 1802)
	Scombridae	<i>Euthynnus affinis</i> (Cantor, 1849)
		<i>Rastrelliger kanagurta</i> (Cuvier, 1816)
		<i>Scomberomorus commerson</i> (Lacepède, 1800)
		<i>Scomberomorus guttatus</i> (Bloch & Schneider, 1801)
	Serranidae	<i>Cephalopholis hemistiktos</i> (Rüppell, 1830)
		<i>Epinephelus areolatus</i> (Forsskål, 1775)
		<i>Epinephelus bleekeri</i> (Vaillant, 1878)
		<i>Epinephelus coioides</i> (Hamilton, 1822)
		<i>Epinephelus diacanthus</i> (Valenciennes, 1828)
		<i>Epinephelus epistictus</i> (Temminck & Schlegel, 1842)
		<i>Epinephelus malabaricus</i> (Bloch & Schneider, 1801)
		<i>Epinephelus polylepis</i> Randall & Heemstra, 1991
		<i>Epinephelus stoliczkae</i> (Day, 1875)
	Siganidae	<i>Siganus canaliculatus</i> (Park, 1797)
	Sillaginidae	<i>Sillago arabica</i> McKay & McCarthy, 1989
		<i>Sillago attenuata</i> McKay, 1985
		<i>Sillago sihama</i> (Forsskål, 1775)
	Sparidae	<i>Acanthopagrus arabicus</i> Iwatsuki, 2013
		<i>Acanthopagrus berda</i> (Forsskål, 1775)
		<i>Acanthopagrus bifasciatus</i> (Forsskål, 1775)
		<i>Argyrops spinifer</i> (Forsskål, 1775)
		<i>Crenidens crenidens</i> (Forsskål, 1775)
		<i>Diplodus kotschy</i> (Steindachner, 1876)
		<i>Rhabdosargus haffara</i> (Forsskal, 1775)

Survey of the marine fishes

		<i>Rhabdosargus sarba</i> (Forsskål, 1775)
		<i>Sparidentex hasta</i> (Valenciennes, 1830)
	Sphyraenidae	<i>Sphyraena obtusata</i> Cuvier, 1829
		<i>Sphyraena qenie</i> Klunzinger, 1870
	Stromateidae	<i>Pampus argenteus</i> (Euphrasen, 1788)
	Terapontidae	<i>Terapon jarbua</i> (Forsskål, 1775)
		<i>Terapon puta</i> Cuvier, 1829
		<i>Terapon theraps</i> Cuvier, 1829
		<i>Pelates quadrilineatus</i> (Bloch, 1790)
	Trichiuridae	<i>Trichiurus lepturus</i> Linnaeus, 1758
Pleuronectiformes	Cynoglossidae	<i>Cynoglossus arel</i> (Bloch & Schneider, 180)
		<i>Cynoglossus kopsii</i> (Bleeker, 1851)
	Paralichthyidae	<i>Pseudorhombus arsius</i> (Hamilton, 1822)
		<i>Pseudorhombus javanicus</i> (Bleeker, 1853)
	Psettodidae	<i>Psettodes erumei</i> (Bloch & Schneider, 1801)
	Soleidae	<i>Brachirus orientalis</i> (Bloch & Schneider, 1801)
		<i>Solea elongata</i> Day, 1877
		<i>Solea stanalandi</i> Randall & McCarthy, 1989
		<i>Zebrias captivus</i> Randall, 1995
Tetraodontiformes	Balistidae	<i>Abalistes stellaris</i> (Bloch & Schneider, 1801)
	Molidae	<i>Ranzania laevis</i> (Pennant, 1776)
	Monacanthidae	<i>Paramonacanthus choirocephalus</i> (Bleeker, 1851)
		<i>Paramonacanthus oblongus</i> (Temminck & Schlegel, 1850)
		<i>Thamnaconus modestoides</i> (Barnard, 1927)
	Tetraodontidae	<i>Arothron stellatus</i> (Anonymous, 1798)
		<i>Lagocephalus lunaris</i> (Bloch & Schneider, 1801)
	Triacanthidae	<i>Triacanthus biaculeatus</i> (Bloch, 1786)
		<i>Pseudotriacanthus strigilifer</i> (Cantor, 1849)


Abbas J. Al-Faisal and Falah M. Mutlak

Plate (1): Cartilaginous fish species from the Iraqi marine waters



Chiloscyllium arabicum

Carcharhinus leucas



Rhizoprionodon acutus

Brevitrygon imbricata



Brevitrygon walga



Himantura uarnak



Maculabatis gerrardi



Pastinachus sephen



Glaucostegus granulatus



Gymnura poecilura

Survey of the marine fishes

Plate (2): Bony fish species from the Iraqi marine waters.



Pegasus volitansi



Pseudosynanceia melanostigma



Ariomma indica

Platax teira



Gerres limbatus

Gerres longirostris



Cryptocentrus lutheri

Pomadasys maculatus



Pomadasys taeniatus

Lactarius lactarius



Lethrinus nebulosus

Lutjanus ehrenbergli



Lutjanus lutjanus



Lutjanus malabaricus



Mene maculate



Nemipterus peronii



Pomacanthus maculosus



Neopomacentrus sindensis



Priacanthus tayenus



Rachycentron canadum

Survey of the marine fishes



Argyrosomus hololepidotus

Protonibea diacantha



Euthynnus affinis

Cephalopholis hemistiktos



Siganus canaliculatus

Acanthopagrus bifasciatus



Sphyaena obtusata

Cynoglossus kopsii



Paramonacanthus choirocephalus *Pseudotriacanthus strigilifer*

Abbas J. Al-Faisal and Falah M. Mutlak

LITERATURE CITED

- Al-Daham, N. K. 1982. The ichthyofauna of Iraq and the Arab Gulf: A check-list. *Publications of the Basrah Natural History Museum*, 4: 1-102.
- Ali, T. S. 1985. Preliminary study on the nature of bony fish assemblage in Khor Al-Zubair. M. Sc. Thesis, College of Science, University of Basrah, 108 pp. (In Arabic)
- Ali, T. S. and Hussain, N. A. 1990. Composition and seasonal fluctuations of intertidal fish assemblage in Khor Al-Zubair, northwest Arabian Gulf. *Journal of Applied Ichthyology*, 6(1): 24-36.
- Al-Mudaffar, N. F. and Mahdi, B. A. 2014. Iraq's inland water quality and their impact on the North-Western Arabian Gulf. *Marsh Bulletin*, 9(1): 1-22.
- Al-Yamani, F. Y. 2008. Importance of the freshwater influx from the Shatt-Al-Arab River on the Gulf marine environment, 207-222. In: Abuzinada, A., Barth, H., Krupp, F., Böer, B. and Al-Abdessalaam, T.Z (Eds.). Protecting the Gulf's Marine Ecosystems from Pollution. Birkhäuser Basel publisher, 1st edition, XVI+ 285pp.
- Assadi, H. and Dehgani, R. P. 1997. Atlas of the Persian Gulf & the Sea of Oman Fishes. Iranian Fisheries Research and Training Organization, Tehran, Vol. 1, xx+226 pp.
- Bishop, J. M. 2003. History and current checklist of Kuwait's Ichthyofauna. *Journal of Arid Environments*, 54: 237-256.
- Blegvad, H. and Løppenthin, B. 1944. Fishes of Iranian Gulf. Danish Scientific investigation in Iran, Copenhagen, 247 pp.
- Carpenter, K. E.; Krupp, F.; Jones, D. A. and Zajonz, U. 1997. FAO species identification guide for fishery purposes. The living marine resources of Kuwait, Eastern Saudi Arabia, Bahrain, Qatar, and the United Arab Emirates. Rome, FAO, 293 pp.
- Froese, R. and Pauly, D. (Eds.) 2018. FishBase. world wide Web electronic publication. Available at: www.fishbase.org (version 02/2018).
- Hussain, N. A., Ali, T. S. and Naama, A. K. 1994. The effect of a heavy river flood on the fish assemblage structure in Khor Al-Zubair, northwest Arabian Gulf, Iraq. *Acta Ichthyologica et Piscatoria*, 24(2): 25-34.
- Hussain, N. A., Mohamed A. R. M. and Ali T. S. 1999. The seasonal formation of thermocline, halocline and water masses in the Iraqi marine waters. *Marina Mesopotamica*, 14(2): 299-312.
- Hussain, N. A. and Naama, A. K. 1989. Survey of fauna of Khor Al-Zubair, northwest Arabian Gulf. *Marina Mesopotamica*, 4(2): 161-197.
- Hussain, N. A., Naama, A. K. and Al-Hassan, L. A. J. 1988. Annotated checklist of the fish fauna of Khor Al-Zubair, north west of the Arabian Gulf, Iraq. *Acta Ichthyologica et Piscatoria*, 18(1): 17-24.

Survey of the marine fishes

- Hussain, N. A. and Younis, K. H. 1997. Fish assemblage of Khor Shettana, northwestern Arabian Gulf, Iraq. *Marina Mesopotamica*, 12(2): 331-356.
- Iwatsuki, Y., Jawad, L. A., Tanaka, F., Al-Busaidi, H., Al-Mamry, J. M. and Al-Kharusi, L. H. 2013. Omani fishes collected in the vicinity of Mutrah, Gulf of Oman and Madrasah, southern Oman through 3 to 13 October 2010. *Bulletin of Faculty of Agriculture*, University of Miyazaki, 59: 29-43.
- Jabado, R. W., Al Ghais, S. M., Hamza, W., Shivji, M. S. and Henderson, A. C. 2015. Shark diversity in the Arabian/Persian Gulf higher than previously thought: insights based on species composition of shark landings in the United Arab Emirates. *Marine Biodiversity*, 45(4): 719-731.
- Khalaf, K.T. 1961. The marine and freshwater fishes of Iraq. Al Rabbita press. Baghdad, 104pp.
- Kuronuma, K. and Abe, Y. 1986. Fishes of Arabian Gulf. Kuwait Institute for Scientific Research, 357 pp.
- Mahdi, N. 1962. Fishes of Iraq. Ministry of Education, Baghdad, 82 pp.
- Mahdi, N. and Georg, P. V. 1969. A systematic list of the vertebrates of Iraq. *Iraq Natural History Museum, Publications*, No 26: 1-104.
- Mohamed, A. R. M., Hussain, N. A. and Ali, T. S. 2001. Estuarine components of the ichthyofauna of the Arabian Gulf. *Marina Mesopotamica*, 16(2): 209-224.
- Nelson, J. S. 2006. Fishes of the World. 4th Edition. John Wiley & Sons Inc., New Jersey, xiv + 601 pp.
- Randall J. E. 1995. Coastal fishes of Oman. Crawford House Publishing Pty Ltd., Hawaii, 456 pp.
- Younis, K. H. 1990. A study of fish assemblage in the north of Khor Abdullah. M. Sc. Thesis, Marine Science Centre., University of Basrah, 95 pp. (In Arabic)

Abbas J. Al-Faisal and Falah M. Mutlak

Bull. Iraq nat. Hist. Mus.

December, (2018) 15 (2): 163-177

مسح لأنواع الأسماك البحرية في العراق

عباس جاسم الفيصل و فلاح معروف مطلق
مركز علوم البحار، جامعة البصرة، البصرة، العراق

تاريخ القبول: 2018/08/28

تاريخ الاستلام: 2018/04/19

الخلاصة

أجري مسح لأنواع الأسماك المتواجدة في المياه البحرية العراقية للمدة من تشرين الثاني 2014 ولغاية آذار 2018. شملت الدراسة تسجيل 214 نوعاً تعود الى 75 عائلة.

وجدت عائلة Carangidae هي الأكثر سيادة وتمثلت بـ 24 نوعاً ثم تلتها عائلة Haemulidae بـ 11 نوعاً ثم عائلتي Serranidae و Sparidae بتسعة انواع لكل منهما، بينما ضمت العوائل الـ 34 عائلة نوعاً واحداً فقط.