

DIETARY REPLACEMENT OF SOYBEAN MEAL BY TOASTED SUNFLOWER SEEDMEAL IN THE DIET OF *Clarias gariepinus*: EFFECT ON GROWTH, BODY COMPOSITION, DIGESTIBILITY, HAEMATOLOGY AND HISTOPATHOLOGY OF THE LIVER.

JIMOH Wasiu Adeyemi

Senior Lecturer

Dept. Aquac. and Fisher. Fac. Agric. University of Ilorin, PMB 1515, Ilorin, Nigeria

jimoh.wa@unilorin.edu.ng,

ABSTRACT

Dietary replacement of soybean meal by toasted sunflower meal in *Clarias gariepinus* diet was investigated in a 56-day feeding trial using growth, body composition, digestibility, haematology and liver histology as indices of assessment. Sunflower seed meal was toasted at 204°C for 5, 10 and 15 minutes. Control diet consist of soybean meal which was replaced by differently toasted sunflower seed meal at a rate 15, 30 and 45% to produce nine 40% crude protein, 18kj/kg test diets. Triplicate groups of fish in 70-litre capacity aerated rectangular plastic tanks were assigned to each dietary treatment at a stocking rate of 15 fingerlings (4.97±0.42g average weight) per tank in a completely randomized design. Fish were fed to satiation. Data generated from the experiment were subjected to one-way analysis of variance. The results of the experiment revealed that up to 30% replacement level of the differently toasted sunflower had a statistically similar results with soybean-based control diets using growth, body composition, digestibility and haematology as indices. Liver histology however show that massive fatty infiltration recorded in the liver of fish fed TSF145 and TSF 345 and diffuse vacuolar degeneration of hepatocytes and kupffer cell hyperplasia observed in the liver of fish fed TSF245 and TSF330.

Keywords: histology, toasted, triplicate groups, seed meal.

اديمي

مجلة العلوم الزراعية العراقية -2020: 51(4): 1088-1103

تأثير الاستبدال الجزئي لبذور عباد الشمس بديلا عن كسبة فول الصويا في عليقة سمكة *Clarias gariepinus*

على النمو ومكونات الجسم والهضم وبعض معايير الدم ونسيج الكبد

واسيو اديمي

مدرس أقدم

قسم تربية الأحياء المائية ومصايد الأسماك، كلية الزراعة، جامعة إيلورين، صندوق بريد خاص 1515، إيلورين نيجيريا

المستخلص

نفذت تجربة استبدال جزئي لبذور عباد الشمس بديلا عن كسبة فول الصويا في عليقة سمكة *Clarias gariepinus* لمدة 56 يوما لدراسة النمو ومكونات الجسم والهضم وبعض معايير الدم ونسيج الكبد. عرضت بذور عباد الشمس الى درجات حرارة 24 م° لمدة 5 و 10 و 15 دقيقة . تم استبدال عليقة السيطرة الحاوية على كسبة فول الصويا ببذور عباد الشمس بنسب 15 و 30 و 45% للحصول على 40% بروتين كلي، 18 جول / كغم في العليقة. استخدم ثلاث مكررات في معاملات الاسماك باستعمال احواض بلاستيكية بسعة 70 لتر للحوض الواحد وزعت صغار الاسماك عشوائياً بواقع 15 سمكة وبمعدل وزن 4,97 ± 0,42 غم لكل حوض اذ غذيت الاسماك لحد الاشباع . اتبع نظام التباين باتجاه واحد في تحليل النتائج احصائيا. بينت نتائج التجربة بان معاملة الاستبدال 30% بذور عباد الشمس مشابهة الى نتائج عليقة السيطرة في صفات النمو ومكونات الدم والهضم والدم . اظهرت نتائج الفحص النسيجي للكبد ارتشاح كبير للدهون في الاسماك المغذات على TSF145 و TSF 345 مع تنكس دهني لخلايا الكبد وتضخم خلايا كويفر في الاسماك المغذاة على TSF245 و TSF330 .

الكلمات المفتاحية: تغذية، دم، بروتين، احواض الاسماك، عليقة السيطرة

*Received:24/8/2019, Accepted:30/12/2019

INTRODUCTION

Aquaculture contribution to total fish production around the globe is increasing every year with global production estimated to be 110.2 million in 2016 (22). This rapid growth of aquaculture in the world is facilitated by proliferation of semi- and intensive culture system which rely majorly on complete feed production (23). The major ingredient in complete feed manufacture is fishmeal but its use is not sustainable owing to its dwindling availability as a result of ever-increasing demand, overfishing of wild stock and prohibitive cost (6; 31; 70-72). Fishmeal has exceptional nutritive profile that are essential, particularly for fish. Fishmeal production is quantitatively and qualitatively variable because it depends on catch gotten from the fishery (60). Soybean meal have successfully been tested and now being used to replace at least 50% of the fishmeal portion in fish feed (67). McGoogan and Gatlin III (47) reported that 90% of the protein from soybean meal could produce comparable weight gain as 100% of protein from fish meal. El-Saidy and Gaber (14) reported the possibility of total replacement of fishmeal with 55% soybean meal supplemented with 0.5% L-lysine in Nile tilapia fingerlings diet. This is because soybean meal has the best amino acid profile of the plant protein sources available, highly palatable and digestible (9). However, owing to its different use as food for man and other livestock, its price is becoming prohibitive in the world especially among fish farmers in the developing countries. Ng and Romano (53) reported that numerous alternatives to soybean meal have to be explored in order to witness cost-effective and sustainable aquaculture growth. Such a veritable plant protein that could be used to replace soybean meal must be less costly, readily available locally and must possess nutrient density that is relatively comparable to soybean meal that would provide greater flexibility to feed millers and farmers to improve cost-effectiveness (18). Sunflower protein has such potential to replace soybean meal. It is widely produced in Nigeria. Fagbenro et al (20) reported its protein content ranging from 35.98% for sunflower cooked at 100°C for 30 minutes to 44.29% crude protein for sunflower toasted at

204°C for 10 minutes. Sunflower seed meal has been tested in the diets of many fishes: Tilapia, *Oreochromis niloticus*, (13; 15; 44; 55; 65); Trout, *Oncorhynchus mykiss*, (50); Silver barb, *Puntius gonionotus*, (48; 49); hybrid striped bass, *Morone chrysops* x *M. saxatilis*, (29); Russian sturgeon, *Acipenser gueldenstaedtii*, (62); Indian major carp, *Cirrhinus cirrhosus* (1); Salmon, *Salmo salar*, (30); Shrimp, *Penaeus schmitti*, (25); Gilthead sea bream, *Sparus aurata*, (4; 43); Abalone *Haliotis midae* (58; 59; 63; 64); European Eel, *Anguilla Anguilla*, (11; 27). Jackson et al (36) reported good growth in tilapia (*Sarotherodon mossambicus*) fed rations containing 35.2% sunflower meal replacing 50% of the fish meal protein. Martinez (46) reported that there was no loss in growth performance and diet utilization efficiency when rainbow trout (*Salmo girdineri*) was fed diet containing 22 and 37.3% sunflower meal though he added L-methionine to that of 37.5% sunflower meal. Tacon et al (69) earlier reported that 36.5% sunflower seed meal had a statistically similar growth with soybean control ration, that supplementation with 0.2% exogenous L-methionine did not yield beneficial effect on fish growth and feed utilization efficiency. Stickney et al (66) found that sunflower protein concentrate can replace 25% fish meal protein in rainbow trout diets. Sanz et al (61) observed better results when replacing 40% of the animal protein in rainbow trout with sunflower protein supplemented with EAA. However, the antinutrient present in sunflower meal restricted its use as soybean replacer in fish feed. Fagbenro et al (20) reported differently processed sunflower meal to contain significantly reduced content of trypsin inhibitors, lectin, tannin, phytin, saponin and oxalates which was completely eliminated at 30 minutes cooking and 15 minutes toasting at 204°C. Francis et al (26) reported how inclusion level could affect physiological functioning of fish with respect to antinutrients present in fish feed ingredients that fish generally has compensatory mechanism to cope with antinutrients in feed when it does not exceed the critical limit. The level of destruction of antinutrients in feed ingredients is generally related to moisture conditions, particle size, duration of heating

and temperature (37; 41). This study attempts to examine the effect of different toasting time and inclusion level of sunflower protein on the growth, body composition, digestibility, haematological and histopathological condition of the liver of *Clarias gariepinus*

MATERIALS AND METHODS

Diet preparation

Toasted sunflower meal were produced by toasting sunflower seed at 204⁰C for 5, 10 and 15 minutes and separately milled into powdery form by the use of split-disc grinder. The differently toasted sunflower meal was taken for proximate (5) and amino acid analysis. Based on their proximate composition of differently toasted sunflower seedmeal (Table 1), ten dietary treatments were developed, consisting of a control diets and nine test diets. The control diet was prepared with soybean replacing up to 50% of the fishmeal and other feeding ingredients to produce 40% crude protein diet. The nine test diets were prepared by replacing soybean portion of the control diet with each of the differently toasted sunflower meal at 15, 30 and 45% replacement level. Proximate analysis (5) of the dietary treatments was done in triplicates (Table 2).

Experimental fish and system design

Clarias gariepinus fingerlings (4.97±0.42g average weight) were obtained from Alagbaka fish farm Akure and transported live in an

aerated polythene bags to the Fish Nutrition Laboratory of Federal University of Technology, Akure. The fish were acclimated to laboratory condition for 14 days during which they were fed complete imported feed. Triplicate groups of randomly picked fifteen fingerlings in aquaria containing 70-litre water were randomly assigned to each dietary treatment with continuous aeration. Group feeding of fish were done two times a day at 5% body weight on equal proportion in the morning 8:00-9:00hr and evening 17:00-18:00hrs. Batch weighing were conducted every other week and the diet rations were adjusted according to their body weight. Mortality and water quality parameters were monitored routinely during the period of the feeding trial.

Whole-body composition

Six fingerlings at the beginning and six fish from each treatment at the end of the experiment were randomly selected for proximate analysis (5).

Digestibility study

Faecal samples from each tank were pooled together and oven dried for 24 hours at 55⁰C. The feed and the faecal samples were analyzed for acid insoluble ash (AIA) using the method explained Jimoh et al (2010) and the apparent organic matter (AOMD) and nutrient

Table 1. Proximate (g/100g dry matter) and amino acid (g/100g protein) composition and sunflower seed meal

Proximate	TSF5	TSF10	TSF15
Moisture	9.23 ± 0.08	9.36 ± 0.18	9.48 ± 0.06
Crude Protein	42.08 ± 1.68	44.29 ± 2.26	39.76 ± 1.42
Crude Lipid	13.28 ± 1.09	12.33 ± 0.84	12.05 ± 2.92
Crude Fibre	14.22 ± 2.2	14.38 ± 0.66	15.37 ± 0.56
Ash	5.99 ± 1.01	6.32 ± 1.31	6.23 ± .45
NFE	15.18 ± 2.03	13.33 ± 1.20	17.79 ± 4.4
Amino Acid			
Lysine	3.64 ± 0.02	4.32 ± 0.01	5.11 ± 0.03
Histidine	2.24 ± 0.08	2.04 ± 0.01	1.91 ± 0.01
Arginine	8.57 ± 0.03	8.90 ± 0.02	9.05 ± 0.04
Aspartic Acid	8.98 ± 0.04	8.95 ± 0.04	9.09 ± 0.03
Threonine	3.10 ± 0.07	3.66 ± 0.04	3.81 ± 0.03
Serine	3.77 ± 0.01	3.61 ± 0.01	3.66 ± 0.01
Glutamic Acid	14.29 ± 0.01	14.33 ± 0.04	14.47 ± 0.02
Proline	2.66 ± 0.02	2.81 ± 0.01	2.75 ± 0.03
Glycine	3.03 ± 0.02	6.22 ± 0.02	3.27 ± 0.04
Alanine	3.01 ± 0.03	3.11 ± 0.01	3.31 ± 0.03
Cystine	1.45 ± 0.04	1.36 ± 0.01	1.00 ± 0.14
Valine	3.92 ± 0.04	3.93 ± 0.05	3.78 ± 0.01
Methionine	1.27 ± 0.04	1.21 ± 0.01	1.09 ± 0.01
Isoleucine	4.80 ± 0.03	5.81 ± 0.01	5.16 ± 0.01
Leucine	5.06 ± 0.04	5.16 ± 0.02	5.77 ± 0.04
Tyrosine	1.41 ± 0.02	1.57 ± 0.02	1.72 ± 0.02
Phenylalanine	4.36 ± 0.02	3.93 ± 0.04	3.77 ± 0.03

digestibility coefficient (ADC) of the nutrients in each diets estimated using the formula

$$ADC (\%) = 100 - \left(100 \left(\frac{\% \text{ AIA in Feed}}{\% \text{ AIA in Faeces}} \right) \times \frac{\% \text{ Nutrients in Faeces}}{\% \text{ Nutrients in Feed}} \right)$$

Blood sampling and hematological analysis

At the end of the 56-day feeding trial, the fish were left unfed for 24 hours and were euthanized in 100ml/l clove oil. The blood samples of the six fish from each tank were collected and pooled together in EDTA bottle using heparinized syringe. Haematological analysis were carried out following the

standard procedures explained in Fagbenro et al (21) and Jimoh et al (38)

Hepatosomatic Index (HSI)

The six fish per dietary treatment that have been priorly individually weighed were euthanized in 100ml/l clove oil. The fish were dissected and the livers were removed. The removed livers weight from each dietary treatment were measured to calculate Hepatosomatic index (HSI).

Histological examination

Figure 1 describes the steps followed in conducting histological examination for the excised livers. The interpretation of the digital images of the liver of each dietary treatment taken was conducted at the Department of Veterinary Anatomy, University of Ibadan, Ibadan, Nigeria.




Figure 1. flow-chart describing histological examination of the excised liver

RESULTS AND DISCUSSION

Growth Performance and Nutrient Utilization

Growth performance and nutrient utilization of *Clarias gariepinus* fed varying levels toasted sunflower meal based experimental diets is presented in Table 3. There was no significant difference ($p > 0.05$) in the mean initial weight in the experimental fish. The fish responded well to all the dietary treatment given as evidenced in the percentage survival that was very high and did not significantly differ ($p > 0.05$). At the end of the feeding trial, the final weight of fish fed up to 30% replacement level of variously toasted sesame seed meal did not significantly differ ($p > 0.05$) from the fish fed control diets, except the fish fed

TSF145, TSF245 and TSF345. The mean weight gain also followed the same trend as observed for mean final weight. The percentage weight gain of fish fed TSF215 was the highest. Specific growth rate of fish fed various dietary treatment did not show any significant difference from the control except those fed TSF145, TSF245 and TSF345. The highest value of feed conversion ratio was recorded in fish fed TSF145. The protein efficiency ratio of the fish fed various dietary treatments followed a reversed trend with the results observed for feed conversion ratio. The protein retention of the fish fed various dietary treatment were not significantly ($p > 0.05$) different from each other.

Table 2. Gross (g/100g) and proximate (g/100g dry matter) composition of experimental diets at varying replacement levels of differently toasted sunflower seed meals

Ingredients	CTR	TSF115	TSF130	TSF145	TSF215	TSF230	TSF245	TSF315	TSF330	TSF345
Fishmeal	27.24	27.24	27.24	27.24	27.24	27.24	27.24	27.24	27.24	27.24
Soybean Meal	46.71	39.71	32.70	25.70	39.71	32.70	25.70	39.71	32.70	25.70
Toasted Sunflower	-	7.13	14.25	21.38	6.78	13.55	20.33	7.55	15.09	22.64
Corn Meal	11.25	11.25	11.25	11.25	11.25	11.25	11.25	11.25	11.25	11.25
Fish Oil	5.09	5.09	5.09	5.09	5.09	5.09	5.09	5.09	5.09	5.09
*Vit/Min Premix	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00	5.00
Starch	4.71	4.58	4.47	4.34	4.93	5.17	5.39	4.16	3.63	3.08
Proximate										
Moisture	9.45±0.19	9.49±0.30	9.41±0.23	9.46±0.31	9.45±0.15	9.51±0.13	9.58±0.09	9.46±0.19	9.54±0.06	9.52±0.24
Crude Protein	40.18±0.03	40.20±0.23	40.25±0.15	40.12±0.11	40.16±0.11	40.10±0.24	40.18±0.08	40.28±0.27	40.28±0.24	40.18±0.23
Crude Lipid	12.17±0.08	12.17±0.21	12.21±0.11	12.15±0.07	12.28±0.07	12.19±0.14	12.15±0.11	12.23±0.12	12.22±0.03	12.16±0.11
Ash	8.73±1.46	8.41±0.77	8.62±0.46	8.99±0.24	8.61±0.73	8.84±0.71	8.48±1.00	8.35±0.11	8.57±0.98	9.35±0.08
Crude Fibre	6.36 ±0.12 ^{ab}	6.50±0.12 ^a	6.14±0.13 ^b	6.26±0.09 ^{ab}	6.48±0.17 ^a	6.27±0.10 ^{ab}	6.40±0.16 ^{ab}	6.35±0.05 ^{ab}	6.38±0.02 ^{ab}	6.51±0.34 ^a
NFE	23.12±1.46	23.24±0.73	23.37±0.65	23.01±0.51	23.03±0.50	23.09±1.05	23.21±1.02	23.33±0.65	23.01±0.77	22.28±0.21
AIA (%)	0.76±0.06	0.81±0.02	0.78 ±0.03	0.79±0.26	0.77±0.05	0.78±0.02	0.80±0.02	0.81±0.04	0.81±0.01	0.81±0.02
Energy (kcal/100g)	435.37±6.56	436.01±3.04	437.21±1.92	434.48±1.93	435.93±3.09	435.03±2.07	435.58±4.47	437.42±0.30	435.99±4.12	431.86±1.16

* Specification: each kg contains: Vitamin A , 4,000,000IU; Vitamin B, 800,000IU; Vitamin E, 16,000mg, Vitamin K₃, 800mg; Vitamin B₁, 600mg; Vitamin B₂, 2,000mg; Vitamin B₆, 1,600mg, Vitamin B₁₂,8mg; Niacin,16,000mg; Caplan, 4,000mg; Folic Acid, 400mg; Biotin, 40mg; Antioxidant 40,000mg; Chlorine chloride, 120,000mg; Manganese, 32,000mg; Iron 16,000mg; Zinc, 24,000mg; Copper 32,000mg; Iodine 320mg; Cobalt,120mg; Selenium, 800mg manufactured by DSM Nutritional products Europe Limited, Basle, Switzerland

Row means with no superscripts are not significantly different ($p>0.05$)

Row means with different superscripts are significantly different ($p<0.05$)

Comparable growth performance that was not statistically different from control was observed in this study for *Clarias gariepinus* fed diets containing differently toasted sunflower meal replacing up to 30% of soybean meal. At higher replacement level of 45% of the differently toasted sunflower meal, there was significant reduction in growth of *Clarias gariepinus* plausibly attributed to incomplete detoxification of anti-nutrients in sunflower seedmeal by toasting, or increased deficiencies of essential amino acids. This result corroborates the finding of Olvera-Novoa et al (55) who observed growth depletion with diets containing higher sunflower seed meal percentage and attributed it to increasing phenylalanine and methionine deficiencies and high non-digestible fibre content. Hertrampf and Piedad-Pascual (35) reported higher contents of crude fibre in sunflower meal reduced its utilization as dietary ingredients in aquaculture. González-Pérez and Vereijken (32) further explains that the phenolic compound found in sunflower meal such as caffeic acid and chlorogenic acid has the capacity to reduce protein solubility. Eusebio et al (17) and Zhou et al (75) attributed the lowered digestibility recorded at higher inclusion of plant protein to imbalanced amino acid; increased antinutrient and cellulose content as the plant protein increases. Köprücü and Sertel (40) also recorded lower growth in carp, *Ctenopharyngodon Idella*, fed sunflower meal when compared to soybean meal fed control group. Earlier work by Jackson et al (36); Tacon et al (69); Stickney et al (66) and Sintayehu et al (65) also reported lower growth performance of fish at higher inclusion of sunflower seedmeal. Francis et al (26) reported that fish generally has compensatory mechanism to cope with antinutrients in feed when it does not exceed the critical limit which practically suggests that at lower inclusion level of sunflower seedmeal, the critical level of the different anti-nutrients in them has not been exceeded.

Whole body composition

Table 4 revealed the whole-body composition of fish at the beginning and at the end of the experiment. There was significant difference ($p < 0.05$) between the initial and final body composition of fish used during the experiments with respect to moisture crude protein, crude lipid and ash content. Among the fish fed the experimental diets, the highest value of crude protein was recorded in fish fed CTR which was not significantly ($p > 0.05$) fish fed other dietary treatments except those fed TSF145, TSF245 and TSF345. Carcass lipid of fish fed the experimental diets did not show any significant variation ($p > 0.05$). Carcass ash in fish fed the TSF345, TSF245 and TSF230 did not show any significant difference ($p > 0.05$) from each other. No significant difference was found in carcass crude protein and crude lipid among the control group and sunflower fed groups up to 30% replacement level following same result trends in the growth studies. Although no significant variations were recorded in the protein retention among the different dietary treatment groups, there was marked differences in carcass protein and lipid. at higher replacement level of 45%. The reduction in carcass lipid at higher replacement level might not be unconnected with the phenolic compound, chlorogenic acid present in sunflower, an anti-oxidant that has the capacity to reduce carcass lipid (68). Hassaan et al (34) observed same trends of results on growth and carcass composition of *Oreochromis niloticus* fed fermented sunflower meal despite the fermentation process higher inclusion could

Table 3. Growth performance and nutrients utilization of *Clarias gariepinus* fed toasted sunflower meal-based diets

Parameters	Experimental Diets									
	CTR	TSF115	TSF130	TSF145	TSF215	TSF230	TSF245	TSF315	TSF330	TSF345
Initial Weight	5.24±0.37	5.49±0.16	5.04±0.49	4.67±0.00	4.62±0.07	4.79±0.47	4.81±0.47	5.08±0.07	4.90±0.55	5.07 ±0.65
Final Weight	15.57±0.41 ^a	15.79±0.14 ^a	15.23±0.56 ^a	13.90±0.04 ^b	15.62±0.41 ^a	15.23±0.19 ^a	13.46±0.26 ^b	15.42±0.73 ^a	15.14±0.55 ^a	13.39±0.32 ^b
¹ Mean Weight Gain	10.33±0.04 ^a	10.31±0.02 ^a	10.19±0.06 ^a	9.23±0.04 ^b	11.00±0.34 ^a	10.45±0.28 ^a	8.65±0.73 ^b	10.34±0.01 ^a	10.24±0.01 ^b	8.32±0.98 ^a
² % Weight Gain	197.90±13.49 ^{ab}	187.96±5.96 ^{ab}	203.20±18.69 ^{ab}	197.65±0.91 ^{ab}	238.07±3.70 ^a	219.65±27.51 ^{ab}	181.31±32.73 ^{ab}	205.61±29.90 ^{ab}	210.59±23.28 ^{ab}	166.70±40.64 ^b
³ SGR	2.72±0.03 ^a	2.73±0.01 ^a	2.69±0.04 ^a	2.60±0.00 ^b	2.72±0.03 ^a	2.70±0.01 ^a	2.57±0.02 ^b	2.70±0.04 ^a	2.69±0.03 ^a	2.57±0.03 ^{ab}
⁴ FCR	1.22±0.03 ^c	1.24±0.02 ^{bc}	1.26±0.02 ^{bc}	1.30±0.01 ^a	1.25±0.00 ^{bc}	1.24±0.01 ^{bc}	1.26±0.0 ^{bc}	1.24±0.01 ^{bc}	1.25±0.02 ^{bc}	1.28±0.02 ^{ab}
⁶ PER	2.05±0.04 ^a	2.01±0.03 ^{ab}	1.99±0.03 ^{ab}	1.92±0.02 ^c	2.00±0.00 ^{ab}	2.02±0.01 ^{ab}	1.99±0.02 ^{abc}	2.02±0.02 ^{ab}	2.00±0.02 ^{ab}	1.96±0.03 ^{bc}
⁷ Protein retention	49.53±2.54	49.65±4.74	46.87±2.29	47.61±2.97	46.44±1.27	46.04±1.90	50.91±4.20	49.88±2.34	47.52±0.99	51.24±8.19
⁸ Survival Rate (%)	96.67±4.71	96.67±4.71	93.33±0.00	86.67±0.00	96.67±4.71	96.67±4.74	90.00 ±4.71	96.67±4.71	90.00±4.71	86.67± 0.00

Row means with no superscripts are not significantly different ($p>0.05$)

Row means with different superscripts are significantly different ($p<0.05$)

¹ Mean weight gain= final mean weight –initial mean weight

² Percentage weight gain= [final weight-initial weight /initial weight] X 100

³ Specific growth rate= [In final weight-In initial weight] X 100

⁴ Feed conversion ratio=dry weight of feed fed /Weight gain (g)

⁵ Protein efficiency ratio=fish body weight (g)/ Protein fed ⁶Protein retention= [protein gain/protein fed] X 100

⁷ Percentage survival = {(total number of fish- mortality)/total number of fish] X 100

Table 4. Whole body composition of *Clarias gariepinus* fed varying replacement levels of toasted sunflower seedmeal based diets

Parameter	Initial	Experimental Diets									
		CTR	TSF115	TSF130	TSF145	TSF215	TSF230	TSF245	TSF315	TSF330	TSF345
Moisture	76.80±0.34 ^c	72.50±0.08 ^{bc}	72.59±0.09 ^{bc}	72.78±0.13 ^{bc}	71.96±1.64 ^c	72.60±0.15 ^{bc}	72.74±0.05 ^{bc}	73.03±0.06 ^b	72.53±0.10 ^{bc}	72.74±0.02 ^{bc}	73.09±0.32 ^b
Crude Protein	15.14±0.09 ^c	17.66±0.08 ^a	17.66±0.15 ^a	17.57±0.10 ^a	17.38±0.10 ^b	17.66±0.11 ^a	17.55±0.08 ^a	17.32±0.05 ^b	17.65±0.09 ^a	17.55±0.06 ^a	17.26±0.26 ^b
Crude Lipid	5.04±0.23 ^a	6.25±0.12 ^b	6.21±0.20 ^b	6.18±0.22 ^b	6.36±0.03 ^b	6.26±0.09 ^b	6.26±0.13 ^b	6.39±0.03 ^b	6.25±0.11 ^b	6.25±0.11 ^b	6.42±0.11 ^b
Ash	3.03±0.18 ^b	3.58±0.09 ^{ab}	3.55±0.14 ^{ab}	3.47±0.14 ^{ab}	4.30±1.68 ^a	3.47±0.05 ^{ab}	3.44±0.09 ^{ab}	3.26±0.04 ^b	3.56±0.12 ^{ab}	3.45±0.06 ^{ab}	3.23±0.04 ^b

Row means with no superscripts are not significantly different ($p>0.05$)

Row means with different superscripts are significantly different ($p<0.05$)

not avail better growth performance. This result is in tandem with earlier finding by Zhou et al (75) on sea bream, *Acanthopagrus schlegelii*, juveniles fed fermented soybean meal in diets and by Lim et al (42) on Korean rockfish, *Sebastes schlegeli* fed dehulled soybean meal. Other researchers who reported decrease in carcass protein and lipid with increasing plant protein are Jimoh et al (39); Hassaan et al (33) and Saha and Ghosh (57).

Apparent digestibility coefficients of the nutrients in each diet

Table 5 showed the proximate composition of faecal samples of *Clarias gariepinus* fed varying levels of toasted sunflower meal-based diets. There was reduction in the nutrient contents of the faecal samples tested when compared with the nutrient contents of the feed. The highest faecal protein in fish fed TSF345 while the least faecal protein was recorded in fish fed CTR. There were no significant differences ($p>0.05$) in the faecal protein of the fish fed TSF145, TSF245 and TSF345. However, fish exposed to other test diets apart from the above did not any significant variation ($p>0.05$) from the fish fed CTR. The highest faecal lipid was observed in fish fed TSF230 while the least faecal lipid was recorded in fish fed TSF315. There was no significant difference ($p>0.05$) in faecal lipid of fish fed control diets and fish fed all other test diets except the fish fed TSF230. Faecal energy of fish fed CTR were not significantly different ($p>0.05$) from the fish fed other dietary treatment. The faecal AIA of fish fed TSF345 recorded the least value; the fish fed TSF315 recorded the highest value. There was no significant variation ($p>0.05$) in the AIA of fish fed various dietary treatment. Table 6 presents apparent digestibility coefficient of nutrients in toasted sunflower meal-based diets fed to *Clarias gariepinus* fingerlings. The fish fed various dietary treatment showed significant difference in ($p<0.05$) in their nutrient digestibility's values. CTR had the highest apparent organic matter digestibility (AOMD) which was not significantly different from AOMD of fish fed TSF115, TSF130, TSF215, TSF230, TSF315 and TSF330 while TSF345 had the least AOMD. Same trend as above was observed for apparent protein digestibility (APD)

coefficient. There was no significant difference ($p>0.05$) in the APD of the fish fed CTR and fish fed TSF115, TSF130, TSF215, TSF230, TSF315 and TSF330. Fish fed TSF315 had the highest apparent lipid digestibility (ALD) coefficient which was not significantly different ($p>0.05$) from fish fed diet CTR, TSF145 and TSF215. The lowest ALD coefficient was recorded in fish fed TSF345 which was not significantly different ($p>0.05$) from fish fed TSF230, TSF245, TSF130, TSF115, TSF330 and TSF215. Similarly, the highest apparent energy digestibility (AED) coefficient was found in fish fed TSF315 while the lowest AED coefficient was found in fish fed 345. There was no significant difference ($p>0.05$) in the AED coefficient among the fish fed TSF315, CTR, TSF215, TSF130, TSF230 and TSF115. Hassaan et al (34) reported that higher inclusion of fermented sunflower seedmeal reduced protein and lipid digestibility in tandem with what was observed in this study. Eusebio et al (17) and Zhou et al (75) attributed the lowered digestibility recorded at higher inclusion of plant protein to imbalanced amino acid.

Haematological profile

The summary of comparison of haematological parameters of *Clarias gariepinus* fed toasted sunflower meal-based diets replacing soybean meal at different levels is presented in Table 7. There was no significant difference ($p>0.05$) in the primary haematological parameters (Hb, PCV, RBC, WBC) of the fish fed CTR and those fed differently toasted sunflower meal up to 30% replacement level. However, significant variations ($p<0.05$) existed between these fish and the fish fed 45% of the differently toasted sunflower seedmeal.

Table 5. Proximate composition (g/100g dry matter) of faecal samples of *Clarias gariepinus* fed varying replacement levels of toasted sunflower seedmeal based diets

	CTR	TSF115	TSF130	TSF145	TSF215	TSF230	TSF245	TSF315	TSF330	TSF345
Moisture	9.49 ±0.44	9.24±0.74	8.45±0.37	8.83±0.85	9.43±0.44	9.19±1.01	9.10 ±0.72	9.56 ±0.52	8.81±0.66	8.90±0.63
Crude Protein	14.13±1.51 ^a	14.36±1.04 ^a	14.78±1.33 ^a	19.81±0.71 ^b	14.30±1.15 ^a	15.04±1.32 ^a	19.96±0.84 ^b	14.37±0.24 ^a	14.81±1.07 ^a	19.93±2.10 ^b
Crude Lipid	6.19±0.42 ^{bc}	6.83±0.70 ^{abc}	7.10±0.54 ^{ab}	6.15±0.99 ^{bc}	6.95±0.14 ^{abc}	7.73±0.33 ^a	6.87±0.30 ^{abc}	5.97±0.85 ^c	6.63±0.16 ^{bc}	6.87±0.32 ^{abc}
Ash	8.27±1.30 ^{ab}	6.58±2.16 ^b	8.32±0.49 ^{ab}	8.66±0.34 ^a	8.59±0.65 ^a	9.37±0.64 ^a	8.47±1.13 ^{ab}	8.87±0.61 ^a	8.67±1.23 ^a	9.51±0.45 ^a
Crude Fibre	13.46±0.14 ^a	13.07±0.38 ^{ab}	12.67±0.46 ^{ab}	12.26±0.75 ^b	13.38±0.78 ^{ab}	12.95±0.22 ^{ab}	12.75±0.79 ^{ab}	13.67±0.81 ^a	12.75±0.15 ^{ab}	12.54±0.71 ^{ab}
NFE	48.47±0.84 ^{ab}	49.93±0.40 ^a	48.67±1.45 ^{ab}	44.29±1.53 ^{cd}	47.35±1.73 ^{ab}	45.72±1.06 ^{bc}	42.85±2.55 ^{cd}	47.56±1.87 ^{ab}	48.33±1.72 ^{ab}	42.16±2.45 ^d
AIA (%)	3.36±1.73	3.44±0.26 ^a	3.42±0.21 ^a	3.22±0.09 ^{ab}	3.37±0.31 ^{ab}	3.37±0.09 ^{ab}	3.27±0.05 ^{ab}	3.54±0.19 ^a	3.32±0.09 ^{ab}	3.04±0.03 ^b
Energy (kcal/100g)	336.67±7.1 ^{bc}	349.98±11.19 ^{ab}	349.80±4.39 ^{ab}	350.93±7.02 ^a	340.24±5.46 ^{abc}	345.05±7.61 ^{abc}	352.79±7.53 ^a	332.17±3.29 ^c	344.08±7.48 ^{abc}	349.73±54.21 ^{ab}

Row means with different superscripts are significantly different (p<0.05)

NFE: Nitrogen free Extract AIA: Acid insoluble Ash

Table 6. Apparent digestibility coefficient of nutrients in toasted sunflower meal-based diets fed to *Clarias gariepinus*

	CTR	TSF115	TSF130	TSF145	TSF215	TSF230	TSF245	TSF315	TSF330	TSF345
AOMD	77.49±0.52 ^a	76.41±1.18 ^{ab}	77.19±0.98 ^{ab}	75.42±1.19 ^b	77.05±1.73 ^{ab}	76.83±0.75 ^{an}	75.38±0.57 ^b	77.23±0.79 ^{ab}	75.58±0.83 ^{ab}	73.36±0.96 ^c
APD	92.09±0.80 ^a	91.56±0.91 ^a	91.21±0.89 ^a	87.87±0.70 ^b	91.82±1.04 ^a	91.32±0.50 ^a	87.77±0.37 ^b	91.87±0.45 ^a	91.02±0.65 ^a	86.80±1.31 ^b
ALD	88.54±0.97 ^{ab}	86.74±1.82 ^{bcd}	86.72±1.44 ^{bcd}	87.62±1.58 ^{abc}	87.02±0.76 ^{abcd}	85.23±0.22 ^d	86.07±0.72 ^{cd}	88.93±1.15 ^a	86.75±0.55 ^{bcd}	84.97±0.51 ^d
AED	82.59±0.49 ^{ab}	81.04±1.62 ^{abc}	81.75±1.00 ^{abc}	80.15±0.89 ^{cd}	82.08±1.60 ^{ab}	81.63±0.27 ^{abc}	80.06±0.67 ^{cd}	82.71±0.67 ^{ab}	80.73±0.85 ^{bc}	78.43±0.93 ^d
AAD	78.65±1.33 ^{ab}	81.65±5.41 ^a	77.99±0.38 ^b	76.33±1.11 ^{bc}	77.13±0.98 ^b	75.42±0.41 ^{bc}	75.41±1.91 ^{bc}	75.85±0.57 ^{bc}	75.36±0.21 ^{bc}	72.95±0.58 ^c
AF D	52.34±1.52 ^a	52.59±2.13 ^a	52.96±1.02 ^a	51.96±1.58 ^a	52.71±2.13 ^a	52.12±1.24 ^a	51.03±1.25 ^a	51.01±1.55 ^a	51.18±2.08 ^a	48.71±0.81 ^b
ACD	52.71±2.60	49.20±4.52	52.50±2.23	52.64±3.73	52.77±4.59	54.06±3.08	54.54±1.84	53.57±2.15	48.71±1.87	49.57±3.90

Row means with different superscripts are significantly different (p<0.05)

Row means with no superscripts are not significantly different (p>0.05)

AOMD Apparent Organic Matter Digestibility

APD Apparent Protein Digestibility

ALD Apparent Lipid Digestibility

AED Apparent Energy Digestibility

AAD

Apparent Ash Digestibility

AF D

Apparent Fibre Digestibility

ACD

Apparent Carbohydrate Digestibility

Table 7. Haematological profile of blood of *Clarias gariepinus* fed toasted sunflower meal based diets

	CTR	TSF115	TSF130	TSF145	TSF215	TSF230	TSF245	TSF315	TSF330	TSF345
Hb	10.20±0.10 ^a	10.05±0.05 ^a	10.01±0.3 ^a	9.39±0.21 ^b	10.07±0.01 ^a	9.82± 0.33 ^a	9.41±0.32 ^b	10.06±0.04 ^a	10.01±0.05 ^a	9.37 ± 0.21 ^b
PCV	31.66±0.24 ^a	29.21±0.19 ^b	29.03±0.23 ^{bc}	26.15±0.91 ^d	29.26±0.06 ^b	28.02±0.18 ^c	26.12±0.81 ^d	29.23±0.09 ^b	29.06 ± 0.26 ^{bc}	25.78±0.79 ^d
WBC	7150±70.71 ^c	7200±70.71 ^{abc}	7300±70.71 ^{abc}	7400±70.71 ^a	7175±35.36 ^{abc}	7225±10.61 ^{abc}	7385±91.92 ^{ab}	7200±11.31 ^{abc}	7315±91.92 ^{abc}	7340±127.27 ^{abc}
RBC	3.37 ± 0.04 ^a	3.30±0.01 ^a	3.27±0.10 ^a	3.07±0.08 ^b	3.28±0.10 ^a	3.25±0.03 ^a	3.01±0.04 ^b	3.28±0.13 ^a	3.24±0.04 ^b	3.02±0.04 ^b
MCHC	32.22±0.07 ^a	34.40±0.05 ^b	34.48±0.17 ^b	35.89±0.46 ^c	34.40±0.10 ^b	35.05±0.93 ^b	36.01±0.10 ^c	34.42±0.25 ^a	34.44±0.13 ^a	36.34±0.31 ^b
MCV	94.09±0.03 ^a	88.50±0.20 ^{ab}	88.78±1.07 ^{ab}	85.38±5.12 ^b	89.23±2.50 ^{ab}	86.22±1.32 ^b	86.90±1.69 ^b	89.18±0.25 ^a	89.82±0.18 ^a	85.38±3.80 ^b
MCH	30.31± 0.03	30.44±0.01	30.62±0.22	30.64±1.44	30.70±0.95	30.22±1.26	31.30±0.69	30.70±1.32	30.93±0.18	31.02±1.12
ESR	3.73±0.03 ^a	3.76±0.04 ^{ab}	3.82±0.01 ^{bc}	3.94±0.02 ^c	3.76±0.01 ^{ab}	3.83±0.04 ^{bc}	3.92±0.01 ^{de}	3.76±0.05 ^{ab}	3.86±0.05 ^{cd}	3.99±0.03 ^e
HSI	1.80±0.01 ^{ab}	1.79±0.02 ^{ab}	1.76±0.01 ^{bc}	1.73±0.01 ^c	1.82±0.04 ^a	1.76±0.03 ^{bc}	1.72±0.00 ^c	1.83±0.01 ^a	1.79±0.01 ^{ab}	1.72±0.04 ^c

Row means with different superscripts are significantly different (p<0.05), Row means with no superscripts are not significantly different (p>0.05), Hb: Haemoglobin content (gm/100ml) PCV: Packed Cell Volume (%), WBC: White Blood Cell Count (10⁴mm³), RBC: Red Blood Cell Count (10⁶mm³) MCHC: Mean Corpuscular Haemoglobin Concentration (%), MCV: Mean Corpuscular Volume (u³) MCH: Mean Corpuscular Haemoglobin (pg), ESR: Er.ythrocyte Sedimentation Rate (mm/hr).

MCV followed similar result trend as recorded above. However, reverse trend of results was recorded for MCHC; The blood of fish fed CTR was the lowest which is significantly different ($p < 0.05$) from the MCHC of the blood of the fish fed other experimental diets. No significant variation ($p > 0.05$) existed in the MCHC of fish fed TSF115, TSF215, TSF315, TSF330, TSF130 and TSF230. The MCHC of the blood of fish fed TSF145, TSF 245 and TSF345 also showed no significant variation ($p > 0.05$). Fish fed TSF345 had the highest MCHC. There was no significant variation ($p > 0.05$) in the mean cell haemoglobin (MCH) among fish fed various experimental diets. The erythrocyte sedimentation rate (ESR) of fish fed CTR was the lowest which was not significantly different ($p > 0.05$) from the ESR of the blood of fish fed TSF115, TSF215 and TSF315. So also, no significant variation existed in the ESR of the blood of fish fed TSF145, TSF245 and TSF345. The Hepatosomatic index (HSI) of fish fed TSF315 had the highest value while the lowest value of HSI was recorded in fish fed TSF245. There was no significant variation in the HSI of fish fed CTR, TSF215, TSF315 TSF115 and TSF330. Primary haematological parameters (Hb, PCV, RBC, WBC) reveal health status of animal (7). Feist and Longshaw (24) reported that blood profile can be altered by diet composition. The non-significant difference in the primary haematological parameters recorded among the fish fed soybean-based control diet and those fed differently toasted sunflower meal up to 30% replacement level in this study further elucidate that the physiological functioning of fish is not hampered by toasted sunflowers up to 30% replacement. The blood profile and health status of the fish is not compromised by toasted sunflower based diets. This results trend of reduced primary haematological parameters with increasing plant protein was also reported by Blom et al (10) and Rinchar et al (56) plausibly due to higher contents of anti-nutrients, poor quality amino acid contents and higher fibre content of the diet (75). The blood profile of *Clarias gariepinus* fell within the range of normal haematology of a healthy fish (19; 51; 54). By contrast, other researchers reported increase in

haematological parameters include but not limited to Barros et al (8) for channel catfish fed cotton seedmeal replacing soybean meal; El-Saidy and Gaber (16) for Nile Tilapia fed the same seedmeal; Akintayo et al (3) who fed the toasted sunflower seed meal to *Clarias gariepinus*, (38) who fed watermelon seedmeal to Nile tilapia; Yue and Zhou (74) who fed cotton seed meal to juvenile hybrid Tilapia.

Histology of the liver

Photomicrographs (5 μ m sectioned, H&E stained, x100 Magnification) of liver of *Clarias gariepinus* fed varying levels of differently toasted sunflower based diet is presented in Figure 2. No visible lesions were seen in the liver of fish fed diet CTR, TSF115, TSF130, TSF215, TSF230, TSM315. Severe hepatic necrosis, with massive fatty infiltration was observed in the liver of TSF145-fed fish. Diffuse vacuolar degeneration of hepatocytes and kupffer cell hyperplasia were observed in fish fed TSF245. Diffuse vacuolar degeneration of hepatocytes was observed in fish fed TSF330 while there was severe hepatic necrosis, with massive fatty infiltration TSF345. The pathological condition of severe hepatic necrosis, with massive fatty infiltration recorded in the liver of fish fed TSF145 and TSF 345 in this study revealed a degraded liver as classified by Martínez-Llorens et al (45) to be grade 3 liver with necrosis, pyknotic nuclei, hyaline cytoplasm and/or large sinusoid spaces. Diffuse vacuolar degeneration of hepatocytes and kupffer cell hyperplasia observed in the liver of fish fed TSF245 and TSF330 depicted a grade 2 liver associated with an intermediate condition of the liver possessing displaced nuclei. The non-visible lesion recorded among the liver of fish fed control diets and those fed differently toasted up to 30% replacement level for 5- and 10-minute toasted sunflower and 15% replacement level for 15-minute toasted sunflower aptly categorized as grade 1 liver described as normally shaped and sized hepatocytes with uncongested vessels and normal sinusoid organization based on semi-quantitative assessment criteria of Martínez-Llorens et al (45). The pathological condition of the liver of fish fed higher content of toasted sunflower meal could be due to incomplete destruction of anti-nutrients in the

seedmeal which increased with increasing sunflower meal in the diet (2; 12; 18). Nero et al (52) reported that liver being vulnerable to destruction owing to its function as xenobiotic

compounds detoxifier. The massive fatty infiltration of the hepatocytes is dietary lipid induced (28; 73).








Figure 2. Photomicrographs (5 μ m sectioned, H&E stained, x100 Magnification) of liver of *Clarias gariepinus* fed diet (a) CTR: No visible lesions seen; (b) TSF145: There is severe hepatic necrosis, with massive fatty infiltration. (c) TSF245: Diffuse vacuolar degeneration of hepatocytes, kupffer cell hyperplasia; (d) TSF330: There is diffuse vacuolar degeneration of hepatocytes; (e) TSF345: There is severe hepatic necrosis, with massive fatty infiltration

CONCLUSION

It is evident from the results of this study that dietary replacement of soybean meal by differently toasted sunflower seedmeal in *Clarias gariepinus* diet can be done up to 30% level using survival, growth, carcass composition, digestibility, haematology and histology of the liver as indices. The physiological condition was comparable with soybean-based control diet. At higher replacement level of 45% for the differently toasted sunflower meal, the fish physiological functioning was becoming compromised though survival was not affected, there was reduced growth and changes to the blood and histopathological parameters of the liver.

ACKNOWLEDGEMENT

The effort of Mr Musa Mustapha in the translation of abstract of this study into Arabic language is highly acknowledged and appreciated.

REFERENCES

1. Abbas K, I. Ahmed and H. Rehman. . 2005. Growth performance as influenced by partial replacement of fish meal with plant proteins in the diet of major carps. *Indus J. Biol. Sci* 2:219-26
2. Agius C, and R. Roberts . 2003. Melano-macrophage centres and their role in fish pathology. *Journal of fish diseases* 26:499-509
3. Akintayo I, S. Obasa W, Alegbeleye and A., Bangbose . 2008. Evaluation of toasted sunflower (*Helianthus annuus*) seed meal in the diets of African catfish (*Clarias gariepinus*) fingerlings. *Livestock Research for Rural Development* 20:28-46
4. Amerio M, C. Vignali L., Castelli L, Fiorentini and E. Tibaldi . 1998. . Vegetable protein sources, protein evaluation indices and 'ideal protein' of sea bream (*Sparus aurata*). *Rivista Italiana di Acquacoltura* 33 135–45
5. AOAC. 2010. Official Methods of Analysis. Association of Official Analytical Chemists, Washington DC. Gaithersburg, Maryland : AOAC International
6. Ayadi FY, KA Rosentrater and K, Muthukumarappan . 2012. Alternative protein sources for aquaculture feeds. *Journal of Aquaculture Feed Science and Nutrition* 4:1-26
7. Bahmani M, R. Kazemi and P. Donskaya . 2001. A comparative study of some hematological features in young reared sturgeons (*Acipenser persicus* and *Huso huso*). *Fish Physiology and Biochemistry* 24:135-40
8. Barros MM, C. Lim and PH. Klesius . 2002. Effect of soybean meal replacement by cottonseed meal and iron supplementation on growth, immune response and resistance of Channel Catfish (*Ictalurus punctatus*) to *Edwardsiella ictaluri* challenge. *Aquaculture* 207:263-79
9. Barrows FT, TG. Gaylord , WM. Sealey , MJ> Haas and RL. Stroup . 2008. Processing soybean meal for biodiesel production ; effect of a new processing method on growth performance of rainbow trout , *Oncorhynchus mykiss* ☆. *Aquaculture* 283:141-7
10. Blom J , K.J. Lee , J. Rinchar , K.Dabrowski, and J. Ottobre . 2001. Reproductive efficiency and maternal-offspring transfer of gossypol in rainbow trout (*Oncorhynchus mykiss*) fed diets containing cottonseed meal. *Journal of animal science* 79:1533-9
11. De la Higuera M, H.Akharbach , M. Hidalgo , J. Peragon J. Lupianez and M. Garcia-Gallego . 1999. Liver and white muscle protein turnover rates in the European eel (*Anguilla anguilla*): effects of dietary protein quality. *Aquaculture* 179:203-16
12. Ebrahimi M, NH. Daeman , CM. Chong A. Karami and V. Kumar et al. 2017. Comparing the effects of different dietary organic acids on the growth, intestinal short-chain fatty acids, and liver histopathology of red hybrid tilapia (*Oreochromis* sp.) and potential use of these as preservatives. *Fish physiology and biochemistry* 43:1195-207
13. El-Saidy D and M. Gaber . 2002. Evaluation of hulled sunflower meal as a dietary protein source for Nile tilapia, *Oreochromis niloticus* (L.), fingerlings. *Annals Agriculture Science* 40:831-41
14. El-Saidy DM and MM. Gaber . 2002. Complete replacement of fish meal by soybean meal with dietary L-lysine supplementation for Nile tilapia *Oreochromis niloticus* (L.) fingerlings. *Journal of the World Aquaculture Society* 33:297-306

15. El-Saidy DM and MM. Gabe. 2003. Replacement of fish meal with a mixture of different plant protein sources in juvenile Nile tilapia, *Oreochromis niloticus* (L.) diets. *Aquaculture research* 34:1119-27
16. El-Saidy DM and MM. Gaber . 2004. Use of cottonseed meal supplemented with iron for detoxification of gossypol as a total replacement of fish meal in Nile tilapia, *Oreochromis niloticus* (L.) diets. *Aquaculture Research* 35:859-65
17. Eusebio PS , R.M. Coloso and RE., Mamauag .. 2004. Apparent digestibility of selected ingredients in diets for juvenile grouper, *Epinephelus coioides* (Hamilton). *Aquaculture Research* 35:1261-9
18. Fadel AH,MS. Kamarudin , N. Romano ,M. Ebrahimi , CR. Saad and AA. Samsudin AA. 2017. Carob seed germ meal as a partial soybean meal replacement in the diets of red hybrid tilapia. *The Egyptian Journal of Aquatic Research* 43:337-43
19. Fagbenro O, C. Adedire , E. Owoseeni and E. Ayotunde . 1993. Studies on the biology and aquaculture potential of feral catfish *Heterobranchus bidorsalis* (Geoffroy St. Hilaire 1809)(Clariidae). *Tropical zoology* 6:67-79
20. Fagbenro OA,, EO. Adeparusi and WA. Jimoh . 2010. Effect of processing on the nutrient, mineral and antinutrients composition of sunflower (*Helianthus annuus*) seed meal. *International Journal of Chemistry* 20:165-70
21. Fagbenro OA, EO. Adeparusi and WA. Jimoh . 2013. Haematological profile of blood of African catfish (*Clarias gariepinus*, Burchell, 1822) fed sunflower and sesame meal based diets. *Journal of Fisheries and Aquatic Science* 8:80-6
22. FAO. 2018. The State of World Fisheries and Aquaculture 2018-Meeting the sustainable goals. Licence: CC BY-NC-SA 3.0 IGO
23. Fawole FJ, AA. Adeoye , LO.Tiamiyu ,KI. Ajala , SO. Obadara and IO. Ganiyu . 2020. Substituting fishmeal with *Hermetia illucens* in the diets of African catfish (*Clarias gariepinus*): Effects on growth, nutrient utilization, haemato-physiological response, and oxidative stress biomarker. *Aquaculture* 518:734849
24. Feist S and M. Longshaw . 2000. Myxosporidiosis of fish and the bryozoan link with proliferative kidney disease (PKD) of salmonids. *Fish Vet. J* 5:37-46
25. Fraga I, J. Galindo , R. Reyes , S. N, et al. 1996. Assessment of different protein sources for feeding white shrimp *Penaeus schmitti*. *Rev. Cub. Invest. Pesq.* 20:6-9
26. Francis G, HP. Makkar and K. Becker . 2001. Antinutritional factors present in plant-derived alternate fish feed ingredients and their effects in fish. *Aquaculture* 199:197-227
27. Garcia-Gallego M, H. Akharbach and M. De la Higuera . 1998. Use of protein sources alternative to fish meal in diets with amino acids supplementation for the European eel (*Anguilla anguilla*). *Animal science* 66:285-92
28. Gatta PP, L. Parma , I. Guarniero , L. Mandrioli and R. Sirri et al. 2011. Growth, feed utilization and liver histology of juvenile common sole (*Solea solea* L.) fed isoenergetic diets with increasing protein levels. *Aquaculture research* 42:313-21
29. Gaylord T, S. Rawles and IH.D. Gatlin .. 2004. Amino acid availability from animal, blended, and plant feedstuffs for hybrid striped bass (*Morone chrysops* × *M. saxatilis*). *Aquaculture Nutrition* 10:345-52
30. Gill N, DA. Higgs , BJ. Skura , M. Rowshandeli and BS. Dosanjh BS, et al. 2006. Nutritive value of partially dehulled and extruded sunflower meal for post-smolt Atlantic salmon (*Salmo salar* L.) in sea water. *Aquaculture Research* 37:1348-59
31. Giri SS, SK. Sahoo and SN, Mohanty SN. 2010. Replacement of by-catch fishmeal with dried chicken viscera meal in extruded feeds: effect on growth, nutrient utilisation and carcass composition of catfish *Clarias batrachus* (Linn.) fingerlings. *Aquaculture international* 18:539-44
32. González-Pérez S and JM. Vereijken .. 2007. Sunflower proteins: overview of their physicochemical, structural and functional properties. *Journal of the Science of Food and Agriculture* 87:2173-91
33. Hassaan MS, MA.Soltan . and AM. Abdel-Moez . 2015. Nutritive value of soybean meal after solid state fermentation with *Saccharomyces cerevisiae* for Nile

- tilapia, *Oreochromis niloticus*. Animal Feed Science and Technology 201:89-98
34. Hassaan MS, MA. Soltan EY., Mohammady , MA. Elashry , ER. El-Haroun ER and SJ. Davies . 2018. Growth and physiological responses of Nile tilapia, *Oreochromis niloticus* fed dietary fermented sunflower meal inoculated with *Saccharomyces cerevisiae* and *Bacillus subtilis*. Aquaculture 495:592-601
35. Hertrampf JW and F. Piedad-Pascual . 2012. Handbook on ingredients for aquaculture feeds. Springer Science & Business Media
36. Jackson A, B. Capper and A. Matty . 1982. Evaluation of some plant proteins in complete diets for the tilapia *Sarotherodon mossambicus*. Aquaculture 27:97-109
37. Jansman A and A. Van der Poel . 1993. Anti-nutritional factors in legumes seeds. Nutritional effects and (bio-) technological inactivation. Mededelingen-Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen. Universiteit Gent (Belgium)
38. Jimoh WA AA. Ayelaja FO., Ajasin FY, Okemakin , SA. Abdusalami and OF., Adekunle . 2015. Some haematological and biochemical profile of blood of Nile tilapia (*Oreochromis niloticus*) fed on diets containing watermelon (*Citrullus lanatus*) seedmeal. Bayero Journal of Pure and Applied Sciences 8:109–14
39. Jimoh WA, MS. Kamarudin MA Sulaiman MA and AB. Dauda . 2019. Assessment of prebiotic potentials in selected leaf meals of high dietary fibre on growth performance, body composition, nutrient utilization and amylase activities of a tropical commercial carp fingerlings. Aquaculture Research:1–11
40. Köprücü K and E. Sertel .2012. The effects of less-expensive plant protein sources replaced with soybean meal in the juvenile diet of grass carp (*Ctenopharyngodon idella*): growth, nutrient utilization and body composition. Aquaculture international 20:399-412
41. Lim C and D. Akiyama . 1992. Full-fat soybean meal utilization by fish. Asian Fisheries Science 5:181-97
42. Lim S , S. Choi S, X. Wang , K. Kim and I. Shin et al. 2004. Effects of dehulled soybean meal as a fish meal replacer in diets for fingerling and growing Korean rockfish *Sebastes schlegeli*. Aquaculture 231:457-68
43. Lozano NBS, S. AT. Vidal .,SN Martínez-Llorens, .JE Mérida . and Blanco et al. 2007. Growth and economic profit of gilthead sea bream (*Sparus aurata*, L.) fed sunflower meal. Aquaculture 272:528-34
44. Maina J, R. Beames D. Higgs , P. Mbugua G., Iwama and S. Kisia . 2003. Partial replacement of fishmeal with sunflower cake and corn oil in diets for tilapia *Oreochromis niloticus* (Linn): effect on whole body fatty acids. Aquaculture Research 34:601-8
45. Martínez-Llorens S, R. Baeza-Ariño ,S. Nogales-Mérida ,M. Jover-Cerdá and A. Tomás-Vidal . 2012. Carob seed germ meal as a partial substitute in gilthead sea bream (*Sparus aurata*) diets: amino acid retention, digestibility, gut and liver histology. Aquaculture 338:124-33
46. Martinez C. 1986. Advances in the substitution of fish meal and soybean meal by sunflower meal in diets of rainbow trout (*Salmo Gairdneri* L.). An. Inst. Cienc. Mar. Limnol. Univ. Auton. Mexico 13:345-52
47. McGoogan B. and HI. D. Gatlin . 1997. Effects of replacing fish meal with soybean meal in diets for red drum *Sciaenops ocellatus* and potential for palatability enhancement. Journal of the World Aquaculture Society 28:374-85
48. Mohanta KN , SN. Mohanty , J. Jena and SP. Sahu 2007. Effect of different oil cake sources on growth, nutrient retention and digestibility, muscle nucleic acid content, gut enzyme activities and whole-body composition in silver barb, *Puntius gonionotus* fingerlings. Aquaculture Research 38:1702-13
49. Mohanta KN, SN Mohanty J., Jena and NP. Sahu . 2008. Effect of three different oil cake-based diets on pond production performance of silver barb, *Puntius gonionotus* (Bleeker). Aquaculture research 39:1131-40
50. Morales A, G. Cardenete , A. Sanz and M. De la Higuera . 1999. Re-evaluation of crude fibre and acid-insoluble ash as inert markers, alternative to chromic oxide, in digestibility studies with rainbow trout (*Oncorhynchus mykiss*). Aquaculture 179:71-9
51. Myburgh JG, CJ. Botha , DG. Booysse and F. Reyers . 2008. Provisional clinical

- chemistry parameters in the African Sharptooth catfish (*Clarias gariepinus*). Journal of the South African Veterinary Association 79:156-60
52. Nero V, A. Farwell , A. Lister and Van Der Kraak G and L. Lee , et al. 2006. Gill and liver histopathological changes in yellow perch (*Perca flavescens*) and goldfish (*Carassius auratus*) exposed to oil sands process-affected water. Ecotoxicology and environmental safety 63:365-77
53. Ng WK and N. Romano . 2013. A review of the nutrition and feeding management of farmed tilapia throughout the culture cycle. Reviews in Aquaculture 5:220-54
54. Okore O. 2016. Growth and Haematological Studies of African Catfish (*Clarias gariepinus*) Juveniles Fed with Housefly Larva (*Musca domestica*) as Feed Supplement. International Journal of Agriculture and Earth Science 2:21-30
55. Olvera-Novoa MA, L. Olivera-Castillo and CA. Martínez-Palacios 2002. Sunflower seed meal as a protein source in diets for Tilapia rendalli (Boulanger, 1896) fingerlings. Aquaculture research 33:223-9
56. Rinchar J, K. Lee , S. Czesny , A. Ciereszko and K. Dabrowski . 2003. Effect of feeding cottonseed meal-containing diets to broodstock rainbow trout and their impact on the growth of their progenies. Aquaculture 227:77-87
57. Saha S. and K. Ghosh .2013. Evaluation of nutritive value of raw and fermented de-oiled physic nut, *Jatropha curcas* seed meal in the formulated diets for rohu, *Labeo rohita* (Hamilton) fingerlings. Proc. Proceedings of the zoological society, 2013, 66:41-50: Springer
58. Sales J, Britz P. 2002. Evaluation of the reference diet substitution method for determination of apparent nutrient digestibility coefficients of feed ingredients for South African abalone (*Haliotis midae* L.). Aquaculture 207:113-23
59. Sales J and P. Britz . 2003. Apparent and true availability of amino acids from common feed ingredients for South African abalone (*Haliotis midae* L.). Aquaculture Nutrition 9:55-64
60. Sánchez-Muros M-J and FG. Barroso and F,Manzano-Agugliaro F. 2014. Insect meal as renewable source of food for animal feeding: a review. Journal of Cleaner Production 65:16-27
61. Sanz A, AE. Morales , M. de la Higuera G, Cardente . 1994. Sunflower meal compared with soybean meal as partial substitutes for fish meal in rainbow trout (*Onorhynchus mykiss*) diets: protein and energy utilization. Aquaculture 128:287-300
62. Şener E, M. Yıldız and E. Savaş . 2006. Effect of vegetable protein and oil supplementation on growth performance and body composition of Russian sturgeon juveniles (*Acipenser gueldenstaedtii* Brandt, 1833) at low temperatures. Turkish Journal of Fisheries and Aquatic Sciences 6:23-7
63. Shipton T and P. Britz . 2000. Partial and total substitution of fishmeal with plant protein concentrates in formulated diets for the South African abalone, *Haliotis midae*. J. Shellfish Res 19:534
64. Shipton T and P. Britz . 2001. The partial and total replacement of fishmeal with selected plant protein sources in diets for the South African abalone, *Haliotis midae* L. Journal of shellfish Research 20:637-46
65. Sintayehu A . E. Mathies , KH.Meyer-Burgdorff , H. Rosenow and KD. Günther . 1996. Apparent digestibilities and growth experiments with tilapia (*Oreochromis niloticus*) fed soybean meal, cottonseed meal and sunflower seed meal. Journal of Applied Ichthyology 12:125-30
66. Stickney RR, RW. Hardy , K. Koch ,R. Harrold , D. Seawright and KC. Massee . 1996. The effects of substituting selected oilseed protein concentrates for fish meal in rainbow trout *Oncorhynchus mykiss* diets. Journal of the world aquaculture society 27:57-63
67. Storebakken T. 2000. Soy products as fat and protein sources in fish feeds for intensive aquaculture. Soy in animal nutrition:127-70
68. Sun W, X. Li , H. Xu , J. Chen , X. Xu and X. Leng . 2017. Effects of dietary chlorogenic acid on growth, flesh quality and serum biochemical indices of grass carp (*Ctenopharyngodon idella*). Aquaculture nutrition 23:1254-63

69. Tacon A, J. Webster and C. Martinez . 1984. Use of solvent extracted sunflower seed meal in complete diets for fingerling rainbow trout (*Salmo gairdneri* Richardson). *Aquaculture* 43:381-9
70. Tacon AG, MR. Hasan and M, Metian . 2011. Demand and supply of feed ingredients for farmed fish and crustaceans: trends and prospects. *FAO Fisheries and Aquaculture technical paper*:I
71. Tacon AG and M. Metian . 2008. Global overview on the use of fish meal and fish oil in industrially compounded aquafeeds: Trends and future prospects. *Aquaculture* 285:146-58
72. Tacon AG and M. Metian . 2015. Feed matters: satisfying the feed demand of aquaculture. *Reviews in Fisheries Science & Aquaculture* 23:1-10
73. Valente L, F. Linares , J. Villanueva, J. Silva and M. Espe . 2011. Dietary protein source or energy levels have no major impact on growth performance, nutrient utilisation or flesh fatty acids composition of market-sized Senegalese sole. *Aquaculture* 318:128-37
74. Yue Y-R, and Q.C.Zhou . 2008. Effect of replacing soybean meal with cottonseed meal on growth, feed utilization, and hematological indexes for juvenile hybrid tilapia, *Oreochromis niloticus* × *O. aureus*. *Aquaculture* 284:185-9
75. Zhou F, W. Song , Q. Shao , X. Peng and J. Xiao et al. 2011. Partial replacement of fish meal by fermented soybean meal in diets for black sea bream, *Acanthopagrus schlegelii*, juveniles. *Journal of the World Aquaculture Society* 42:184-97