

SEX DETERMINATION OF FETUS PRENATAL FROM MATERNAL PLASMA BY USING DUPLEX PCR TECHNIQUE IN GOAT

Talib Ahmed Jaayid*, Tahir A. Fahid** ,NisreenYasirJasim**

* Department of Animal Production, College of Agriculture , University of Basrah , Basrah, Iraq

** Department of Internal Medicine, Surgery and Obstetrics, College of Veterinary medicine, University of Basrah , Basrah, Iraq

(Received 18 September 2013 ,Accepted 23 September 2013)

Keyword; Fetal sex, Goat , Plasma, Duplex PCR

ABSTRACT

The present study included determination of fetal sex prenatal by analyzing free fetal DNA cells in maternal plasma by using duplex PCR . At the first amplifying the DNA which extracted from male and non-pregnant female goat plasma which used as control to confirm the accuracy of this technique. After that blood were collected from 5 pregnant females goat prepared the plasma and extract the DNA by using heat-based direct method for plasma and then investigate the sequences GAPDH and DYS14 through the use of duplex polymerase chain reaction technique and the results were 2 (40%) males show a double bands one at 97 bp representing GAPDH and the second at 198 bp representing DYS14 and 3 (60%) females show a single band at 97 bp representing GAPDH. This technique of sexing is applied for the first time on animals especially in Iraq.

INTRODUCTION

There is a simple and effective method for noninvasively detecting fetal sex using circulation fetal DNA from first trimester maternal plasma. Fetal sex determination is important in some endocrine diseases, such as congenital adrenal hyperplasia, where there is masculinization of the female fetus, which is preventable with antenatal treatment. Generally, early fetal sex determination has been performed by invasive procedures, such as chorionic villus sampling or amniocentesis. However, these invasive procedures still carry a 1 to 2% risk of miscarriage (1,2) and cannot be performed until 11 week of gestation. Moreover, reliable determination of fetal sex

using ultrasonography cannot be performed in the first trimester, because the development of external genitalia is not complete (3).

Circulating fetal DNA arises from trophoblasts in the placenta and is released directly into the circulation system of the mother (4, 5). It represents 3–6% of the total cell-free DNA that is present in maternal circulation and is rapidly cleared from the maternal circulation after delivery (6,7). Therefore, circulating fetal DNA has been proposed as a potential material for noninvasive prenatal diagnosis (NIPD). Many researchers have explored the possibility of circulating fetal DNA on NIPD of fetal sex, RhD blood typing, and single-gene disorders, such as cystic fibrosis (8, 9).

Promising results have been reported from studies using the combination of markers such as Detection Y-Specific gene (DYS14) presence of this gene indicates a male fetus and glyceraldehyde-3-phosphate dehydrogenase(GAPDH) sequence found in both sex (8, 9). (6) demonstrated that free fetal DNA (ffDNA) circulating in maternal blood has a relatively high concentration in early pregnancy, which increases during gestation and degrades in a few hours after delivery (7,10,11). Detection of ffDNA in maternal blood during pregnancy has given rise to the possibility of developing new noninvasive approaches for early prenatal diagnosis (12).

This approach seem to be more useful in human as compared with animals because it enable the diagnosis of the fetal gender in early gestation period this led to eliminate the risk of some disease such as Mendelian diseases, also prediction of RhD genotype and in addition to fetal gender (12, 13).

MATERIALS AND METHODS

1.Samples collection

Peripheral blood samples obtained from 5 healthy goat males were used to determine the PCR specificity and sensitivity, whereas peripheral blood samples obtained from 5 non pregnant goat female served as negative controls and also from 5 pregnant goat females use to determine the gender of fetus at agricultural research station in agriculture college, university of Basrah.

2.Plasma preparation

Plasma in this study was prepared according to (14):-

- 1) For plasma preparation, collect 2-3 ml into an EDTA-containing tube for plasma separation .
- 2) The blood samples were centrifuged at 3000 rpm for 10 min and the plasma was carefully removed from their tubes and transferred into plain polypropylene tubes.
- 3) The plasma samples then submitted a second centrifugation at 3000 rpm for 10 min and these recentrifuged plasma samples weretransferred into fresh polypropylene tubes.
- 4) The samples were stored at -20 °C until DNA extraction.


3.DNA extraction from plasma using the heat-based direct method

The DNA extracted from plasma using the heat-based direct method according to (15):-

- 1) 200 µl of plasma in a 0.5 µl sterile Eppendorf tube was heated at 99 C for 5 min.
- 2) The heated sample was then centrifuged at 12000 rpm for 10 minutes in a cooling microcentrifuge.
- 3) After which the clear supernatant was collected and 10 µl used for PCR reaction.

4.Measurement quantity of extracted DNA from plasma maternal using Nanodrop technique

The DNA quantity is measured by using Nanodrop technique (Thermo Scientific company/USA) as in figure (1) to detect genomic size (DNA quantity ng / µl).



Figure(1): Showing measurement DNA quantity ng/ μ using Nanodrop technique.

5. Amplification of DYS14 gene and GAPDH using duplex PCR

The DYS14 and GAPDH genes were studied according to protocol of (16). DYS14 and GAPDH genes were co-amplified in a duplex PCR setting for the detection of fetal DNA in maternal plasma. The PCR amplification mixture showed in table (1).

Table (1): Showing PCR product composition (μ l)

Chemical material	Master Mix	Primers	DNA Template	Nuclease free water	Final volume
Volume	25	2.5 of each forward and reverse primer	10	5	50

Table (2): Oligonucleotide primers sequences used for A duplex PCR amplification of DYS14 and GAPDH genes according to (16)

Primer name	Description and size of primer	sequence (5→3)	PCR product size (bp)
Y1.7	(21)	5' CATCCAGAGCGTCCCTGGCTT-3)	198
Y1.8	(21)	5' CTTTCCACAGCCACATTTGTC-3)	
GAPDH	forward primer (21)	5'CCCCACACACATGCACTTACC-3)	97
	Reverse primer (21)	5' CCTAGTCCCAGGGCTTTGATT-3)	

Table (3): A duplex PCR amplification program for DYS14 and GAPDH genes detection according to (16).

Step	Temperature (C°)	Time (Min.)	NO. of cycles
Initial denaturation	95	15	1
Denaturation	94	1	40
Annealing	57	1	
Extension	72	1	
Final extension	72	10	1

6.PCR result analysis

The results of PCR were performed after the amplification process. 5 µl from amplified sample was directly loaded in a 2% agarose gel that prepare containing 0.5 µl /25 ml ethidium bromide with the addition of loading buffer and DNA size marker as standard in electrophoresis and run at 80 V for 40 minutes then the products were visualized by gel documentation system(S 140, Votornix, USA).

RESULTS

The PCR analysis was applied for DNA extracted from plasma by heat – based direct method which characterized by using the extracted DNA directly in PCR reaction because when the extracted DNA delayed to use will not give results when amplification and the PCR reaction were failed. After DNA extraction method, the quantity of DNA measured by Nanodrop technique (figure 2 and3). The PCR were positive for both DYS14 and GAPDH sequences in DNA samples from all 5 healthy males. While in DNA samples extracted from non pregnant females the PCR were negative for the DYS14 sequence, but positive for GAPDH sequence in all 5 non pregnant females. The results of gender diagnosis were 2 (40%) male and 3 (60%) females as show in (table 4) from the DNA samples was extracted from 5 pregnant females (figure 4) .




Figure (2): Measurement quantity of DNA (ng/μl) from non pregnant female plasma using Nanodrop technique.




Figure (3): Measurement quantity of DNA (ng / μ l) from pregnant females plasma using Nanodrop technique.

Table (4): Results of amplification of DYS14 and GAPDH using duplex PCR for gender diagnosis from maternal plasma in pregnant females.

No . of pregnant goat females	Amplification of DYS14	Amplification of GAPDH	Gender detected
1	+	+	Male
2	+	+	Male
3	-	+	Female
4	-	+	Female
5	-	+	Female




Figure (4): Showing duplex PCR setup for DYS14 (198 bp) and GAPDH (97 bp) electrophoresis in 2% agarose. M Lane = DNA ladder, Lane 2 =male (has two bands one at 198 bp (DYS14) and other at 97 bp (GAPDH), Lane 3, 5, 6=female, has one band at 97 bp (GAPDH).

DISCUSSION

This study used to diagnosis the gender of embryo from pregnant female circulation this done by extraction the DNA from maternal plasma by using the heat – based direct method, this method characterized by using the extracted DNA directly in PCR reaction (15).

At the first amplified the DNA extracted from male and non pregnant female goat plasma which used as control to confirm the accuracy of this technique to detect the gender and the results were accordance with actual sex. In this study the technique applied on 5 pregnant goat female to detect the gender. The results were 2 (40%) male and 3 (60%) female. When the embryo are male the result of amplified were two bands, one 198 bp represent the DYS14 sequence, and other 97 bp represent the GAPDH sequence, while when embryo are female the results of PCR are one band at 97 bp represent GAPDH sequence. These results agreed with many investigators (14.,16, 17; 18). Finally have been concluded from this study that sex determination of fetus prenatal by analysis of ffDNA in maternal plasma is reliable technique but need more sterilized condition to avoid the mistake results.

تحديد جنس الجنين قبل الولادة من البلازما الأمومي باستخدام تقنية التفاعل التسلسلي للبلزمة المتعدد في المعز

طالب احمد جايد* ، طاهر عبدالحسين فهد** ، نسرين ياسر جاسم**

قسم الثروة الحيوانية، كلية الزراعة، جامعة البصرة، البصرة، العراق.

قسم الجراحة والتوليد، كلية الطب البيطري، جامعة البصرة، البصرة، العراق.**

الخلاصة

تضمنت الدراسة الحالية تحديد جنس الجنين قبل الولادة بتحليل خلايا ال DNA الجنينية المتحررة الى البلازما الامومي باستخدام التفاعل التسلسلي للبلزمة المتعدد في البداية تم تضخيم ال DNA المستخلص من بلازما اناث المعز الغير حامله وذكور المعز التي استخدمت ك سيطرة لتأكيد دقة التقنية. بعد ذلك تم جمع الدم من 5 اناث معز حوامل وتم تحضير البلازما واستخلاص ال DNA بطريقة التسخين المباشر للبلازما heat –based direct method ومن ثم التحري عن التتابع GAPDH و DYS14 من خلال استخدام تقنية تفاعل البلمرة المتعدد وكانت النتائج (40%) ذكور تظهر زوجين من الحزم الاول عند 97 زوج قاعدي يمثل GAPDH والثاني عند 198 زوج قاعدي يمثل DYS14 و(60%) اناث تظهر حزمة واحدة عند 97 زوج قاعدي يمثل GAPDH. تقنية التجنيس هذه لأول مرة تطبق على الحيوانات خصوصا في العراق.

REFERENCES

1. Tabor, A.; Philip, J.; Madsen, M.; Bang, J.; Obel, E.B. and Nørgaard-Pedersen, B. (1986). Randomised controlled trial of genetic amniocentesis in 4606 low-risk women. *Lancet*. **1**:1287–1293.
2. Mujezinovic, F. and Alfirevic, Z. (2007). Procedure-related complications of amniocentesis and chorionic villous sampling: a systematic review. *Obstet. Gynecol.*, **110**: 687–694.
3. Odeh, M.; Granin, V.; Kais, M.; Ophir, E. and Bornstein, J. (2009). Sonographic fetal sex determination. *Obstet. Gynecol. Surv.*, **64**:50–57.
4. Tjoa, M. L.; Cindrova-Davies, T.; Spasic-Boskovic, O.; Bianchi, D. W. and Burton, G.J. (2006). Trophoblastic oxidative stress and the release of cell-free fetoplacental DNA. *Am. J. Pathol.*, **169**: 400–404.
5. Alberry, M.; Maddocks, D.; Jones, M.; Abdel Hadi, M.; Abdel-Fattah, S.; Avent, N. and Soothill, P.W. (2007). Free fetal DNA in maternal plasma in anembryonic pregnancies: confirmation that the origin is the trophoblast. *Prenat. Diagn.*, **27**: 415–418.
6. Lo, Y. M.; Tein, M. S.; Lau, T. K.; Haines, C. J.; Leung, T. N.; Poon, P. M.; Wainscoat, J. S.; Johnson, P. J.; Chang, A. M. and Hjelm, N. M. (1998). Quantitative analysis of fetal DNA in maternal plasma and serum:

- implications for noninvasive prenatal diagnosis. *Am. J. Hum. Genet.*, **62**: 768–775.
7. **Lo, Y. M.; Zhang, J.; Leung, T. N.; Lau, T. K.; Chang, A. M. and Hjelm, N. M. (1999).** Rapid clearance of fetal DNA from maternal plasma. *Am. J. Hum. Genet.*, **64**: 218–224.
 8. **Wright, C.F. and Burton, H. (2009).** The use of cell-free fetal nucleic acids in maternal blood for non-invasive prenatal diagnosis. *Hum. Reprod. Update.* **15**: 139–151.
 9. **Bustamante-Aragones, A.; Gonzalez-Gonzalez, C.; de Alba, M. R.; Ainse, E. and Ramos, C. (2010).** Noninvasive prenatal diagnosis using cfDNA in maternal blood: state of the art. *Expert Rev. Mol. Diagn.*, **10**: 197–205.
 10. **Angert, R.M.; LeShane, E.S.; Lo, Y.M.; Chan, L.Y.; Delli-Bovi, L.C. and Bianchi, D.W. (2003).** Fetal cell-free plasma DNA concentrations in maternal blood are stable 24 hours after collection: analysis of first and thirdtrimester samples. *ClinChe.*, **49** (1): 195–198.
 11. **Benachi, A.; Steffann, J. and Gautier, E. (2003).** Fetal DNA in maternal serum: does it persist after pregnancy? *Hum Genet.*, **113** (1): 76–79.
 12. **Picchiassi, E.; Coata, G.; Fanetti, A.; Centra, M.; Pennacchi, L. and Di Renzo,G. (2008).** The best approach for early prediction of fetal gender byusing free fetal DNA from maternal plasma. Department of Obstetrics and Gynecology and Centre of Perinatal and Reproductive Medicine, University of Perugia, Perugia ,Italy. *PrenatDiagn.* **28**: 525–530.
 13. **Clausen, F.B.; Krog, G.R. and Rieneck, K. (2005).** Reliable test for prenatal prediction of fetal RhD type using maternal plasma from RhD negative women. *PrenatDiagn.*,**25**: 1040–1044.
 14. **Tungwiwat, W.; Fucharoen, G. and Ratanasiri, T .(2003).**Non-invasive fetalsex determination using a conventional nested PCR analysis of fetal DNA in maternal plasma. *ClinChimActa.*, **334**: 173-177.
 15. **Emanuel, S.L. and Pestka, S. (1993).** Amplification of specific gene products from human serum. *Genet Anal Tech Appl.*, **10**: 144-6.
 16. **Lale, N.S. T.; Tufan, A.; Babur, K.; Yildirim, B.; Semercü, C. and Baúci, H.(2005).** Analysis of cell-free fetal dna from maternal plasma and serum using a conventional multiplex PCR: factors influencing success. *Turk .J .Med .Sci.*, **35**: 85-92
 17. **Lo, Y.M.; Corbetta, N. and Chamberlain, P.F .(1997).**Presence of fetal DNA in maternal plasma and serum. *Lancet.* **350**: 485-7.
 18. **Houfflin-Debarge, V.; O’Donnell, H.and Overton, T.(2000).** High sensitivity of fetal DNA in plasma compared to serum and nucleated cells using unnested PCR in maternal blood. *Fetal DiagnTher.*, **15**: 102-7.