

Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
 Investigate Their Biological Activity

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

Synthesis a Series of 1,3,4-Thiadiazole Derivatives and Investigate Their
 Biological Activity

Waseela Abdul Reda*, Mustafa K. Shneshil*, Salih Mahdi Salman **

*Department of Chemistry, college of education for pure Sciences university of Diyala

** Department of chemistry, college of Medicine University of Diyala

Received 10 November 2014 ; Accepted 23 December 2014

Abstract

Eight compounds of new 1,3,4-thiadiazole biologically active for the series 1,4-di(5-(4-substitutedphenyldiazenyl)-1,3,4-thiadiazol-2-yl) benzene were prepared through the reaction of terephthalic acid with thiosemicarbazide .The product was coupled with substituted benzene through a diazotization reaction. The synthesized compounds (2-9) were identified using the analytical and spectral means. The biological activity for those compounds was investigated against three types of bacteria Escherichia coli (Gram negative bacteria), Enterobacter (Gram- negative bacteria) and Staphylococcus aureus (Gram-negative).

Keyword: Thiadiazol, biological active compounds, diazotization reaction

تحضير سلسلة جديدة من مشتقات 1,3,4 ثايدايازول والتحري عن فعاليتها البيولوجية

وسيلة عبد الرضا* ، مصطفى شنشل* ، صالح مهدي سلمان**

*قسم الكيمياء- كلية التربية للعلوم الصرفة- جامعة ديالى

**قسم الكيمياء- كلية الطب- جامعة ديالى

الخلاصة

تم تحضير ثمانية مركبات 1,3,4- ثايدايازول جديدة فعالة بيولوجيا تنتمي لسلسلة مركبات 1,4-di(5-(4-substitutedphenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene وذلك من خلال مفاعلة حامض التريفثاليك مع ثايوسيمي كاربازيد ثم تفاعل (diazotization) الناتج مع بنزين معوض. تم تشخيص المركبات المحضرة بالوسائل التحليلية والطيفية، وتم دراسة الفعالية البيولوجية لهذه المركبات (2-9) باستخدام وسط زرع لبعض انواع البكتيريا.

مفاتيح الكلمات : الثايدايازول ، المركبات الفعالة بيولوجيا، تفاعل الثايدايازيشن.

**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

Introduction

The resistance towards available drugs is rapidly becoming a major worldwide problem due to the conditioning of various virus and bacteria to current chemical used for treatments. The need to design new chemical compounds to overcome this problem has become one of the most important areas of research nowadays. Thiadiazole is a versatile moiety that exhibits a wide variety of biological activities. Thiadiazole moiety acts as a hydrogen binding domain and two – electrons donor system. It also acts as constrained pharmacophore. Many drugs containing thiadiazole nucleus are available in the market such as acetazolamide, methazolamide, sulfamehazole, etc.

It was shown that substituted 1,3,4-thiadiazoles exhibit antimicrobial [1] and antitubercular [2-4] activities , while other compounds act on the CNS as anticonvulsants [5-7] or as antidepressant and anxiolytic [8] agents. A family of selective 1,3,4-thiadiazoles phosphodiesterases inhibitors [9] , and selective orally active cyclooxygenase inhibitors were reported[10].

Moreover, many reports indicate that acylthiosemicarbazides and their corresponding cyclized 1,3,4 thiadiazole derivatives possess anti- inflammatory [11-13] and analgesic [14] activities. 1,3,4 – Thiadiazoles are thus a group of heterocycles whose derivatives are important in drugs industry, medicine and agriculture [13,15-21]. Accordingly, in continuation of our work in this area [22-27] , a variety of heterocyclic derivatives have been prepared from saccharide derivatives, involving some new thiadiazole, oxathiazoles , and their chemistry and effect of the derivatives on the enzyme tyrosinase was studied [28-31] , which is the rate limiting step in melanin biosynthesis [32] .

Members of this ring system have found their way into such diverse application as pharmaceuticals, oxidation inhibitors, cyanine dyes and metal complexing agents. The literature review showed that the thiadiazole nuclei have antimicrobial, anti-inflammatory, anticancer, anticonvulsant, antidepressant, antioxidant, radio protective, and anti-leishmanial activities [33-34].

The aim of this work is to synthesis new series of 1,3,4-thiadiazole which are expected to show biological activity.

Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

Experimental

Instrumental

The FTIR spectral data were recorded on FTIR-8300 Fourier Transform Infrared Spectrophotometer *SHIMADZU* using potassium bromide disc doublebeam UV-VISIBLE spectrophotometer. Melting points (°C) were recorded on hot stage Gallen Kamp melting point apparatus and were uncorrected. NMR spectra were recorded on Jeol and spectrometer at 400 MHz for ¹H.

The IR spectral data and the study of physical properties were done at the Department of Chemistry, College of Education for Pure Science, Diyala University.

1. Synthesis of 1,4-di(-2-amino-1,3,4-thiadiazol)-5-yl-benzene (1)

A mixture of appropriate terephthalic acid (0.01mol) and (1.82g, 0.02 mol) of thiosemicarbazide with (5mL) of phosphorus oxychloride was refluxed gently for (5 hrs.). After cooling (50mL) of water was added, the mixture was then refluxed for (7hrs.) and filtrated, neutralized with potassium hydroxide. The Precipitate was washed with water and recrystallized from (ethanol-water) to give compounds (1).

2. Synthesis of 1,4-di (5-(phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene series (2-9):

1,4-Di(2-amino-1,3,4-thiadiazol)-5-yl-benzene (0.01mole) was dissolved in slightly acidified distilled water, a mixture of (HCl and NaNO₂) (0.02 mole) was added and the reaction was carried in an ice bath. Then different substituted benzene (G: *p*-CH₃, *p*-OH, *p*-OCH₃, *p*-CH₂CH₃, *p*-isopropyl, *p*-tertiary butyl, *p*-n-propyl) (0.01 mole) were used to prepare the target molecules that were filtered and purified.


**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

Results and Discussion

1. Synthesis

Scheme 1. Synthesis scheme of 1,3,4-thiadiazoles derivatives



The precursor 1,4-di(-2-amino-1,3,4-thiadiazol)-5-yl-benzene (1) was synthesized by the reaction of terephthalic acid and thiosemicarbazide in presence of phosphorus oxychloride [35]. Compound (1) was characterized based on FTIR and ¹H-NMR, the IR spectrum shows two strong peaks at (3140-3215) cm⁻¹ related to the symmetric and asymmetric (-NH₂) group absorption vibration and peak at 1620 cm⁻¹ related to (C=N) of the 1,3,4-thiadiazole ring, while the ¹H-NMR spectrum was the chemical shift at 7.6 ppm for benzene ring and 8.2 ppm for (N-H) (figure 1).

**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman




Figure 1: ¹H NMR Spectrum for compound 1,4-di(5-(phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene [2]

The target compounds (2-9) 1,3,4-thiadiazoles derivatives were synthesized by diazotization reaction of 1,4-di(-2-amino-1,3,4-thiadiazol)-5-yl-benzene with benzene and other substituted benzene compounds (*p*-CH₃, *p*-OH, *p*-OCH₃, *p*-CH₂CH₃, *p*-isopropyl, *p*-tertiary butyl, *p*-*n*-propyl). The physical properties of the target molecules as shown in (table 2). The authenticity of the product was confirmed by spectral data (FTIR) shown in (table 1).

Table 1: The FTIR (KBr cm⁻¹) spectrum data (stretching vibrations) for the compounds (2-9).

Substituent G	Comp.	O-H	C-H aromatic	C-H aliphatic	C=N	N=N
H	2	-	3080	2955	1615	1600
<i>p</i> -CH ₃	3	-	3110	2987-3000	1610	1550
<i>p</i> -OH	4	3356	3112		1689	1560
<i>p</i> -OCH ₃	5	-	3024	2990-2866	1684	1580
<i>p</i> -C ₂ H ₅	6	-	3100	2980-2890	1643	1581
<i>p</i> - <i>i</i> -pro	7	-	3103	298-2885	1647	1555
<i>p</i> - <i>t</i> -Bu	8	-	3050	2995-2810	1644	1548
<i>p</i> - <i>n</i> -pro	9	-	3090	2985-2880	1656	1588

**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**
Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

Table 2: The physical properties of the compounds (2-9).

Com. No.	M.P. (°C)	% Yield	IUPAC Name
2	200	90	1,4-di(5-(phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene
3	220	85	1,4-di(5-(4-methyl phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene
4	300	87	1,4-di(5-(4-hydroxy phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene
5	198	80	1,4-di(5-(4-methoxy phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene
6	208	77	1,4-di(5-(4-ethyl phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene
7	210	79	1,4-di(5-(4-isopropyl phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene
8	209	78	1,4-di(5-(4-tert-butyl phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene
9	200	98	1,4-di(5-(4-n-propyl phenyldiazenyl)-1,3,4-thiadiazol-2-yl)benzene

2. Biological activity

The antibacterial activity results are given in figure (3-4). The synthesized 1,3,4-thiadiazoles derivatives exhibit various antibacterial activities which can make those compounds useful for the target application. The compounds [2-9] have been screened for their antibacterial activity by agar growth technique against three types of bacteria *Escherichia coli* (Gram negative bacteria), *Enterobacter* (Gram-negative bacteria) and *Staphylococcus aureus* (Gram-negative). Each compound was dissolved in DMSO to give a final concentration of 0.01 mg/ml. From the data obtained, all the compounds (3, 7, 8, and 9) have the highest activity against *E. coli* (G-ve) and *Enterobacter* (G-ve) than others, in case of compounds (2) and (5) show no activity against *E. coli* and *Enterobacter* while with *Staphylococcus aureus*, compounds (3, 4, 6, 7 and 9) shows the higher activity than others.

**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**
Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

Table 3: Biological Activity of compounds (2-9)

	2	3	4	5	6	7	8	9
Escherichia coli	+	+++	++	+	++	+++	+++	+++
Entrobacter (G⁻)	+	+++	+++	++	+++	+++	++	+++
Candida albicans	- - -	+++	++	+++	+++	++	+++	+++

Note:

- = Inactive

+ = slightly active inhibition zone <5 mm

++ = Moderately active inhibition zone <5 mm

+++ = Highly active inhibition >12 mm

All the compounds 3 , 7,8 and 9 were found to be highly active against Escherichia coli (G⁻) while the compounds (3 , 4 , 6 and 7) were found to be highly active against Entrobacter (G⁻) . But the compounds (3, 5,6,8 and 9) were found to be highly active against Candida albicans. But the compound (4) and (7) were found to be med active against Candida albicans.

References

1. Dogan, H.N.; Duran, A.; Rollas, S.; Sener, G.; Uysal, M.K.; Gulen, D. Synthesis of new 2,5-disubstituted-1,3,4-thiadi azoles and preliminary evaluation of anticonvulsant and antimicrobial activities. Bioorg. Med. Chem. Lett. 2002,10,2893-2898.
2. Mamolo, M.G.; Falagiani, V.; Zampieri, D.; Vio, L.; Banfi, E.; Scialino, G. Synthesis and antimycobacterial activity of (3,4-diaryl-3H-thiazol-2-ylidene)-hydrazide derivatives, Farmaco 2003, 58,631-637.
3. Foroumadi, A.; Kiani, Z.; Sonani, F. Antituberculosis agents VIII—Synthesis and in vitro antimycobacterial activity of alkyi a-[5-(5-nitro-2-thienyl)-1,3,4-thiadiazole-2-yltfo] acetates. Farmaco 2003, 58, 1073-1076.

**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

4. Oruc, E.E.; Rollas, S.; Kandemirli, F.; Shvets, N.; Dimoglo, A.S. 1,3,4-thiadiazole derivatives. Synthesis, structure elucidation, and structure-antituberculosis activity relationship investigation. *J. Med. Chem.* 2004, 47, 6760-6767.
5. Stillings, M.R.; Welbourn, A.P.; Walter, D.S. Substituted 1,3,4-thiadiazoles with anticonvulsant activity. 2. Aminoalkyl derivatives. *J. Med. Chem.* 1986, 29, 2280-2284.
6. Chapleo, C.B.; Myers, P.L.; Smith, A.C.; Tulloch, I.F.; Walter, D.S. Substituted 1,3,4-thiadiazole with anticonvulsant activity. *J. Med. Chem.* 1987, 30, 951-954.
7. Chimirri, A.; Grasso, S.; Monforte, A.M.; Zappala, M. Synthesis and anticonvulsant properties of 3-(1,3,4-thiadiazol-2-yl) imidazolidin-4-ones. *Farmaco* 1991, 46, 935-943.
8. Clerici, F.; Pocar, D.; Maddatena, G.; Loche, A.; Perlini, V.; Brufani, M. Synthesis of 2-amino-5-sulfanyl-1,3,4-thiadiazole derivatives and evaluation of their antidepressant and anxiolytic activity. *J. Med. Chem.* 2001, 44, 931-936.
9. Vergne, F.; Bernardelli, P.; Lorthiois, E.; Pham, N.; Proust, E.; Oliveira, C.; Mafroud, A.; Royer, F.; Wrigglesworth, R.; ScheHhaas, J.K.; et al, Discovery of thiadiazoles as a novel structural class of potent and selective PDE7 inhibitors. Part 1. design, synthesis and structure- activity relationship studies. *Bioorg. Med. Chem. Lett.* 2004, 14, 4607-4613.
10. Song, Y.; Connor, D.T.; Sercel, A.D.; Sorenson, R.J.; Doubleday, R. Synthesis, structure activity relationships, and in vivo evaluations of substituted di-tert-butylphenols as a novel class of potent, selective, and orally active cyclooxygenase-2 inhibitors, 2. 1,3,4- and 1,2,4-thiadiazole series. *J. Med. Chem.* 1999, 42, 1161-1169.
11. Labanauskas, L.; Kalacs, V.; Gaidelis, P.; Brukstus, A.; Dauksas, V. Synthesis of 3-(3,4-dimethoxy phenyl)-1H-1,2,4-triazole-5-thiol and 2-amino-3,4-dimethoxy phenyl)-1,3,4-thiadiazole derivatives exhibiting antiinflammatory Activity. *Pharmazie* 2001, 56, 617-619.

**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

12. Sahin, G.; Palaska, E.; Kelicen, P.; Demirdamar, R.; Altinok, G.A. Synthesis of some new 3-acyl thiosemicarbazides, 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazole-1-thiones and their anti-inflammatory activities. *Forsch Drug Res.* 2001, 51, 478-484.
13. Palaska, E.; Sahin, G.; Kelicen, P.; Durlu, N.T.; Altinok, G. Synthesis and anti-inflammatory activity of 1-acylthiosemicarbazides, 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazole-3- thiones. *II Farmaco* 2002, 57, 101-107.
14. Amir, M.; Shikha, K. Synthesis and antiinflammatory, analgesic, ulcerogenic and lipid peroxidation activities of some new 2-[(2,6-diehloroanilino)phenyl]acetic acid derivatives. *Eur. J. Med. Chem.* 2004, 39, 535-545.
15. Mishra, L.; Singh, V.K.; Dubey, N.K.; Mishra, A.K. Synthesis and fungicidal activity of some 5- membered heterocyclic derivatives containing benzimidazoles. *Biosci. Biotechnol. Biochem.* 1993,57,989-991.
16. Dogan, H.N.; Rollas, S.; Erdeniz, H. Synthesis, structure elucidation and antimicrobial activity of some 3-hydroxy-2- naphthoic acid hydrazide derivatives. *II Farmaco* 1998, 53,462-467.
17. Mamolo, M.G.; Vio, L.; Banfi, E. Synthesis and antimicrobial activity of some 2,5-disubstituted 1,3,4-thiadiazole derivatives, *II Farmaco* 1996, 51, 1-74.
18. Chufan, E.E.; Pedregosa, J.C.; Baldini, O.N.; Bruno-Blanch, L. Anticonvulsant activity of analogues of acetazolamide, *U Farmaco* 1999, 54, 838-841.
19. Krutovskikh, G.N.; Rusanov, A.M.; Gomaeva, G.F.; Vartanyan, L.P.; Kolesova, M.B. Radioprotective action of thiadiazole derivatives. *Pharm. Chem. J. (Engl. Transl.)* 1977, II, 484-488; *Kim.-Farm. Zh.* 1977, 11,48-52.
20. Chou, J.Y.; Lai, S.Y.; Pan, SJL.; Jow, G.M.; Chem, J.W.; Guh, J.H. Investigation of anticancer mechanism of thiadiazole- based compound in human nonsmall cell lung cancer A549 cells. *Biochem. Pharmacol.* 2003, 66, 115-124.
21. Oleson, J.J.; Sloboda, A.; Troy, W.P.; Halliday, S.L.; Landes, MJ.; Angfer, R.B.; Semb, J.; Cyr, K.; Williams, J.H. The carcinostatic activity of some 2-amino-1,3,4-thiadiazoles. *J. Am. Chem. Sac.* 1955,77,6713-6714.

**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

22. El-Sadek, M.M.; Mostafa, M.A.; Abdel Rahman, M.M.; Zagzoug, N.B. Reactions of periodate oxidized methyl - 4,6-O-benzylidene- α -D-glucopyranoside with hydrazines. Bratislava Symposia on Saccharides, Smollenice, Czechoslovakia 1984,67.
23. El-Sadek, M.M.; Faidailah, H.M.; Hassan, S.Y. Structure and reactions of 3-benzoyl-2-methyl-5- (D-arabino- tetrahydroxybutyl)pyrrole. Carbohydr. Res. 1990, 199,248-254.
24. Hassan, S.Y.; Faidailah, H.M.; El Massry, A.; El Sadek. M.M. Synthesis and reactions of 5-(D-aroAmo-tetrahydroxybutyl)-3-(2,3-dihydro-1,3,4-oxadiazole-2-thion-5-yl)-2-methylfuran and 5-(Z)-ara6wo-tetrahydroxybutyl)-3-(2substituted ammo-1,3,4-oxadiazol-5-yl)-2-methyl furan. Carbohydr. Res. 1997, 298,123-126.
25. El-Sadek, M.M.; El Socary, N.N. Synthesis of N-[1-acetyl-5- (d-arab } no-tetraacetoxybutyl) -2- methyl-3-pyrroyl]-N-phthalimidoacetamide. Carbohydr. Res. 1991,222,267-269.
26. El-Sadek, M.M.; Abdel-baky, S.A.; El Socary, N.N. Synthesis and reactions of 2-methyl 5-(D- arabifio-tetTahydroxy butyl)-3-pyiTolecarbohydrazide. Carbohydr. Res. 1992,223, 311-319.
27. El-Sadek, M.M.; Zagzoug, N.B.; El Socary, N.N. Reactions of 2-methyl-5-(D-arabirw tetrahydroxybutyl)-3-furoylhydrazine. Carbohydr. Res. 1993, 250, 323-326.
28. Barton, D.E.; Kwon, B.S.; Francke, U. Human tyrosinase gene, mapped to chromosome 11 (q14 \rightarrow q21), defines second region ofhomology with mouse chromosome 7. Genomics 1988, 3, 17-24.
29. Getting, W.S.; Fryer, J.P.; Shriram, S.; King, R.A. Oculocutaneous albinism type 1: The last 100 years. Pigment Cell Res. 2003, 16, 307-311.
30. Potterf, S.B.; Hearing, V.J. Tyrosine transport into melanosomes is increased following stimulation ofmelanocyte differentiation. Biochem. Biophys. Res. Commun. 1998, 248,795-800.
31. Borden, E.C. Melanoma: Biologically Targeted Therapeutics; Humana Press: New York, NY, USA, 2002.

**Synthesis a Series of 1,3,4-Thiadiazole Derivatives and
Investigate Their Biological Activity**

Waseela Abdul Reda , Mustafa K. Shneshil and Salih Mahdi Salman

32. Arias-Carrion, O.; Poppel, E. Dopamine, learning, and reward-seeking behavior. *Acta Neurobiol. Exp.* 2007, 67,481-488.
33. Seo, S.Y.; Sharma, V.K.; Sharma, N.J. Mushroom Tyrosinase: Recent Prospects. *Agric. Food Chem.* 2003, 51, 2837-2853.
34. Gonzalez, G. Reactions of mono saccharides with p-keto-esters. *Adv. Carbohydr. Chem.* 1956, 11, 97-143.
35. Rou'il I. H., Synthesis and Characterization of some Novel Hetrocyclic Liquid Crystalline Compounds and Polymer, PhD thesis, 2006.

