

Synthesis of Several New Schiff Bases Linked to Sulfonamido Naphthalimide Moiety with Expected Biological Activity.

Ahlam Marouf Al-Azzawi* Ahmed Solaiman Hamed**

***University of Baghdad - College of Science**

****University of Anbar- College of Veterinary Medicine**

Abstract

Several new Schiff bases linked to sulfonamidonaphthalimide moiety have been synthesized. The first step in this work involved introducing of 4-(N-naphthalimidyl) phenyl sulfonylchloride in amination reaction with hydrazine hydrate producing 4-(N-naphthalimidyl) phenyl sulfonylhydrazine. Introducing of the prepared sulfonyl hydrazine in condensation reaction with different substituted aromatic aldehydes in the second step afforded the target new Schiff bases. Structures of the prepared compounds were elucidated on the basis of FTIR, ¹HNMR, ¹³CNMR spectral data which agreed with the proposed structures. The newly synthesized compounds are expected to have biological activity since they are built from biologically active components including naphthalimide, sulfonylimide and Schiff base.

Key Words; Naphthalimide, Sulfon amide, 4-(N-naphthalimidyl) phenyl , Sulfonylhydrazine

Introduction

Schiff bases which contain azomethin group attract much interest in synthetic chemistry (1,2). They are used as substrate in the preparation of industrial and biologically active compounds .Moreover they are also known to have biological activities such as antibacterial, antifungal, anti-tumor and antioxidant activities (3-6).

On the other side naphthalimides first discovered by Brana and coworkers have been found to exhibit diverse biological activities, some of them have shown high anticancer activity against a variety of murine and human tumor cells while others have shown analgesic properties (7-9).

Besides sulfonylimide drugs were the first effective chemotherapeutic agents to be employed systemically for the prevention and cure of bacterial infection in human beings (10,11).

Encouraged by all these observations it was thought worthwhile to synthesize new Schiff bases linked to naphthalimide through sulfonylimido group with expected biological activity.

NATERIALS AND METHODS

Melting points were determined on Thomas Hoover apparatus and were uncorrected. FTIR spectra were recorded on SHIMADZU FTIR-8400 Fourier Transform Infrared spectro-photometer using KBr discs. ¹HNMR and ¹³CNMR spectra were recorded on Bruker 300MHz instrument in Al-Albate University in Jordan using DMSO-d₆ as a solvent and trimethylsilane (TMS) as internal reference.

Synthesis of 4-(N-naphthalimidyl) phenyl sulfonylhydrazine [1]

To a solution of (0.01 mol, 3.71g) of 4-(N-naphthalimidyl)phenyl sulfonyl-chloride in (15 mL) of absolute ethanol , (0.01 mol) of hydrazine hydrate was added dropwise with stirring and keeping temperature at zero C° (12) .

The resulted mixture was refluxed for six hours then cooled to room temperature before pouring on crushed ice with stirring . The resulted precipitate was filtered , washed with cold water, dried and finally recrystallized from ethanol.

synthesis of Schiff bases [2-10]

A mixture of 4-(N-naphthalimidyl) phenyl sulfonylhydrazine (0.01mol , 3.67g) of aromatic aldehyde (0.01 mol) and (2-3) drops of glacial acetic acid in absolute ethanol (20 mL) was refluxed for 6 hours (13).


After cooling the obtained precipitate was filtered then washed with cold ethanol , dried and recrystallized from a suitable solvent.

Physical properties of Schiff bases [2-10] are listed in Table (1).

RESULT AND DISSCUTION

Since both naphthalimides and Schiff bases belong to a widely used group intermediates important for production of many types of pharmaceuticals and have wide spectrum of biological applications the target of the present work is synthesis of new molecules containing these two active moieties naphthalimide and Schiff base linked together through phenyl sulfonamide component.

Steps for performing this target are shown in scheme (1). (1).



The first step involved preparation of 4-(N-naphthalimidyl)phenylsulfonylhydrazine [1] via reaction of 4-(N-naphthalimidyl)phenylsulfonylchloride with hydrazine hydrate under reflux conditions.

The starting compound 4-(N-naphthalimidyl)phenylsulfonylchloride was prepared via reaction of N-phenyl-naphthalimide with chlorosulfonic acid according to literatures (14).

Compound [1] was purified by recrystallization from ethanol and was afforded as off-white crystals in (87%) percent yield and having melting point ($>300^{\circ}\text{C}$).

In the second step compound [1] was introduced in condensation reaction with different aromatic aldehydes producing the desired Schiff bases [2-10].

Physical properties of Schiff bases [2-10] are listed in Table (2).

Structures of the prepared compounds were confirmed by FTIR, ^1H NMR and ^{13}C NMR spectra data.

FTIR spectrum of compound [1] showed clear absorption bands at 3417 cm^{-1} and 3309 cm^{-1} due to asym. and sym. $\nu(\text{NH}_2)$. Absorption bands belong to $\nu(\text{N-H})$ amide, $\nu(\text{C-H})$ aromatic and $\nu(\text{C=O})$ imide appeared at 3236 cm^{-1} and 3062 cm^{-1} and 1705 cm^{-1} while absorption bands due to $\nu(\text{C=C})$ aromatic, $\nu(\text{C-N})$ imide, asym. $\nu(\text{SO}_2)$ and sym. $\nu(\text{SO}_2)$ appeared at 1620 cm^{-1} , 1361 cm^{-1} , 1381 cm^{-1} and 1180 cm^{-1} respectively (15).

^1H NMR spectrum of compound [1] showed clear signal at $\delta = 2.27\text{ ppm}$ belong to (N-H) amide proton, signals at $\delta = (7.38-7.53)\text{ ppm}$ belong to phenyl ring

protons, signals at $\delta = (7.84-8.51)$ ppm belong to naphthyl ring protons and signal at $\delta = (8.71)$ ppm belong to (NH₂) protons.

¹³CNMR spectrum of compound [1] showed signals at $\delta = (122.14-134.95)$ ppm belong to aromatic carbons and signal at $\delta = (160.95)$ ppm belong to (C=O) imide carbons.

FTIR spectra of prepared Schiff bases [2-10] showed clear absorption bands at (3230-3260) cm⁻¹, (3040-3113) cm⁻¹ and (1701-1740) cm⁻¹ which are belong to ν (N-H) amide, ν (C-H) aromatic and ν (C=O) imide respectively (15).

Other absorption bands were appeared at (1660-1680) cm⁻¹, (1585-1596) cm⁻¹ and (1290-1350) cm⁻¹ belong to ν (C=N) imide, ν (C=C) aromatic and ν (C-N) imide respectively while absorption bands due to asym. ν (SO₂) and sym. ν (SO₂) were appeared at (1342-1390) cm⁻¹ and (1178-1192)cm⁻¹ respectively (15).

FTIR spectrum of compound [3] showed clear absorption bands at (1087) cm⁻¹ belong to ν (C-Cl), while FTIR spectra of compound [4] and [5] showed absorption bands at (1508, 1523)cm⁻¹ and (1435, 1438)cm⁻¹ belong to ν (C-NO₂) and FTIR spectra of compounds [7] and [9] showed absorption bands at (3417, 3411) cm⁻¹ belong to ν (OH). All details of FTIR spectral data of Schiff bases [2-10] are listed in Table (2).

¹HNMR spectrum of compound [3] showed clear signal at $\delta = 5.8$ ppm belong to (N-H) amide proton, signals at $\delta = (7.65-8)$ ppm belong to phenyl ring protons, signals at $\delta = (8.43-8.7)$ ppm belong to naphthyl ring protons and signal at $\delta = (8.79)$ ppm belong to imine proton.

¹³CNMR spectrum of compound [3] showed signals at $\delta = (122.13-137.79)$ ppm belong to aromatic carbons present in two phenyl rings and naphthyl ring. The spectrum showed also signals at $\delta = (160.73-161)$ ppm belong to (C=N) carbon and signal at $\delta = (171.16)$ ppm belong to (C=O) imide carbon.

¹HNMR spectrum of compound [5] showed signal at $\delta = 5.79$ ppm belong to (N-H) amide proton, signals at $\delta = (7.38-7.89)$ ppm belong to phenyl rings protons, signals at $\delta = (8.29-8.54)$ ppm belong to aromatic proton in naphthyl rings and signal at $\delta = (8.69)$ ppm belong to imine proton.

¹³CNMR spectrum of compound [5] showed signals at $\delta = (111.91-134.98)$ ppm belong to aromatic carbons present in two phenyl rings and naphthyl ring, signal at $\delta = (160.0)$ ppm belong to (C=N) carbon and signal at $\delta = (172.2)$ ppm belong to (C=O) imide carbon.

¹HNMR spectrum of compound [6] showed two clear signals at ($\delta = 2.98$ ppm and $\delta = 3.04$ ppm) belong to two methyl groups protons and signal at $\delta = (5.80)$ ppm belong to (N-H) amide proton. Signals appeared at $\delta = (7.62-8.45)$ ppm are belong to aromatic protons in two phenyl ring and naphthyl rings while the signal appeared at $\delta = (8.5)$ ppm is belong to imine proton.

¹³CNMR spectrum of compound [6] showed two signals at $\delta = (19.3$ and $24.64)$ ppm belong to carbons of two methyl group and signals at $\delta = (118.13-148.14)$ ppm are belong to aromatic carbons of two phenyl rings and naphthyl rings. Other signals appeared at $\delta = (159.5-160.42)$ ppm and at $\delta = (179.66)$ ppm which are belong to (C=N) and (N=O) imide carbons respectively.





CONCLUSION

A series of new Schiff bases containing two biologically active components was synthesized successfully by performing two steps synthesis. The newly synthesized compounds were expected to possess high biological activity since they contain three known biologically active moieties cyclic imide, sulfonamide and Schiff base.

ACKNOWLEDGMENT

Authors are thankful to all who help in this work or share their knowledge.

Table (1) physical properties of Schiff bases [2-10]

Comp.No.	Compound structure	Yield%	Color	Melting point C ^o	Recrystall. solvent
2		87	yellow	232-233	Acetone
3		85	Light brown	183-185	Acetone
4		78	yellow	207-209	Ethanol
5		75	yellow	210-212	Ethanol

6		87	Red	239-241	Acetone
7		72	Brown	235-237	Dioxane
8		85	Brown	202-204	Acetone
9		80	Redish Brown	205-207	Dioxane
10		70	Deep yellow	223-225	cyclohexane

Table (2) FTIR spectral data (cm⁻¹) of Schiff bases [2-10]

CompN o.	v(NH) Amide	v(C-H) Aromatic	v(C=O) Imide	v(C=N) Imide	v(C=C) aromatic	Asym.vSO ₂ Sym. vSO ₂	v(C-N) Imide	Others
2	3230	3040	Asym.1740 Sym.1700	1670	1590	1390 1185	1315	-
3	3236	3066	1701	1662	1585	1381 1188	1330	v(C-Cl) 1087
4	3260	3040	Asym.1725 Sym.1700	1680	1590	1385 1185	1346	v(C-NO ₂) 1508 1435
5	3236	3078	1705	1674	1589	1342 1188	1290	v(C-NO ₂) 1523 1438
6	3255	3103	Asym.1740 Sym.1700	1660	1596	1363 1178	1332	v(C-H) Aliphatic 2912
7	3234	3113	1705	1674	1593	1346 1192	1292	v(O-H) 3417
8	3260	3060	1708	1668	1587	1375 1188	1350	-
9	3238	3110	1708	1675	1588	1348 1190	1295	v(O-H) 3411
10	3255	3105	1708	1665	1590	1345 1189	1345	-

REFERENCES

- [1] Chandra K.B. and Kaushik A.J., (2012) pharmacological examination and synthesis of some Schiff bases and thiazolidinone derivatives of 5-amino-1H-imidazole-4-carboxamide, *Der pharm chemical*, 4(5) :1873-1878.
- [2] Kumar P.P. and Rani B.L., (2011) Synthesis and characterization of new Schiff bases containing pyridine moiety and their derivatives as antioxidant agent, *Int. J. chem. Tech. Res.*, 3(1): 155-160.
- [3] Abdel-Salam F. H., (2010) Biological study and complexation behavior of some anionic Schiff base amphiphiles, *J. Surfact. Deterg.*, 13:423-431
- [4] Antia S. and Manish K. S., (2013) Synthesis, characterization and biological activity of Schiff bases derived from 3-(4-substituted)-1-phenyl-

- 1H-pyrazol-4-carbaldehyde and o-aminophenol, Chem. Sci. Trans., 2(3): 871-876.
- [5] Satyanarayana V. S., Sreevani P., Sivakumar A. and Vijayakumar V., (2008) Synthesis and
- [6] Sathe B. S., Jayachandran E. and Jagrap V. A., (2011) Antihelminthic activity of newly synthesized moieties of fluorobenzothiazole Schiff's base, Res. J. Pharm. Bio. And Chm. Sci., 2: 510-515.
- [7] Aibin W., Yufang X. and Xuhong Q., (2010) Novel naphthalimide-indomethacin hybrids as potential antitumor agents, Monatish chem. 141: 393-899.
- [8] Andricopulo A. D., Yunes R. A., Filho V. C., Correa R., Willain A., Santos A. R.S. and Nunes R. J., (1998) Synthesis and analgesic properties of cyclic imides, naphthalimide and bisnaphthalimide derivatives, Acta Farm. Bon., 17 :219-224.
- [9] Tati Jana M., Tina M., Celine B., Nancy D. and Janique D., (2008) UNBS 5162 a novel naphthalimide that decreases CXCL chemokine expression in experimental prostate cancer. Neoplasia, 10 :573-586.
- [10] Santosh K., Niranjan M. S., Chaluvaraju K. C., Jamakhandi C. M. and Dayanand K., (2010) synthesis and antimicrobial study of some Schiff bases of sulfonamides, J. Current Pharm. Res., 1 :39-42.
- antimicrobial activity of new Schiff bases containing comarin moiety and there spectral characterization, ARKIVOC, xvii :221-223.
- [11] Sreenivasa G. M., Jayachandran E., Shivakumar B., Jayaraj K. K. and Vijikumar M. M., (2009) Synthesis of bioactive molecule chlorobenzothiazole compressing potent heterocyclic moieties for anthelmintic activity, Arch. Pharm. Sci. and Res., 1 (2) : 150-157.
- [12] Al-Azzawi A. M. and Hmd A. S., (2014) Synthesis and antimicrobial screening of new naphthalimides linked to oxadiazole, thiadiazole and triazole cycles. Int. Res. Pharm. Chem., 4(2) :283-290.
- [13] Al-Azzawi A. M. and Abd Al-Razzak M. S., (2013) Synthesis, characterization and antimicrobial screening of new Schiff bases linked to phthalimide, Int. Res. Pharm. Chem., 3 (3) :682-690.
- [14] Al-Azzawi A. M. and Abd Al-Razzak M. S., (2011) Synthesis, characterization and antimicrobial activity of several new Schiff bases linked to phthalimide moiety, Kerbala J. Pharm. Sci., 2:124-133.
- [15] Silverstein R. M., Basler G. C. and Kiemel D. J., 7th Ed. John Wiley and Sons, N. J. (2005).

تحضير عدد من قواعد شيف الجديدة المرتبطة بمكونة سلفون

اميدونفتال ايميد ذات فعالية بايولوجية متوقعة

احلام معروف العزاوي احمد سليمان حمد

E-mail : Chem_ah_67@yahoo.com

الخلاصة

تم في هذا البحث تحضير عدد من قواعد شيف الجديدة المرتبطة بمكونة سلفون اميدونفتال ايميد. تضمنت الخطوة الاولى من التحضير ادخال المركب 4-N (نفتال ايميديل) فنيل كلوريد السلفونيل في تفاعل مع الهيدرازين وبذلك تم الحصول على المركب 4-N (نفتال ايميديل) فنيل سلفونيل هيدرازين. اما الخطوة الثانية فقد تضمنت ادخال المركب المحضر 4-N (نفتال ايميديل) فنيل سلفونيل هيدرازين في تفاعل تكاثف مع الديهايدات اروماتية مختلفة لانتاج قواعد شيف الجديدة. تم اثبات صحة تركيب المركبات المحضرة من خلال الاعتماد على مطيافية الاشعة تحت الحمراء والرنين النووي المغناطيسي بنوعيه $^{13}\text{CNMR}$, $^1\text{HNMR}$ حيث كانت النتائج المستحصلة مطابقة للتركيب المقترحة. من المتوقع ان تكون قواعد شيف الجديدة ذات فعالية بايولوجية سيما وان جزيئاتها قد بنيت اساسا من مكونات فعالة بايولوجيا وهي النفثال ايميد والسلفون اميد وقواعد شيف.