

## Detection of Extended Spectrum $\beta$ -Lactamase Gene CTX-M-1 in *Escherchia coli* and *Klebseilla pneumonia* Isolated from Urinary Tract Infection of Pregnant Women in Al-Nassyriah City

Nasser Saleh Lhwak \*

Yahya A.Abbas \*\*

\* Department Of Biology- College Of Science- University Of Thi-Qar

\*\* AL- Nassyriah Technical Institute- Southern Technical University

### Abstract:

This study was performed to detect the presence of CTX-M-1 gene in *Klebseilla pneumonia* and *Escherchia coli* are that causing urinary tract infections in pregnant women in Al- Nassyriah city , during the period from september (2016) to february (2017).

Three hundred and thirty (n= 330) urine samples were collected and cultured on appropriate media for initial isolation. Morphological, conventional biochemical tests and API 20 system were performed to identify the pathological bacteria agents. Ninety samples showed positive culture, the most common isolates were *E. coli* 57 (63.3%) followed by *K. pneumonia* 21(23.3%) and 12(13.3%) from other Gram negative bacteria.

ESBLs gene CTX-M-1(688 bp) was found in 22 isolates 17/57 (29.8%) in *E.coli* and 5/21 (23.8 %) in *K. pneumonia*.

**KEYWORD:** ESBLs, *E.coli*, *K.pneumoniae*, Pregnant Women ,UTI.

### Introduction:

Urinary tract infection (UTI) is an infections caused by the existence and growth of microorganisms anywhere in the urinary tract. It is usually due to bacteria from the tract of digestive which ,climbs to the opening of the urethra and begin to double to cause infection (Okonko *et al.*, 2009). In dissimilarity to men, women are more susceptible to UTI, and this is mainly due to absence of prostatic secretion, shortened urethra, pregnancy and easy contamination of the urinary tract with fecal flora (Haider *et al.*, 2010).

*Escherichia coli* and *Klebsiella pneumonia* were of the common intestinal commensals normal flora , it's a members of enterobacteriaceae family, with importance role in diverse hospital infections and community acquired infections. These bacteria dominant in urinary tract infections. Frequently use of antibiotics to remedy *E. coli* and *K. pneumonia* infectious lead to increasing the multidrug resistance

(MDR) and hence the problem in treating the infection specially UTIs (Trecarichi *et al.*, 2012).

$\beta$ -lactamase production considered as an important mechanism between several model to resistance of antibiotic by *E.coli* and *K. pneumonia* (Nicolas-Chanoine *et al.*, 2012).The ability to production ESBLs is contributors in the trouble of resistance to antimicrobial(Mirzaee *et al.*, 2009). Cefotaxim hydrolyzed (CTX-M-1) ESBLs family has been increasing in frequency around the world (Woerther *et al.*, 2013).

ESBLs originate mainly because of mutations in  $\beta$ -lactamases encoded by the TEM ,SHV and CTX-M-1 genes, the variants TEM and SHV are most widespread ESBLs, but strains expressing CTX-M-1 ESBLs have initiate to emerge in many countries (Ahmed *et al.*,2013).The  $\beta$ -lactamases CTX-M-1 are characterized by selective degradation of cefotaxime (Paterson and Bonomo 2005).This study aimed to

detected CTX-M-1 ESBLs gene in *E.coli* and *K. pneumoniae* isolated from pregnant women.

## **2. Materials and Methods:**

### **Isolation of Bacteria:**

We collected 330 urine samples from pregnant women from september (2016) to february (2017), who suspected to have urinary tract infection (UTI), consulted for AL- Nassyriah city hospital in Iraq. Urine samples transferred to laboratory and culture according to standard methods, after incubation for 24 h at 37°C, the positive culture were identified by conventional techniques and confirmed by using API 20 E system (Cowan and Steel,2014).

### **Detection of Extended Spectrum $\beta$ -lactamase (ESBLs).**

Double disk synergy test (DDST) method was used to detection of ESBLs production. *E.coli* and *K. pneumoniae* was identification by using tested by spreading on Mueller -Hinton agar plates, then (30 $\mu$ g) disk of cefotaxime, aztreonam (30 $\mu$ g), ceftazidime (30 $\mu$ g) and ceftriaxone (30 $\mu$ g) was dispensed around a disk of amoxicillin -clavulanate (20  $\mu$ g amoxicillin / 10  $\mu$ g clavulonic acid) positioned at a distance of (30mm) (centre to centre). Resulting by increasing the diameter of inhibition zone at the side of amoxicillin-clavulonic acid disk (NCCLS,2007).

### **Detection of CTX-M-1 Gene by PCR Technique.**

The DNA was extracted and purified according to the instructions of the manufacturing company (Geneaid / Korea). The DNA was extracted from (57) isolates of *E.coli* and (21) of *K.pneumoniae*. The CTX-M-1 gene was amplified using primers CTX-M-1 F (5'-TTAGGAARTGTGCCGCTGYA-3') and CTX-M-1R (5'-CGATATCGTTGGTGGTRCCAT-3') (Dallenne *et al.*,2010).

Total DNA (2  $\mu$ l) was subjected to PCR in a 50 $\mu$ l reaction mixture containing 1X PCR buffer [10 Mm Tris-HCl (pH 8.3), 50Mm KCl], 1.5 Mm MgCl<sub>2</sub>, 200 Mm of each deoxynucleotide triphosphate, 1.5  $\mu$ l of each of the primers and 2.5 U of Taq polymrase. PCR Program was performed with following: (1) cycles of (10 min.) initial denaturation at

(94°C), followed by (35) cycles of denaturation at (90°C), (40 sec.) annealing at (60°C), (40 sec.) extension at (72°C), (60 sec.) and a final extension step of (7 min.) at (72°C). (Dallenne *et al.*, 2010).

## **3. Results and Discussion:**

A total of 330 urine samples were collected and cultured at different media, only 90 samples given positive growth results. Gram-negative distributed after diagnosed using conventional techniques and conformed by using API 20 E system to 57(63.3%) *E.coli* and 21 (23.3%) *K.pneumoniae* and 12(13.3%) from other Gram negative bacteria as in Table (1). A high frequency of UTI commonly accepted during pregnancy. Among pregnant women the UTI bacterial infection is the predominant type, the physiological and hormonal changes in urinary tract include changes in ureteral dilatation, tone and bladder volume that may promote the infection in pregnant women. The isolation percentage of this study was in agreement with other studies such as Al- Muk and Hansony (2001) in Basrah (Iraq), Almushaita *et al.*, (2013) in Saudi Arabia and Hamdan *et al.* (2011) in Sudan.

*E.coli* is the most implicated microorganisms and the second common bacterium isolated of Gram negative bacteria was *K.pneumoniae* that causing UTI in pregnant women in the present study. This study correspond with results obtained in Tikrit city by Al-Jebouri and Mdish, (2013), also in AL - Najaf city (Abdulla and Oleiwi,2016) and in Kirkuk (Ali *et al.*, 2007).

Results of detection of ESBLs production were showed that out of 90 isolates 26(28.9%) were ESBLs producing isolates, *E.coli* contain 17(29.8%) and *K.pneumoniae* contain 5(23.8%) ESBLs producing isolates (Table 1).

Table (1): Distribution of ESBLs Production and CTX-M-1 gene in *E.coli* and *K.pneumoniae*.

Isolates	Number of isolates	ESBLs enzyme production	%	ESBLs producing isolates and have CTX-M-1 gene	%
<i>Escherichia coli</i>	57	17	29.8	13	76.5
<i>Klebsiella pneumoniae</i>	21	5	23.8	4	80.0
<i>Klebsiella oxytoca</i>	12	4	25.0	2	50.0
Total	90	26	28.9	19	73.1

The multidrug resistant isolates of *E. coli* has been increasing around the world and is associated with the acquisition of mobile element-encoded  $\beta$ -lactamases (Ramos *et al.*,2012). The CTX-M-1  $\beta$ -lactamases production is a very important mechanism to resist for many antibiotic groups in *K.pneumoniae* and *E.coli* ,this family of plasmid-mediated ESBLs of ambler class A has been detected mainly in Asian countries , Europe and America (Castanheira *et al.*,2008).

In this study, were the production detected of ESBLs in (28.9 %) of Gram negative isolated from the pregnant urine figure(1), this result were correspond with the study reported in Tikrit city 27.5% done by Alsamarai *et al.*, (2016) and this study less than the study in Erbil province 61.6 % by Ahmad and Ali, (2014) .

The study revealed that *E.coli* was the most prevalent ESBL producing isolates 65.4%. This result closed to Al-Nammi (2001) in Baghdad (Iraq), who reported that (72.6%) of *E. coli* isolates gave positive ESBLs, while ESBLs producing *K.pneumoniae* represent 19.2% in present study this near from local study showed 10.5 % done by Al-Charrakh , (2005). Such result was less than the study in Erbil province 40 % by Ahmad and Ali,(2014).




figure (1).Clearance the inhibition of producing of ESBLs due to clavunic acid through expansion of the zone of inhibition between AUG and each of CRO ,CAZ ,ATM and CTX.

The results showed individual band of CTX-M-1 gene were characterized in (688 bp) by comparison with the standard molecular DNA ladder (100-2000bp) figure (2),the CTX-M-1 was considered as a common type of ESBLs that detected at Asia, Europe, North and South America among multidrug-resistant in *E.coli* (Feizabadi *et al.*,2010).

The ESBLs gene CTX-M-1, predominates in this results which present four isolated of *K. pneumoniae* out of 5 isolates producing ESBLs and in 13 isolates of

*E.coli* out of 17 isolates producing ESBLs as in Table (1) ,this result corresponding with result of Al-Mayahie,(2013) in Al-kut city.The CTX-M-1gene predominates in Europe, while in other countries, the ESBL genes are more diverse (Livermore *et al.*, 2012) .In Norway and Portugal,the CTX-M-1 is the ESBLs enzyme most frequently found in *E. coli* (Tofteland *et al.*, 2007). Also,this results are in accordance with (Pitout *et al*, 2005) who reported that repeat UTIs were more likely to be caused by CTX-M-1producing strains than by non-CTX-M-producing strains.



Figure(2): Gel Electrophoresis [Ethidium bromide,Agarose (2%)] of PCR products from extracted Gram negative isolates DNA amplified with primers for ESBLs gene CTX-M-1 .lane(L),DNA Ladder size marker (2Kb ladder).Lanes:21,25,26,32,40,43 and 44 isolates of *E.coli*, lane 20,28, 29 and 45 isolates of *K.pneumoniae* that show positive results for CTX-M-1 of ESBLs genes .lane 30 and lane 47 isolates of *E.coli* show negative result for ESBLs genes .Lane (C) control.

## CONCLUSION:

This study was conducted for detection of extended spectrum  $\beta$ -lactamases enzyme and CTX-M-1 gene among *E.coli* and *K. pneumoniae* isolates that causes urinary tract infections for pregnant women at AL-Nassyriah city in Iraq.

## References:

- Abdulla, M. M. and Oleiwi, I. F. (2016).** Prevalence, associated factors, bacterial and antimicrobial susceptibility patterns of urinary tract infection in a sample of Iraqi pregnant women in Al-Najaf. *European Journal of Biology and Medical Science Research.* 4 (6) :7-29.

- Ahmad ,S.S. and Ali, F.A. (2014 ).** Detection of ESBL, AMPCand Metalo betalactmase Mediated Resistance in Gram- negative bacteria Isolated from Women with Genital Tract Infection. *European Scientific Journal* . 10 (9) : 1857 – 7881.
- Ahmed, O. I. ; El-Hady, S. A.; Ahmed, T.M. and Ahmed, I. Z. (2013).**Detection of bla SHV and bla CTX-M genes in ESBLproducing *Klebsiella pneumoniae* isolated from Egyptian patients with suspected nosocomial infections. *The Egyptian Journal of Medical Human Genetics*. (14): 277–283.
- Al-Charrakh, A. ( 2005).**Bcteriological and genetic study on extended-spectrum  $\beta$ -lactamases and bacteriocins of *Klebsiella* isolated from Hilla city. Ph.D. Thesis. College of Science Baghdad University.
- Al-Jebouri, M. and Mdish, S. ,(2013).**Antibiotic Resistance Pattern of Bacteria Isolated from Patients of Urinary Tract Infections in Iraq.*Open Journal of Urology*. (3): 124-131.
- Al-Mayahie ,S.M. (2013) .**Phenotypic and genotypic comparison of ESBL production by vaginal *Escherichia coli* isolates from pregnant and non-pregnant women. *Annals of Clinical Microbiology and Antimicrobials Journal*. (12): 7-13.
- Al-Muk ,J. M. and Hasony, H. J. (2001) .**Isolation of Gardnerella vaginalis from Pregnant Women with Bacterial Vaginosis in Basrah, Iraq. *Bahrain Medical Bulletin Journal*.23 (3): 124-126.
- Al-mushaita ,M. ; Mohammed ,H. ; Al-Harthy, D. and Abdullah, A. (2013).**Prevalence and Predisposing Factors of Urinary Tract Infections among Pregnant Women in Abha General Hospital.*International Journal of Sciences: Basic and Applied Research*. 11(1): 18-29.
- Al-Nammi, A. A. (2001).** Synergistic effect of some  $\beta$ -lactamase inhibitors with  $\beta$ -lactam antibiotics on some gram- negative bacteria and other  $\beta$ -lactamase producers. M.Sc. thesis. Baghdad: University of Al-Mustansiriya.College of Science. Iraq.
- Al-samarai ,A.M. ;Latif, I.M. and Abdul-Aziz, M.M.( 2016).**Antibiotic susceptibility of Extended Spectrum Betalactmase (ESBLs)Producing *Escherichia coli* .*World journal of pharmacy and pharmaceutical sciences*.5 (7) :195-215.
- Ali, C.;Anwer, S.; ssa, S.and Khorsheed, S.( 2007).**Study of Urinary Tract Infection Among Pregnant Women in Kirkuk.*Tikrit Medical Journal*; 13(2):39- 43.
- Castanheira ,M. ; Mendes ,R.E. ; Rhomberg, P.R.and Jones, R.N. ( 2008).** Rapid emergence of blaCTX-M among Enterobacteriaceae in U.S. Medical Centers: molecular evaluation from the MYSTIC Program. *Microbial Drug Resistance Journal*. 14(3): 211-6.
- Cowan,S.T.and Steel,K.J.(2014).** Manual for the Identification of Medical Bacteria. 3rd edition.G. I. Barrow and R. K. A. Feltham, Eds. Cambridge: Cambridge University .
- Dallenne ,C. ; Costa ,A.D ; Decre, D. ; Favier ,C. and Arlet, G. (2010).** Development of a set of multiplex PCR assays for the detection of genes encoding important b-lactamases in *Enterobacteriaceae*. *Journal of Antimicrobial Chemotherapy*.(65): 490–495.
- Feizabadi ,M.M. ; Delfani, S. ; Raji, N. ; Majnooni, A. ; Aligholi, M. ; Shahcheraghi ,F. ; Parvin, M. and Yadegarinia ,D. (2010) .** Distribution of bla(TEM), bla(SHV), bla(CTX-M) genes among clinical isolates of *Klebsiella pneumoniae* at Labbafinejad Hospital, Tehran, Iran.). *Microbial Drug Resistance Journal*. 16(1): 49-53.
- Haider ,G.; Zehra,N.; Afrozem, A. and Haider ,A. (2010).** Risk factors of urinary tract infection inpregnancy. *Journal of the Pakistan Medical Association*. (60): 213-216.
- Hamdan, Z. ; AbdelHalim, M. and Salah, K.( 2011).** Epidemiology of urinary tract infections and antibiotics sensitivity among pregnant women at Khartoum North Hospital.*Annals of Clinical Microbiology and Antimicrobials*. (2):10-12.
- Livermore, D.M . (2012) .**Current epidemiology and growing resistance of Gram-negative pathogens. *Korean Internal Medicine Journal*. (27): 128-142.
- Mirzaee ,M. ;Owlia, P. and Mansouri, S. (2009).** Distribution of CTX-M beta lactamase genes among *Escherichia coli* strains isolated from patients in Iran. *Lab Medicine* (40): 724-727.
- NCCLS (National Committee for Clinical Laboratory Standards). ( 2007).** Performance standards for antimicrobial susceptibility

testing. NCCLS approved standard M100-S17. National Committee for Clinical Laboratory Standards, Wayne, PA.

- Nicolas-Chanoine ,M.H.; Jarlier, V.; Robert, J.and Arlet, G. (2012).** Patient's origin and lifestyle associated with CTX-M-producing *Escherichia coli*: a case control- control study. 1 (7): 30498.
- Okonko ,J.; Ijandipe ,L.and Ihusanya, O. (2009).** Incidence of urinary tract infection among Pregnant women in Ibadan, South-Western Nigeria. African Journal of Biotechnology. (8):6649-6657.
- Paterson ,D.L. and Bonomo, R.A.( 2005).** Extended-spectrum  $\beta$ -lactamases: aclinical update. Clinical Microbiology Reviews Journal.(18): 657–86.
- Pitout, J.D.; Nordmann, P.; Laupland, K.B.and Poirel, L. (2005).**Emergence of *Enterobacteriaceae* producing extended-spectrum  $\beta$ -lactamases (ESBLs) in the community Journal of Antimicrobial Chemotherapy. (56):52-59.
- Ramos, N. L.; Sekikubo, M. and Dzung, D.T. (2012).** Uropathogenic *Escherichia coli* Isolates from Pregnant Women in Different Countries. Journal of Clinical Microbiology. 50(11): 3569-3574.
- Tofteland, S. ; Haldorsen, B.; Dahl, K.H.; Simonsen, G.S. ; Steinbakk, M. and Walsh, T.R. (2007).** Effects of phenotype and genotype on methods for detection of extended spectrum  $\beta$ - lactamase producing clinical isolates of *Escherichia coli* and *Klebsiella pneumoniae* in Norway. Journal of Clinical Microbiology.(45):199-205.
- Trecarichi ,E.M.; Cauda, R. and Tumbarello,M. (2012)** .Detecting risk and predicting patient mortality in patients with extended-spectrum beta-lactamase-producing *Enterobacteriaceae* bloodstream infections. Future Microbiology journal.(7): 1173–89.
- Woerther ,P.L.; Burdet ,C.; Chachaty ,E. and Andreumont, A. (2013).**Trends in human fecal carriage of extended-spectrum beta-lactamases in the community: toward the globalization of CTX-M. Clinical Microbiology Reviews.(26): 744–58.