

## M1 TRANSITIONS IN ER ISOTOPES

A. M. ASHIJ

*Department of Physics-College of Science- University of Anbar,*

*Received: 14/2/2007*

*Accepted 11/4/8/2007:*

**ABSTRACT:** We presented the results of schematic calculations of Magnetic dipole transitions in Erbium isotopes from 158 to 168 by considering the Hamiltonian of the neutron- proton Interacting Boson Model (IBM-2). New idea for counting bosons number at N=64 has been used. A comparison with available M1 experimental data is presented.

*Key words: M1 TRANSITIONS , Er ISOTOPES*

### INTRODUCTION

The even-even Er isotopes lie in the mass region between N=90 to N=100, their low lying states have been successfully described within framework of either the phenomenological or microscopic models[1,2,3,4]. Soloviev et.al[5] calculated the reduced probabilities of M1 and M2 in Er isotopes between excited states within the framework of Quasiparticle-phonon nuclear model. The quadrupole properties of k=2 gamma-vibrations in this well deformed nuclei have been extensively studied [6]. But our knowledge of the magnetic dipole properties remain severely limited. In all collective model the M1 transitions are strictly forbidden. However, it is experimentally observed between inter band states and cross band states [7,8,9]. Bohr and Mottelson [1] shown that gamma-ground M1 transitions

can't be explained via K=2 mixing alone, but they require, in addition, the existence of  $\Delta k=1$  mixing with higher lying K=1 band. Warner [10] shows that the difference in the strength of the pairing for neutron and proton gives rise to different neutron and proton deformations which in term of generate M1 transition. The IBM-1 [11], which deals only with states completely symmetric in the proton-neutron, considered zero terms for M1 transition while IBM-2 gives a complete calculation for M1 matrix elements by considering F-spin ( $F(\max) = (N_{\pi} + N_{\nu})/2$ ) [12-14]. Scholton et.al [15] described the existence of M1 transitions between collective states in deformed nuclei as a nature of states which are not fully

symmetric in the neutron and proton degree of freedom.

The aim of this work is to calculate the M1 matrix elements in deformed Er isotopes, using the IBM-2, and to compare the results with the experimental data.

**THE MODEL**

In the IBM-2 the structure of the collective states in even-even nuclei is calculated by considering a system of interacting neutron (v) and proton (π) boson s(l = 0) and d(l = 2). The boson Hamiltonian can be written as [15]:

$$H = E_d(n_{dv} + n_{d\pi}) + KQ_v^{(2)} \cdot Q_\pi^{(2)} + M_{v\pi} \dots\dots\dots(1)$$

where

$$Q_p = (s_p^+ d_p + d_p^+ s_p)^{(2)} + X_p (d_p^+ d_p)^{(2)}$$

$$p = v, \pi$$

and

$$M_{v\pi} = L_2(s_v^+ d_\pi^+ - d_v^+ s_\pi^+)^{(2)}(s_v d_\pi - d_v s_\pi)^{(2)} - 2 \sum_{k=1,2} L_k (d_v^+ d_\pi^+)^k (d_v d_\pi)^k$$

The majorana term  $M_{\pi v}$  introduced the force which creates the K=1 band in deformed nuclei and push up all states with an amount that depends only on their symmetry with respect to interchange of neutron and proton:

$$T(E2) = e_\pi Q_\pi + e_v Q_v \dots\dots\dots(2)$$

Where  $e_\pi$  and  $e_v$  are proton and neutron effective charges.

The M1 transition operator is :

$$T(M1) = \sqrt{3/\pi} (g_\pi L_\pi + g_v L_v) \dots\dots\dots(3)$$

where  $L_v(L_\pi)$  is the neutron and (proton) angular momentum operator

$$L_p^{(1)} = \sqrt{10} (d^+ d)^{(1)}$$

where  $P = \pi, v, g_\pi$  and  $g_v$  are the effective boson (proton,neutron) geomagnetic –factors.

**THE RESULTS AND DISCUSSION**

In Erisotopes the number of valance nucleons outside the closed shell are large(Number of protons 18 and neutrons 8-18).Which means that we have 9 proton bosons and neutron bosons that start from 4 to 9.These numbers are outside the IBM-2 space from the computer point of viewing .To solve this problem we depend on the sub closed shell at 64 ,which suggested by Priston[2] .This idea reduced the total bosons number to 6 for  $Er^{158}$  to 11 for  $Er^{168}$  . With these bosons number and model parameters listed in table-1.

NPBOS [16] code used to get the best fit for experimental energy level ,then NPBOS code has been used to calculate the transition matrix elements .Electric quadrupole transition probability (E2) have been calculated using the effective charge  $e_\pi = 0.17$  eb and  $e_v = 0.225$  eb which have been estimated using the method described in reference [17].These values are slightly different from the general values estimated by Sala at.al[18]  $e_\pi = e_v = 0.13eb$ , for rare earth nuclei. The results of the calculation of the B(E2) matrix elements are shown in figure(1) and (2). The fitting of E2 matrix elements is essentials for the calculation of the M1 matrix elements, because they share with the  $\delta(E2/M1)$ . So we should make sure that the calculated E2 reduced transition bribability are doing good, and then staring with the M1 fraction in the transition. As one can see from the samples on both

figures the agreement is quite acceptable between calculating and experimental value for  $B(E2; 2_1 \rightarrow 0_1)$  and  $B(E2; 2_2 \rightarrow 2_1)$ .

The M1 transition strength has been calculated by using the operator (3). The boson g-factor was determined using the relation shown in reference [19] which is written as :

$$g = (N_\pi / N)g_\pi + (N_\nu / N)g_\nu = Z / A$$

.....(4)

One experimental  $B(M1)$  also used to generate two relation for  $g_\pi$  and  $g_\nu$  from which we get  $g_\pi = 0.446$  and  $g_\nu = 0.369$ . These values were then generalized for all Er isotopes. They are different from those of the rare - earth nuclei, ( $g_\pi - g_\nu = 0.65$ ), suggested by van Isacker et al. However they also used  $g_\pi = 1$  and  $g_\nu = 0$  to reduce the number of the model parameters in their calculation of M1 properties in deformed nuclei. The results of our calculation are listed in table-2. A good agreement between the theory and the available experimental data is achieved. As can be seen from the table yields to a simple prediction that M1 matrix elements values for gamma to ground and transitions should be equal for the same initial and final spin. Also the size of gamma to ground matrix elements seems to decrease as the mass number increases.

#### CONCLUSIONS

- 1- By fitting  $B(M1)$  from  $2_\gamma$  to  $2_g$  we always get small value for  $g_\pi - g_\nu$  compared with the value basis on the microscopic calculations  $g_\pi - g_\nu = 1$ .
- 2- There are evidences that M1 small mode exists in all spectra.

- 3- In the present work we are paying no attention to the sign of M1 matrix elements which might effect the results.

We can not judge on the agreement between theoretical and experimental data from the above table due to the lack of experimental data. However both experiments and theory predicts small M1 component which is due to symmetry and forbiddances of band crossing gamma transitions.

#### References

- [1] A. Bohr and B.R.Motelson "nuclear structure" vol. 2 (benjamin Reading) (1975)
- [2] M.A. Priston.and.K.Bhaduri "Structure of the Nucleus" (Addison-Wesley) (1975)
- [3] D.J. Rowe "Nuclear collective motion" (Methuen and CO.LTD) (1970)
- [4] K. Kumar "nuclear models and Search for unity in Nuclear Physics" University tsferlaget (1984)
- [5] V.G.Soloviev, A.v. Sushkov and N.Y. Shirikova, J. Phys. G: Nucl.Part.Phys.,20,113, (1994)
- [6] P. Phallon, Acta Physica Polonica B, Vol 36, 1003, (2005)
- [7] W. Gelletly , P. van Isacker, D.D. Warner,G. Colvin and K. Schreckenbach, Phy. Lett. B.3, 191, (1987)
- [8] P. von Brentono, A. Gelbert, H. Harter and P. Sala, J.Phys.G: Nucl.Phys., 11, L85, (1985)
- [9] H.Maser and S. Lindenstruth and von Brentano, Phys.Rev.,C53,2749, (1996)
- [10] D.D. Warner, Phys.Rev.lett. 47,25,1819, (1981)
- [11] F. Iachello and A. Arima, "Interacting Boson in Nuclear


- Physics", (Plenum, New York) (1979)
- [12] F. Iachello and A. Arima, "Interacting Boson Model", (Cambridge Press) (1986)
- [13] W.Pfeifer "An Interacting boson IBM), vdf hochschulverlag-Zurich, (1998)
- [14] P. van Isacker, K. Hyde, J Jolie and A. Sevrin, Ann.Phys.,171,2,253, (1985)
- [15] O. Schelton, K. Hyde and P. van Isacker, Phys.Rev. Lett., 55,18,1866, (1985)
- [16] T.Otsuka and N. Yashida, "Program NPBOS, Japan Atomic Energy and Research Institute Report" No. JAERI-M85-095", (1985)
- [17] A.R.H.Subber, P. Park, W.D.Hamilton and K. Kumar, J.Phys. G: Nucl. Phys. 12,881, (1986)
- [18] P.Sala,a. Gelbert and von Brentono, Z.Phys. A 323, 281, (1986)
- [19] P. van Isacker, P.O. lipas, K. Helimaki and D.D. Warner, Nucl.Phys., A476,303,(1986)

**Table-1- The IBM-2 parameters for Er isotopes**


| A                          | E(MeV) | K(MeV) | XN   | XP    |
|----------------------------|--------|--------|------|-------|
| 158                        | 0.43   | -0.19  | 0.29 | -0.30 |
| 160                        | 0.45   | -0.19  | 0.26 | -0.30 |
| 162                        | 0.47   | -0.18  | 0.25 | -0.30 |
| 164                        | 0.47   | -0.16  | 0.15 | -0.30 |
| 166                        | 0.48   | -0.14  | 0.10 | -0.30 |
| 168                        | 0.51   | -0.13  | 0.13 | -0.30 |
| $L_1=L_3=0.091; L_2=0.052$ |        |        |      |       |

**Table-2-reduced transitions probability B(M1) for gamma to ground and the inter band gamma to gamma transitions in Er Isotopes**

| A   | I <sub>i</sub> | I <sub>f</sub> | B(M1)        |        |
|-----|----------------|----------------|--------------|--------|
|     |                |                | Experimental | Theory |
| 158 | 2 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.066  |
|     | 3 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.0380 |
|     | 3 <sub>γ</sub> | 2 <sub>γ</sub> |              | 0.050  |
|     | 4 <sub>γ</sub> | 3 <sub>γ</sub> |              | 0.046  |
| 160 | 2 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.026  |
|     | 3 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.017  |
|     | 3 <sub>γ</sub> | 2 <sub>γ</sub> |              | 0.044  |
|     | 4 <sub>γ</sub> | 3 <sub>γ</sub> |              | 0.031  |
| 162 | 2 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.041  |
|     | 3 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.029  |
|     | 3 <sub>γ</sub> | 2 <sub>γ</sub> |              | 0.047  |
|     | 4 <sub>γ</sub> | 3 <sub>γ</sub> |              | 0.026  |
| 164 | 2 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.012  |
|     | 3 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.012  |
|     | 3 <sub>γ</sub> | 2 <sub>γ</sub> |              | 0.022  |
|     | 4 <sub>γ</sub> | 3 <sub>γ</sub> |              | 0.012  |
| 166 | 2 <sub>γ</sub> | 2 <sub>g</sub> | 0.000115     | 0.0082 |
|     | 3 <sub>γ</sub> | 2 <sub>g</sub> | 0.00054      | 0.0072 |
|     | 3 <sub>γ</sub> | 2 <sub>γ</sub> |              | 0.0144 |
|     | 4 <sub>γ</sub> | 3 <sub>γ</sub> |              | 0.0011 |
| 168 | 2 <sub>γ</sub> | 2 <sub>g</sub> |              | 0.0044 |
|     | 3 <sub>γ</sub> | 2 <sub>g</sub> | 0.0071       | 0.0038 |
|     | 3 <sub>γ</sub> | 2 <sub>γ</sub> | 0.0036       | 0.0038 |
|     | 4 <sub>γ</sub> | 3 <sub>γ</sub> | 0.00034      | 0.0005 |



**Figure 1** The theoretical and experimental reduced transition probabilities  $B(E2)$  in the unit of  $e^2b^2$  between  $2^+_g$  to  $0^+_g$ . The solid line represents the theoretical values and the dots are the experimental values.



**Figure 2** The theoretical and experimental reduced transition probabilities  $B(E2)$  in the unit of  $e^2b^2$  between  $2^+_{\gamma}$  to  $2^+_g$ . The solid line represents the theoretical values and the dots are the experimental values.

. Er

*Email: [Monean\\_m\\_957@yahoo.com](mailto:Monean_m_957@yahoo.com)*

. 2 -  
. 64

168 158