

MRI ASSESSMENT OF CLINICALLY SUSPECTED MENISCAL TEARS

HASIBA MUAHMED SHUKRI, MBChB, DMRD*
SALEEM KHADIR MUSALAH, MBChB, CABR**

Submitted 12/5/2016; accepted 1/5/2017

ABSTRACT

Background: Magnetic resonance imaging (MRI) is the tool commonly used in the diagnosis of meniscal tears. It has been suggested that, for clinically suspected meniscal tears, the adoption of routine MRI before therapeutic arthroscopy will reduce the number and cost of unnecessary invasive procedures.

The aim of this study was to document MRI yield in knee injuries clinically suspected as possible meniscal tears in Duhok city

Subject and Methods: This cross sectional study was conducted during the period from June to December 2014. A consecutive sampling procedure was used to enroll eighty cases of Knee injuries presenting with clinical features of meniscal tears. All patients underwent MRI examination by 1.5 Tesla machine.

Results: The mean age of the patients was 33.5 years. The number of males was 58 (72.5%) with mean age 29.55 years while females constituted 22 (27.5%) with mean age 44years. The results revealed that 64 patients (80%) were affected by tear in the medial meniscus compared to 16 patients (20%) in the lateral meniscus .

Isolated anterior horn tear was significantly more common in the lateral meniscus (12.5%) than in the medial meniscus (1.25%) while isolated posterior horn tear was more common in the medial meniscus (66.25%) than the lateral meniscus (7.5%)

Conclusions: Meniscal tears were more common in males (~ 3/4th of cases) who presented at a younger age. Medial meniscal tears were four times as common as those affecting the lateral meniscus.

Duhok Med J 2017; 11 (1): 32-37.

Keywords: MRI Assessment, Meniscal Tears

The knee meniscus was known to be as non-important vestigial structure that has no function.¹

In 1887 Sutton described the meniscus as the functionless remains of a leg muscle.² In 1948; Fairbank stated that meniscectomy is not wholly innocuous in his report on the radiographic changes of post meniscectomy which will lead to early degenerative knee changes.³

In 1883 Thomas Annandale was the first one to do meniscal repair.⁴ Now the meniscus is known to play an important function in the complex biomechanics of

the knee⁵. Meniscal tears are the result of either trauma in young athletes or degenerative changes in elderly patients.

The radiological assessment of meniscal tear by MRI examination, significantly participated in avoiding unnecessary diagnostic arthroscopy.⁶ Since the introduction of magnetic resonance imaging in 1980, it has been used as diagnostic tool for musculoskeletal disorders so that arthroscopy now is mostly used for treatment purposes, and it is replaced by the noninvasive MRI for diagnosis.⁷ It is safe for the patient and

* Physician, Department of Radiology, Azadi Teaching Hospital, Duhok, Kurdistan Region, Iraq

**Lecturer, Department of Surgery, College of Medicine, University of Duhok, Kurdistan Region, Iraq

32 *Correspondence author to: Saleem Khadir Musalah, salimokhader@yahoo.com Mobil +9647503521*

does not need exposure to ionizing radiation.⁸

Currently, MRI of the knee for detecting internal derangement has an accuracy rate of more than 90% by most studies.⁹ This study has been designed to assess MRI yield in a sample of patients with knee injuries.

PATIENTS AND METHODS

This cross sectional study was done at the department of surgery and MRI unit/ Azadi teaching hospital/ Duhok/ Kurdistan Region during the period June to December 2014.

All patients with knee complaints referred from the department of surgery suspected of having of meniscal tear, were enrolled consecutively and examined by MRI.

the patient lied on supine position with knee slightly external rotation (5-100), small field of view (14-16cm) and slice thickness of 4-5 cm.

MRI protocol used for the examination was sagittal proton density weighted turbo spine echo sequence with fat suppression (TR=3000ms, TE-30ms) and coronal weighted with fat suppression (TR=3000ms, TE-60ms) axial Gradient and T1W coronal also used.

Intra meniscal signals have three grades on MRI which are first described by Stoller and Colleagues.¹⁰ Grade 1 is rounded or amorphous signal intensity that does not disrupt the articular surface.

Grade 2 is liner signal that does not disrupt the articular surface. Grades 1 and 2 represent myxoid and fluid intra substance degeneration.¹¹ Grade 3 in signal that extended to the articular surface and represents a meniscal tear.¹² Data

management and statistical analysis were performed by using the statistical package for social sciences (SPSS) version 10.

RESULTS

The study sample comprised eighty patients; 58(72.5%) males and 22(27.5%) females. The mean age of the studied groups was 33.52 years; 29.55 years for males and 44years for females. (Table1):

Table 1: Study Sample by Age and Gender

Gender	No. (%)	Mean Age ± SD	*P-value
Gender			
Male	58(72.5)	29.55 ±7.554	<0.001*
Female	22(27.5)	44.00 ±14.102	<0.001*

* Fisher's Exact Test

Sixty four patients (80%) had tear in the medial meniscus compared to 16 patients (20%) with tear in the lateral meniscus Isolated anterior horn tear was significantly more common in the lateral meniscus (12.5%) compared to those affecting the medial meniscus (1.25%) while isolated posterior horn tear was more common in the medial meniscus (66.25%) than in the lateral meniscus (7.5%).

(Table 2):

Table 2: Affected Menisci by Site of Injury

Site of tear	Affected Meniscus				Total	
	Medial Meniscus		Lateral Meniscus		(%)	
	No.	(%)	No.	(%)	No.	(%)
Anterior	1	1.25	10	12.5	11	13.73
Posterior	53	66.25	6	7.5	59	73.75
Combined	10	12.5	0	0	10	12.5
Total	64	80	16	20	80	100

Common stressors

Forty patients (50%) had horizontal tear, 8 (10%) had vertical tear, 10 (12.5%) had bucket handle and 22(27.5%) had complex tear the horizontal, complex and bucket handle tears are more common at the medial meniscus (41%, 24% and 12.5%

MRI ASSESSMENT OF CLINICALLY SUSPECTED MENISCAL TEARS

respectively) while the vertical tear are more common at the lateral meniscus (Table 3):

Table 3: Affected Menisci by Type of Tear

Type of tear	Affected Meniscus				Total	
	Medial Meniscus		Lateral Meniscus		No.	(%)
	No.	(%)	No.	(%)		
Horizontal	33	41	7	12.5	40	40
Vertical	2	2.5	6	7.5	8	10
Bucket handle	10	12.5	0	0	10	12.5
Complex	19	24	3	3.5	22	27.5
Total	64	80	16	20	80	100

DISCUSSION

In the current study, MRI was performed on 80 patients referred from the surgical department because of a unilateral knee injury with clinical suspicion of meniscal tear.

Male predominance (72.5%) was documented, in agreement with another study by Magee and Williams.¹³ which also showed male preponderance (73%).

The mechanism of tear was either traumatic or degenerative. The traumatic type was the most commonly occurring (mean age of 29.39 years) and its rate decreased with increasing age. The non-traumatic cause of injury occurred with mean age of 40.79. These findings are comparable to the findings of a study done in Erbil.¹⁴ Traumatic tears more commonly occurred in patients of lower body mass index (BMI) while the degenerative tears were more common in obese patients; this is in agreement with a study done in Australia.¹⁵

There is a significantly higher proportion of medial meniscus tear (80%) than that of the lateral meniscus (20%).

Isolated anterior horn tear was significantly more common in the lateral meniscus (12.5%) than the medial meniscus (1.25%) while, isolated posterior horn tear was more common in the medial meniscus (66.25%) than the lateral meniscus (7.5%).

The other finding was that horizontal tear was the most common tear (50%) and more commonly occurred at the medial meniscus (41%) mostly in the posterior horn. The vertical longitudinal tear occurred most commonly at the lateral meniscus (10%) while, Bucket handle tear was much more common in the medial meniscus (12.5%) than the lateral meniscus, these results are comparable to a study done by Wright et al., 1995.¹⁶

Complex tears which are combination of horizontal and vertical tears comprised (27.5%), these findings were with agreement with the results of Englund et al.¹⁷

REFERENCES

1. Lento PH, Akuthota V. Meniscal injuries: a critical review. *J Back Musculoskelet Rehabil.* 2000; 15(2): 55-2.
2. Amendola, A. and Bonasia, D. E. The menisci: anatomy, healing response, and biomechanics. In: M.Bonnin, N. A.Amendola, J. Bellemans, S. J. MacDonald and J. Menetrey (eds.). *the knee joint: surgical techniques and strategies.* Paris: Springer, 2012 pp.5-9.
3. Verdonk, P., Depaepe, Y., Desmyter, S., De Muynck, Almqvist, K. F., Verstaete, K., and Verdonk, R. Normal and transplanted lateral knee menisci: evaluation of extrusion using magnetic resonance imaging and ultrasound. *KSSTA,* 2004 Sep., 12(5):411-9.
4. Di Matteo, B., Tarabella, V., Filardo, G., Vigano, A., Tomba, P. And Marcacci, M. . Thomas Annandale: the first meniscus repair. *KSSTA,* 2013 Sep., 21(9):1963-6.

5. Thornton, D. and Rubin, D. Magnetic resonance imaging of the knee menisci. *Seminars in Roentgenology* , 2000 July.,35(3):217-230.
6. Ryzewicz, M., Peterson, B., Siparsky, P.N. and Bartz, R.L.. The diagnosis of meniscus tears: the role of MRI and clinical examination. *Clinical Orthopaedics and Related Research*, 2007 February, 455, pp.123-133.
7. Avcu, S., Altun, E., Akpinar, I., Bulut, M. D., Eresov, K., and Biren, T.(2010). Knee joint examinations by magnetic resonance imaging: the correlation of pathology, age, and sex. *NAJMS* 2010 Apr, 2(4):202-4
8. Jackson, D., Jennings, L., Maywood, R., and Berger, P. Magnetic resonance imaging of the knee. *AJSM*, 1988,16(1):29-8
9. Barnett, M. MR diagnosis of internal derangements of the knee: effected of field strength on efficacy. *AJR*1993 March, 161(1):115-8.
10. Baltazar, R., Ma , C., Shankman, S., Beltran, J. *Knee*. Dogra, in J.R Haaga, V.S. Dogra, M. Forsting, R. C. Gilkeson, H. K. Ha and M Sundaram (eds), *CT and MRI of whole body* .5th ed .Philadelphia PA, Mosby Isevier 2009, PP.2317-2371
11. Renton, P. and Green, R. disease of joints. In *Text book of radiology and imaging* .D.Sutton .7th ed. London .Churchill livingstone/ Elsevier., 2003, 1201-1246.
12. Helms, C.A .magnetic resonance imaging of the knee .In W.E.Brant and C.A.Helms .(eds) .*fundamentals of diagnostic radiology* 3rd ed. Philadelphia. Lippincott, Williams and Wilkins, pp.2007,1193-1205
13. Magee, T. and Williams, D 3.0 T MRI of meniscal tears *AJR* 2006
14. Omar, O.D. the value of MRI examination in specific sequences in evaluation of the meniscal tears of the knee joint 2012 dissertation Arab board of radio diagnosis and medical image, Arabic commission of medical specialization. Iraq.
15. Ding, C., Martel-pelletier, j., pelletier, jp., Abram, F., Raynauld, JP Cieocuttini, F., Jones, G. meniscal tear as an osteoarthritis cohort .a cross-sectional study *journal of rheumatology*, 2007,34(4), 776-784.
16. Wright, D.H., DeSmet, A.A. and Norris, M. Bucket-handle tears of the medial and lateral menisci of the knee., value of MR imaging in detecting displaced fragments. *AJR*, 1995, 165(3). 621-625.

پوخته

ئەگەر ئىشقا تېشىكى نامەيى بو ھەلسەنگاندنا جورىن شكاندنا كركپكا چوكى ئەويىت تىنە دەستنىشانكرن ب تيشكا رەننى

پېشەكى: تيشكا رەننىنا موگناتىسى ئىك ژ چارەسەرىئە سەرەككېيە بو دەست نىشانكرنا نەخوشىت وەكى پېچرىنا غەچرونى بو گەھا چوكى.

رېكېن ھەكولېنى: ويىنەيىت رەننى فاست سېن ئىكو پروتون دېئىستى و T2 ويىتد ئىمېچ ھاتىنە وەرگرتن ب سىستەمى 1.5 تسلا ب شىوازى ساچىتال و كورورنال بو (80) نەخوشا ئەويىت شكاندن ل كركپكا چوكى ھەبووين، و ئەو شكاندن تىنە دابەشكرن بو چوار جورا: ئاسويى، ستويى، ئالوز و باكىت ھاندل.

ئەنجام: ب تيشكا رەننى شكاندنىت كركپكا چوكى تىنە دابەشكرن بو ئاسويى، ستويى، ئالوز و باكىت ھاندل، شكاندنىت ئاسويى ژ ھەمى جورىت دى مشەترن (50%)، پشتى وا شكاندنىت ئالوز (27.5%) شكاندنىت باكىت ھاندل (12.5%) و دوماھىكى شكاندنىت ستويى (10%)، شكاندنىت كركپكا چوكى يىت رەخى ژناقدا پترن ژ رەخى دەرڧە، و رەخى پشتى پترن ژ رەخى بەرڧە.

دەرئەنجام: تيشكا رەننى يا چوكا باشتىن رىكا نەيىنقىزىڧ بو شلوفەكرنا ھەمى جورىت شكاندنىت كركپكا چوكى و بو دىتنا ھەر برىندارىەكا دى دناڧ يان دورماندورىت چوكى.

الخلاصة

تقييم التمزق الغضروفي بواسطة الرنين المغناطيسي

الخلفية والأهداف: التصوير بالرنين المغناطيسي (MRI) هو الوسيلة التي عادة ما تستخدم في تشخيص إصابات الغضروف الهلالي لمفصل الركبة. وقد أشارت الدراسات إلى أن اعتماد التصوير الروتيني بالرنين المغناطيسي قبلًا لتطير العلاجي في حالات الإصابات المحتملة سريريا سوف يقلل من عدد وتكلفة الإجراءات المجتاحة لأعضاء الجسم. كان الغرض من هذه الدراسة هو توثيق نتائج التصوير بالرنين المغناطيسي في إصابات الركبة و التي يشتهب فيها سريريا تضرر الغضروف الهلالي لدى عينة من المرضى في مدينة دهوك.

طرق البحث: أجريت هذه الدراسة المقطعية خلال الفترة من حزيران الى كانون الاول 2014. تم استخدام اسلوب الاعتيان المتعاقب لضم ثمانين حالة من إصابات الركبة ممن لديهم ملامح سريرية لإصابة الغضروف الهلالي. اخضع جميع المرضى لفحص التصوير بالرنين المغناطيسي (1.5 Tesla) .

النتائج: اظهرت النتائج بانمتوسط عمرالمرضى 33.5 سنة. وكان عدد الذكور 58 (72.5%) بمتوسط عمر 29.55 سنة، في حين شكلت الإناث 22 (27.5%) بمتوسط عمر 44 سنة. كما كشفت أن 64 مريضا (80%) تضرر لديهم الغضروف الهلالي الأنسي مقارنة مع 16 مريضا (20%) تضرر لديهم الغضروف الهلالي الوحشي. كذلك اظهرت النتائج بان تمزق القرن الأمامي للغضروف كان أكثر شيوعا في الغضروف الوحشي (12.5%) من الغضروف الأنسي (1.25%) بينما كان تمزق القرن الخلفي أكثر شيوعا في الغضروف الأنسي (66.2%) من الغضروف الجانبي (7.5%).

الاستنتاجات: كانت إصابات الغضروف الهلالي أكثر شيوعا في الذكور (~ 4/3 من الحالات) وحدثت في اعمار اصغر مما هي لدى الاناث كما ان إصابات الغضروف الهلالي الانسي اكثر شيوعا باربعة اضعاف من نضيراتها في الغضروف الهلالي الجانبي.