

Proton Beam Radiation Targeted Nucleotides with Negligible Effect on Interferon

Zainab W Abdul Lateef BSc, MSc

Dept. of Physiology & Medical Physics, College of Medicine, Al-Mustansiriya University.

Abstract

- Background** There is increasing evidence that localized irradiation of the tumor may also modify the tumor microenvironment and generate inflammatory cytokines.
- Objective** This study is aimed to clarify the effect of proton beam radiation on the interferon (IFN- α , IFN- β , IFN- γ) and nucleotide.
- Methods** The Microsoft "The Stopping and Range of Ions in Matter (TRIM-SRIM)" version 1998, and 2003 was used. A model of targeting certain interferon (IFN- α , IFN- β , IFN- γ) as well as the nucleotide pair was created. Each target was subjected to proton radiation of hydrogen [H], helium [He], or carbon [C] at different range of energy seeking for the Bragg's peak.
- Result** The results showed that the cross sections IFN- α , IFN- β , IFN- γ and nucleotide targeted by proton therapy were 0.9776, 0.8317, 0.8297 and 0.7305 [keV/($\mu\text{g}/\text{cm}^2$)] for hydrogen ion, and 2.3354, 2.3414, 2.3377, 2.0842 [keV/($\mu\text{g}/\text{cm}^2$)] for helium ion, and 8.3032, 8.3198, 8.3109, 7.5394 [keV/($\mu\text{g}/\text{cm}^2$)] for carbon ion respectively.
- Conclusions** It concludes that targeting of well precise located tumor with proton beam radiation therapy resulted in nucleotide damage of cancer cell without affecting the immune system in term of interferon surveillance.
- Key words** Proton, Nucleotide, Interferon

Introduction

Interferons (INFs) are glycoprotein belong to cytokines that released in response to the presence of virus, bacteria, parasites or tumor cells. They activate natural killer cells and macrophages and they increase recognition of infective or tumor cell to T lymphocytes. IFN- γ has pleiotropic effects in the tumor microenvironment, including the inhibition of cell proliferation and angiogenesis⁽¹⁾. They reversed the signal defect in T lymphocytes in patients with melanoma and the synthetic INF- α_{2b} is useful as an adjuvant therapy for high risk melanoma⁽¹⁾. Because abnormally low levels of IFN- γ are produced by tumor cells and local T lymphocyte in the glioma, it is a promising adjunct to other immunotherapeutic modalities in the treatment of brain tumors⁽²⁾. Radiation is an important treatment for the

local control of cancer based on its ability to directly kill tumor cells.

However, there is increasing evidence that localized irradiation of the tumor may also modify the tumor microenvironment and generate inflammatory cytokines, which can increase the robustness of the immune response^(4,7). Radiotherapy has been demonstrated to cause inflammation, a potentially beneficial state in which IFN- γ is undoubtedly involved as well as it created a tumor microenvironment conducive for T cell infiltration and tumor cell target recognition⁽⁶⁾. Interferon- α potentiated the cytotoxicity of X-ray radiation⁽⁸⁾. *In vitro* model the production of IFN- γ by cells is suppressed by ultraviolet A1 radiation and thereby the immune system is suppressed⁽³⁾. Recently proton radiation gets

access in management of cancer as a preferable therapeutic modality because fewer harmful adverse reactions, more direct impact on the tumor and increased tumor control. Its effect on the hemopoietic system was generally less pronounced compared to gamma rays and X-rays⁽⁹⁾. Proton radiation was significantly modified the pattern of gene expression in T lymphocytes and highly dependent upon total dose and it may enhance their responsiveness at low dose radiation⁽¹⁰⁾

The stopping power (S) is given by:

$$N.S = - (dE/dx)$$

The quantity of S (keV/μ) is referred to specific energy loss

E: charged particle kinetic energy

-dE: the energy increment lost in infinitesimal material thickness (dx)

N: is number of atom /volume

The specific energy loss is expressed by Bethe-Bloch formula

For heavy charged particle:

$$-\frac{dE}{dx} = \frac{4\pi e^4 z^2}{m_o v^2} NB$$

Where

$$B = Z \left[\ln \frac{2m_o v^2}{I} - \ln \left(1 - \frac{v^2}{c^2} \right) - \frac{v^2}{c^2} \right] - S/2$$

With the following definitions:

v velocity of the charged particle

Z charge of the charged particle

N number density of absorber atoms

Z atomic number of absorber atoms

m electron rest mass

e electron charge

I A parameter, treated as experimentally determined, representing average excitation and ionization potential

B is known as the stopping number (atomic number scaled for stopping)

S is the density correction

Bethe-Bloch formula for electrons:

$$-dE/dx = (2\pi e^4 / m_o v^2) NB$$

$$B = Z \left[\ln \frac{m_o v^2 T}{2I^2(1-\beta^2)} - (\ln 2)(2\sqrt{1-\beta^2} - 1 + \beta^2) + 1 - \beta^2 + \frac{1}{8} (1 - \sqrt{1-\beta^2})^2 \right]$$

Where $\beta = \frac{v}{c}$, T is a constant factor

The total stopping power for electron can be given as a combination of collisional (elastic collision with atomic electrons) and radiative (inelastic collision with nucleus) types of interaction:

$$[dE/dx]_{total} = [dE/dx]_{collision} + [dE/dx]_{radiative}$$

For heavy particles, orbital electron interactions are only considered since the probability of nuclear interaction resulting in energy loss is much smaller.

$$-\left(\frac{dE}{dx}\right)_r = \frac{NTZ(Z+1)e^4}{137m_o^2c^4} \left(4 \ln \frac{2T}{m_o c^2} - \frac{4}{3} \right)$$

The percent of the energy loss goes to emitted rays is expressed by:

$$\left(\frac{dE}{dx}\right)_r / \left(\frac{dE}{dx}\right)_{\text{total}} = EZ/1000$$

Where E is in MeV, where Z is the atomic number of the absorber.

The range of a charged particle can be derived from stopping power formula:

$$R = \int_E^0 dx(\text{cm}) = \int_E^0 \frac{dE}{\left(\frac{dE}{dx}\right)} = -\int_0^E \frac{1}{\left(\frac{dE}{dx}\right)} dE = \int_0^E \frac{dE}{S}$$

This study is aimed to explore the effect of energy loss from the immune system using the hydrogen, helium, carbon, and oxygen sources and interferon as the target in Trim-

Shin model of stopping power (KeV/(μg/cm²)) is referred to specific energy loss per cross

section of targeting molecule. Microsoft Excel 2003 was used for calculations and figures

of Physiology/Medical Physics, College of Medicine, Al-Mustansiriya University in

Baghdad, Iraq. The Microsoft "The Stopping Power of Matter" (SRIM) was used to

calculate the Bragg's peak (-dE/dx) of the targeting

particle of proton, helium, carbon, and oxygen

for different targets. The effect of nucleotide pairing was studied from each target

was subjected to proton and helium of hydrogen

targeting molecule but different Bragg's peak

range of energy seeking for the Bragg's peak.

The characteristics of the targets

are shown in Table 1. The constituents of the targets

are shown in Table 1. The constituents of the targets

well as the targeting cross section (Table 2, Figure 1). The Bragg's peak of proton targeting nucleotide is far away than those of interferons with lesser effect on the cross section of nucleotide (Table 2, Figure 1). The results obtained with proton of helium or carbon sources are similar in pattern but not in magnitude to that obtained with hydrogen source in targeting the interferons or nucleotide (Table 2, Figure 1). The cross section of INF-γ targeted by proton of whatever sources (hydrogen, helium or carbon) is less affected than INF-α and INF-β and its targeted depth is more INF-α and INF-β by 100-400 Angstrom. The cross section of nucleotide targeted by proton is less than those observed with interferon despite of higher Bragg's peak and longer projected distance for different sources of proton (Table 2, Figure 1). The spread out cross section of INF-γ targeted by protons in terms of longitudinal and lateral straggling is higher than corresponding INF-α and INF-β (Table 3). Moreover, the spread out effect of proton targeting nucleotide is higher than interferons by 1.3 for all ion sources (Table 3).

Table 1. The constituents of the targets

	IFN-α	IFN-β	IFN-γ	Nucleotide
Density (g/cm ³)	0.98010	0.98276	0.97573	1.1165
Atomic percent (Mass percent) in infinitesimal material				
C	31.82 (53.69)	32.18 (54.45)	31.55 (53.32)	29.62 (38.41)
H	50.02 (7.08)	49.92 (7.09)	50.01 (7.09)	35.83 (3.90)
N	8.39 (16.52)	8.72 (17.21)	8.87 (17.48)	18.51 (28.0)
O	9.43 (21.21)	8.93 (20.13)	9.32 (20.98)	14.81 (25.58)
S	0.33 (1.5)	0.25 (1.12)	0.25 (1.12)	-
P	-	-	-	1.25 (4.12)

C (carbon), H (hydrogen), N (nitrogen), O (oxygen), S (sulfur), P (phosphate)

Table 2. Effect of proton originated from different ions sources on the interferon and nucleotide

Ion	Target	Energy (KeV)	-dE/dx (KeV/μ)	Depth (μm)	Cross section keV/(μg/cm ²)
H	IFN-α	90	81.27	1.39	0.9776
	IFN-β	90	81.74	1.39	0.8317
	IFN-γ	90	80.96	1.40	0.8297
	Nucleotide	100	81.56	1.52	0.7305
He	IFN-α	550	228.9	3.47	2.3354
	IFN-β	550	230.1	3.45	2.3414
	IFN-γ	550	228.1	3.49	2.3377
	Nucleotide	600	232.7	3.70	2.0842
C	IFN-α	2400	813.8	4.33	8.3032
	IFN-β	2400	817.6	4.30	8.3198
	IFN-γ	2400	810.9	4.34	8.3109
	Nucleotide	2800	842.4	4.77	7.5394

Table 3. The lateral and radial struggle of proton of each target at Bragg's peak

Ion	Target	Longitudinal (μm)	Lateral (μm)	Cross section of damage beyond the target (μm ²)
H	IFN-α	0.1106	0.1514	0.01674
	IFN-β	0.1094	0.1498	0.01638
	IFN-γ	0.1110	0.1520	0.01687
	Nucleotide	0.1271	0.1732	0.02201
He	IFN-α	0.2051	0.2689	0.05515
	IFN-β	0.2029	0.2661	0.05399
	IFN-γ	0.2058	0.2698	0.05552
	Nucleotide	0.2304	0.3020	0.06958
C	IFN-α	0.1879	0.2397	0.04503
	IFN-β	0.1862	0.2373	0.04418
	IFN-γ	0.1885	0.2404	0.04531
	Nucleotide	0.2114	0.2729	0.05769




Figure 1. Bragg's peak deposited in different molecules; INF- α , INF- β , INF- γ and DNA targeted by hydrogen [A], helium [B] or carbon [C].

Discussion

The results showed that the Bragg's peak of proton (the maximum energy loss) of nucleotide is differed from that of interferon which means that proton beam radiation targeting the nucleotide will not affect the interferon and thereby not interferes with immune system. Moreover, the spread out effect of proton against the nucleotide at the Bragg's peak was higher by 1.3 fold of interferon at their Bragg's peak which indicated that proton showed selective effect against nucleotide.

Khvostunov et al (2010) found that whole cell nucleus as a function of proton energy shows a distinct peak at 550 keV using biophysical modeling of radiation effects induced by exposure of V79 cells which is approximated to

that obtained with helium in this study ⁽¹¹⁾. In vivo, proton beam was found to be more cytotoxic to A549 lung adenocarcinoma cell than gamma radiation ⁽¹²⁾. Previous studies showed that proton radiation exerts minimal effect on immune system as showed in this study.

The cell death in the splenic white pulp of irradiated whole body ICR mice with proton was lower compared with gamma radiation in spite of an increase damaged DNA ⁽¹³⁾. Moreover, there is an evidence of using interferon, which is not targeted by proton in this study, in cutaneous melanoma patients to prevent metastasis and recommended to use interferon following proton radiation in patients with high risk of metastasis ⁽¹⁴⁾.

This study adds more information that endogenous interferon was not affected by proton when the later targeted the nucleotide which means that the immune system is free from the effect of proton radiation as it happens with conventional X-ray radiation⁽¹⁵⁾. It concludes that targeting of well precise located tumor with proton beam radiation therapy resulted in nucleotide damage of cancer cell without affecting the immune system in term of interferon surveillance.

References

1. Tahrini AA, and Kirkwood JM. Clinical and immunologic basis of interferon therapy in melanoma. *Ann N Y Acad Sci*, 2009 Dec; 1182: 47-57.
2. Kane A, and Yang I. Interferon-gamma in brain tumor immunotherapy. *Neurosurg Clin N Am*, 2010; 21: 77-86.
3. Smit N, Musson R, Romijn F, van Rossum H, and van Pelt J. Effects of ultraviolet A-1 radiation on calcineurin activity and cytokine production in (skin) cell cultures. *Photochem Photobiol*, 2009; 86: 360-366.
4. McBride WH, Chiang CS, Olson JL, Wang CC, Hong JH, Pajonk F et al. A sense of danger from radiation. *Radiat Res*, 2004; 162: 1-19.
5. Boehm U, Klamp T, Groot M, Howard JC. Cellular responses to interferon- γ . *Annu Rev Immunol*, 1997; 15: 749-795.
6. Lugade AA, Sorensen EW, Gerber SA, Moran JP, Frelinger JG, and Lord EM. Radiation-induced IFN-gamma production within the tumor microenvironment influences antitumor immunity. *Immunol*, 2008; 180: 3132-3139.
7. Zou, W. Immunosuppressive networks in the tumour environment and their therapeutic relevance. *Nat Rev Cancer*, 2005; 5: 263-274.
8. Horiguchi-Yamada J, Iwase S, Kawano T, and Yamada H. Pretreatment with interferon-alpha radiosensitizes Daudi cells modulating gene expression and biomarkers. *Anticancer Res*, 2005; 25: 2631-2638.
9. Gridley DS, Rizvi A, Luo-Owen X, Makinde AY, Coutrakon GB, Koss P et al. Variable hematopoietic responses to acute photons, protons and simulated solar particle event protons. *In Vivo*, 2008; 22: 159-169.
10. Gridley DS, Pecaut MJ, Rizvi A, Coutrakon GB, Luo-Owen X, Makinde AY et al. Low-dose, low-dose-rate proton radiation modulates CD4(+) T cell gene expression. *Int J Radiat Biol*, 2009; 85: 250-261.
11. Khvostunov IK, Nikjoo H, Uehara S, and Hoshi M. The consideration of biological effectiveness of low energy protons using biophysical modeling of the effects induced by exposure of V79 cells. *Radiat Biol Radioecol*, 2010; 50: 81-89.
12. Ghosh S, Bhat NN, Santra S, Thomas RG, Gupta SK, Choudhury RK et al. Low energy proton beam induces efficient cell killing in A549 lung adenocarcinoma cells. *Cancer Invest*, Jul; 28(6): 615-22.
13. Finnberg N, Wambi C, Ware JH, Kennedy AR, and El-Deiry WS. Gamma-radiation (GR) triggers a unique gene expression profile associated: with cell death compared to proton radiation (PR) in mice in vivo. *Cancer Biol Ther*, 2008; 7: 2023-2033.
14. Munzenrider JE. Uveal melanomas: conservation treatment. *Hematol Oncol Clin North Am*, 2001; 15: 389-402.
15. Meo SA, Al Drees AM, Zadi SZ, Al Damgh S, and Al-Tuwaijri AS. Hazards of X-ray radiation on the quantitative and phagocytic functions of polymorphonuclear neutrophils in X-ray technicians. *J Occup Health*, 2006; 48: 88-92.

Correspondence to: Zainab W. Abdul Lateef,
E-mail: zainabwahbee@yahoo.com

Received: 26th Apr. 2010, Accepted: 6th Dec. 2010