

**Synthesis and Spectral Analysis of VO(II),Cr (III) ,
Zn(II),Cd(II),Hg(II)and UO₂(II) Complexes with Mixed Ligands
of Bipyridyl and Novel Schiff Base**

Abbas Ali Salih Al-hamdani*,Amer JabbarJarad**,Sargol Abd-AlGhani AL-Atrakchi***
Department of chemistry, College of Science for women ,University of Baghdad
Department of chemistry, Ibn-Al-Haithem College ,University of Baghdad
The Central Laibrary, University of Baghdad

Abstract

Salicylaldehyde was react with 4-amino-2,3-dimethyl-1-phenyl-3-Pyrazoline-5-on to produce the novel Schiff base ligand 2,3-dimethyl-1-phenyl-4-salicylidene-3-pyrazoline-5-on (HL). A new complexes of VO(II), Cr(III), Zn(II), Cd(II), Hg(II) and UO₂(II) with mixed ligands of bipyridyl and new shiff base (2,3-dimethyl-1-phenyl-4-salicylidene-3-pyrazoline-5-on) (HL) were prepared . All prepared compounds were identified by atomic absorption, FT.IR , UV-Visable spectra and molar conductivity. From the above data, the proposed molecular structure for VO(II) complex is squire pyramidal while (Zn(II), Cd(II), Hg(II)) and (UO₂(II),Cr(III)) complexes are forming tetrahedral and octahedral geometry respectively.

Introduction

Synthesis of transition–metal complexes with schiff–base have been a subject of considerable importance ⁽¹⁾ .These complexes are good coordination agents for schiff–base and also good bactericide ⁽²⁾ ,antitumor ⁽³⁾ , antifungal ⁽⁴⁾ and anticancer ⁽⁵⁾ .A search through literature ⁽⁶⁻⁸⁾ reveals that no work has been done on the transition metal complexes of the schiff 2,3-dimethyl-1-phenyl-4-salicylidene-3-pyrazoline-5-on and bipyridyl .

The present paper reports the synthesis and characterization of new VO(II),Cr(III),Zn(II),Cd(II),Hg(II) and UO₂(II) complexes with mixed ligand of bipyridyl and new schiff base.

Experimental


a /Materials:- All chemical used were of reagent grade and used without further purification VOSO₄.H₂O, ZnCl₂, CrCl₃.6H₂O, CdCl₂.H₂O, UO₂(CH₃COO)₂.2H₂O, HgCl₂(Fluka) Salicylaldehyde , KOH, 4-amino -

2,3-dimethyl-1-phenyl-3-Pyrazoline-5-on, DMF, Methanol , Ethanol and acetic acid (B.D.H).

b /Instrumentation:-FT.IR- spectra as CsI discs, in the range (4000 – 200) cm^{-1} were obtained using a Shimadzu , FT.IR-8400 S Fourier Transform Infrared Spectrophotometer .Electronic spectra were recorded in DMF 10^{-3} M using (Shimadzu UV-160A) Ultra Violet-Visible Spectrophotometer. The conductivities of 10^{-3} M solution of the metal complexes in DMF were measured using (Philips Pw – Digital Meter) conductmeter at room temperature. Atomic absorption measurement were obtained using (Shimadzu A.A-160) Atomic Absorption/Flame Emission Spectrophotometer. Magnetic properties were recorded using (Balance Magnetic Susceptibility Model MSB-MKi). In addition melting point were measured using (Gallen kamp Melting Point).

Synthesis of 2,3-dimethyl-1-phenyl-4-salicylidene-3-pyrazoline-5-on (HL) ⁽⁶⁾

(0.26 ml , 2.47 mmole) of salicylaldehyde was added to ethanolic solution of (0.5 g , 2.47 mmole) of 4-amino-2,3-dimethyl-1-phenyl-3-pyrazoline-5-on, The solution mixture was stirred and refluxed for 2 hours, yellow crystalline precipitate observed. The resulting precipitate was filtered off, recrystallized from menthol and dried at 50 °C.The preparation method of the ligand (HL) is represented in scheme-1.



(Schem-1) preparation method of the ligand (HL)

Synthesis of Complexes (general method)

All complexes were prepared by dissolving (0.306 g, 1mmole) of 2,3-dimethyl-1-phenyl-4-salicylidene-3-pyrazoline-5-on in ethanolic (KOH) solution . The solution was added gradually with stirring to the ethanolic metal salt contains 0.181g, 0.266g, 0.228g,0.172g, 0.271g and 0.424g (1mmole) of $\text{VOSO}_4 \cdot \text{H}_2\text{O}$, $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$, ZnCl_2 , $\text{CdCl}_2 \cdot \text{H}_2\text{O}$, HgCl_2 and $\text{UO}_2 (\text{CH}_3\text{COO})_2 \cdot 2\text{H}_2\text{O}$ respectively, an ethanolic solution of (0.156g ,1mmole) bipyridyl was added to the mixture in each case. the solution mixture was refluxed for 4 hours. crystalline precipitate was filtered off and recrystallized from methanol.

Result and Discussion

The isolated complexes were crystalline solids soluble in some common solvents such as ethanol and methanol. The conductivity measurements in DMF (10^{-3}M) at 298°K indicated the electrolyte type ratio (1:1)⁽⁹⁾.(Table-1) includes the physical properties of all prepared compounds.

FT.IR- Spectra

The FT.IR-spectral data of the ligands and all complexes are listed in (Table-2), the FT.IR-spectrum of the ligand(HL), (Fig-1) abroad vibration band at 3290cm^{-1} was assigned to phenolic OH group. The disappearance of this band in the spectra of all the complexes indicated the deprotonation of phenol proton prior to and indicated the coordination the phenolic oxygen with metal ion⁽¹⁰⁾. The bands at 3060cm^{-1} and 2935cm^{-1} refers to $\nu(\text{C-H})$ aromatic and aliphatic in the ligand^(11,12). The band at 1656cm^{-1} is characterizatic of the carbonyl group present in the free schiff base ligand ,this band was shifted to lower frequency 1627cm^{-1} in IR spectrum of Cr(III) complex only, binary and mixed ligand complex, which indicate the involvement of oxygen of carbonyl group in coordination⁽¹³⁾.

The band assigned to the azomethine group in the free schiff base ligand was a bserved at 1600cm^{-1} and shifted to lower frequency in IR spectra of all complexes. This indicates the participation of the nitrogen atom of the azomethine group in coordination^(14,15). The stretching frequency due to N-N in the free ligand were observed at 1038cm^{-1} ⁽¹⁶⁾.

The IR spectra of vanidyl complex (Fig-2) record the presence of new bands which are representing the stretching bidentate SO_4^{-2} group and $\text{V}=\text{O}$, these bands are observed at 1245 cm^{-1} , 509 cm^{-1} and 950 cm^{-1} respectively^(17,18). Also new band observed in Uranyl complex at 907 cm^{-1} due to $(\text{O}=\text{U}=\text{O})$ ⁽¹⁹⁾. Metal–nitrogen and metal–oxygen bands further confirmed by the presence of the bands around $(532\text{--}492)\text{ cm}^{-1}$ and $(487\text{--}340)\text{ cm}^{-1}$ ^(20,21). According to the previous IR spectral data, we can conclude that the schiff base behaves as bidentate ligand through the oxygen of phenolic OH group and nitrogen of azomethine group with VO(II), Zn(II), Cd(II) and Hg(II), but schiff base behaves as tridentate ligand through the oxygen of phenolic OH group and nitrogen of azomethine group and oxygen of carbonyl group with Cr(III) complex only.

The band at 1619 cm^{-1} in IR spectrum of bipyridyl (Fig-3) was shifted to lower frequency in IR spectra of all complexes and appeared at range $(1568\text{--}1572)\text{ cm}^{-1}$, these shifting indicates the coordination between two nitrogen of bipyridyl and metal ions⁽²²⁾

Electronic Spectra

The UV-Vis spectra data for of the free ligands (HL) and bipyridyl and all metal complexes are listed in (Table-3). The UV-Vis spectrum of the ligand (HL) (Fig-4) shows three peaks at 256 nm , 340 nm and 362 nm assigned to $(\pi - \pi^*)$ and $(n - \pi^*)$ electronic transitions ^(23,24). The electronic spectrum of bipyridyl (Fig-5) shows two peaks at 240 nm and 285 nm attributed to $(\pi \rightarrow \pi^*)$ and $(n \rightarrow \pi^*)$ electronic transition respectively⁽²⁵⁾.

The Electronic Spectra of Complexes

1- $\text{K}[(\text{VO})_2(\text{L})(\text{bipy})(\text{SO}_4)_2]$:-The yellowish green complex (Fig-6) showed two absorption peaks at 354 nm and at 380 nm due to charge transfer (C.T). The peak at 610 nm was attributed to $(d - d)$ electronic transition type $({}^2\text{B}_{2g} \rightarrow {}^2\text{B}_{1g})$ which is a good evidence for square pyramidal geometry about V(IV) ⁽²⁶⁾.

2- $[\text{Cr}(\text{L})(\text{bipy})\text{Cl}]\text{Cl}$:-The brown complex gave two absorptions at 256 nm and at 315 nm assigned to ligand field. The peak at 370 nm due to charge transfer (C.T). other two peaks at 425 nm and at 510 nm were found to be caused by $(d-d)$ electronic transition type ${}^4\text{A}_{2g}(\text{F}) \rightarrow {}^4\text{T}_{1g}(\text{F})$ and ${}^4\text{A}_{2g}(\text{F}) \rightarrow {}^4\text{T}_{2g}(\text{F})$ suggesting octahedral geometry about Cr(III) ⁽²⁷⁾


3- $[\text{Zn}(\text{L})\text{bipy}]\text{Cl}$, $[\text{Cd}(\text{L})\text{bipy}]\text{Cl}$, $[\text{Hg}(\text{L})\text{bipy}]\text{Cl}$:-Yellowish green complexes:- The peaks appeared at 263 nm and 348 nm in the spectrum of Zn^{+2} complex, 262 nm and 345 nm in the spectrum of Cd^{+2} complex and finally the peaks at 267 nm and 348 nm in the spectrum of Hg^{+2} complex could be attributed to the ligand field. The peaks at 355 nm , 370 nm and 369 nm due to charge transfer in Zn^{+2} , Cd^{+2} and Hg^{+2} complexes respectively ⁽²⁸⁾. The absence of absorption peaks in the range

(371-1000) nm in the electronic spectra of Zn^{+2} , Cd^{+2} and Hg^{+2} complexes indicated no (d-d) electronic transition happened (d^{10} -system) in the visible region ,that is a good result for Zn^{+2} , Cd^{+2} and Hg^{+2} tetrahedral complexes⁽²⁹⁾.


4-[$UO_2(L)bipy$](CH_3COO):- The reddish brown complex showed two absorption peaks at 265nm and 350nm attributed to ligand field. The third band at 376nm was caused by charge transfer⁽¹⁵⁾.

According to the results obtained and spectral analysis, the chemical structures of the complexes may be suggested as in the following :-

a- Square pyramidal for $K [(VO)_2 (L) bipy (SO_4)_2]$.




b-Tetrahedral for the complexes $[Zn(L) bipy]Cl$, $[Cd(L)bipy]Cl$ and $[Hg(L) bipy]Cl$.



M= Zn^{+2} , Cd^{+2} , Hg^{+2}

C-Octahedral for $[Cr(L)(bipy)Cl]Cl$ and $[UO_2(L)bipy](CH_3COO)$.



References

- 1- Das, G ;Shukla, R; Mandal Singh, R and Bharadwaj, P. K., Inorg. Chem., 36,323 – 329 (1997).
- 2- Sokolik, j; Blahovg , M; Cukanova, G; Kohutora, M; Misikova , E and Mlynarcik , D., Ceska . Slov . Farm., 47,180 – 188 (1998).
- 3- Li , Z.L ; Chen , J.H ; Zhang , K.C Li M and Land Yu, R .Cl., Sci . China B., 36 , 214 -224(1993).
- 4- Buer, A .W; Kirby, W.M.M; Sheries, J and Cand Turck, M., J. Am.Clin. Pathol., 44,493-496 (1966).
- 5- Keypour, H ; Salehzadeh, S and Parish,R.V; J.Molecules.,7,140-144 (2002).
- 6- AL- hamdani,A.A.S., J.Um-Salama.Sci.,2,2(2005) .
- 7-Azawy, S.A.S and AL- hamdani,A.A.S; J.Um-Salama.Sci.,1, 2(2007).
- 8-AL- hamdani,A.A.S; J.Um-Salama.Sci.,2, 2 (2008).
- 9- Geary,w.j., "The Use of Conductivity Measurements Inorganic Solvents for The Characterization of Coordination Compounds"., Coord.Chem. Rev.,7 (1971).
- 10- Yildiz,E and Boztepe,H., J.Chem.Turk.,26, 897-903(2002).
- 11-Korn,M; Ferreira,A; Texeira,L and Costa,A; J.Braz.Chem .Soc .,10, 46(1999).
- 12- Vicente,J;GIL-Rubio,J and Barquero,N., J.Poly.Sci.,43,1-6(2005).

- 13- Colthup,N.B; Daly,L.H and Wiberley,S.E; " Introduction to Infrared and Raman spectroscopy",2nd.Ed.Academic Press.Inc.J.R.Dyer(1975).
- 14- Stuart,B;George,B and Mdnrtyre,P; "Modren Infrared Spetroscopy"John Wiley and Sons,New York (1998).
- 15- SI,Y; Hu,Q; Huang,Z; Yang,G and Yin,J., Turk.J.Chem.,29, 135-140(2005) .
- 16- Maradiya,H.R and Patel,V.S., J.Serb.Chem.Soc.,66,87-93(2001).
- 17-Schumann,E; Altman,J; Karaghiosoff,K and Beek,W., Inorg.Chem.,34,154-165(2002).
- 18- Raman,N; Raja,Y.P and Samy,A.K., Proc.Indian.Acad.Sci.,113,3(2001).
- 19-Martijn,M.G ;Antonisse,B.H.M and Davaid,N.R., J.Org.Chem.,63,26(1988)
- 20- Pabha Karam,C.P and Patel,C.C.J., Inorg.Nucl.Chem.,31,3316(1969) .
- 21- AL-Janabi,M.Y., "The Physical Methods in Inorganic Chemistry"(1983).
- 22- Khan,Z; Ali,S.I; Gupta,D and Khan,A.A; Indian.J.Chem;35A,320-323,Apriil (1996).
- 23- Bair,M.L and Larsen,E., J.Am.Chem.,10,84-90(1971).
- 24- Sonmez,M and Sekerci,M., J.Chem.,76,121-126(2002).
- 25- Suzuki,T; Shiotsuki,M; Wada,K; Kondo,T and Mitsudo,T., Organo. Metal. lics.,18,18(1999).
- 26- Thompson,L; Tandon,S.S and Mannel,M.E., Inorg.Chem;34,425(1995).
- 27- Daniel,E; Seema,C; Melryn,R.C and Kim,M,K., Inorg.Chem.,40,7(2001).
- 28- Guang,P.X.Y and Guang,P.F., J.Med. Gen. Mede.,22,6(2002).
- 29-Sandhu,S.S; Jaswal,J.S and Sandhu,G.K; Indian.J.Chem;35,218-222(1996).

Diala , Jour , Volume , 36 , 2009

bipy	—	3050	—	—	1619	—	—	—	—	—
HL	3290	3060	2935	1656	1600	1038	1303	—	—	—
K[(VO)₂(L)(bipy)(SO₄)₂]	-	3062	2931	1654	1572 1586	1040	1303	492	455	—
[Cr(L)(bipy)Cl]Cl	-	3050	2939	1627	1568 1588	1039	1303	510	420	391
[Zn(L)bipy]Cl	-	3061	2919	1655	1570 1579	1038	1303	523	412	—
[Cd(L)bipy]Cl	-	3043	2945	1654	1568 1591	1039	1303	541	340	—
[Hg(L)bipy]Cl	-	3076	2934	1655	1569 1589	1039	1303	516	487	—
[UO₂(L)bipy](CH₃COO)	-	3062	2939	1656	1569 1582	1040	1305	532	485	—

Table (3): The Electronic spectra for the free ligands and their complexes in DMF(10^{-3} M)

Compounds	λ_{nm}	ABS	νcm^{-1}	$\epsilon_{max} L.mole^{-1}.cm^{-1}$
HL	256	1.365	39062.5	2632
	340	2.265	29411.7	4458
	362	1.354	27624.3	2354
bipy	240	2.432	41666.6	2432
	285	2.926	35087.7	2926
K[(VO) ₂ (L)(bipy)(SO ₄) ₂]	260	0.975	38461.5	975
	354	1.157	26315.7	1157
	380	2.158	28248.5	2158
	610	0.043	16393.4	43
[Cr(L)(bipy)Cl]Cl	265	1.37	37735.8	1370
	351	0.925	27027	925
	370	1.88	28490	1880
	425	0.560	23529.7	560
	510	0.068	19607.8	68
[Zn(L)bipy]Cl	263	1.08	38022.8	1080
	348	0.85	28735.6	850
	355	2.2	28169	2200
[Cd(L)bipy]Cl	262	1.093	38167.9	1093
	345	0.65	28985.5	650
	370	2.15	27027	2200
[Hg(L) bipy]Cl	267	1.35	37453.1	1350
	348	0.85	28735.6	850
	369	2.45	27100.2	2450
[UO ₂ (L)bipy](CH ₃ COO)	265	1.382	37735.8	1382
	350	1.303	28571.4	1303
	376	2.401	265095.5	2401




Fig. (1): FTIR spectrum of 2,3-Dimethyl-1-phenyl-4-salicylidene-3-pyrazoline-5-on (HL)




Fig. (2): FTIR spectrum of the K[(VO)₂(L)(bipy)(SO₄)₂] complex




Fig. (3): FTIR spectrum of bipyridyl




Fig. (4): UV- Vis spectrum of 2,3-Dimethyl-1-phenyl-4-salicylidene-3-pyrazoline-5-on (HL)




Fig. (5): UV- Vis spectrum of bipyridyl




Fig. (6): UV-Vis spectrum of the $\text{K}[(\text{VO})_2(\text{L})(\text{bipy})(\text{SO}_4)_2]$ complex

تحضير وتحليل طيفي لمعقدات VO(II) ، Cr(III) ، Zn(II) ، Cd(II) ، Hg(II) مع UO₂(II) ليكائنات مختلطة للبايريدائل وقاعدة شف جديدة

عباس علي صالح الحمداني* ، عامر جبار جراد** ، سر كول عبد الغني الاطرقجي***
*قسم الكيمياء كلية العلوم للبنات جامعة بغداد ، ** قسم الكيمياء كلية التربية - ابن الهيثم جامعة بغداد
*** المكتبة المركزية جامعة بغداد

الخلاصة

حضرت معقدات جديدة من تفاعل الفنادايل (II) ، الكروم (III) ، الخارصين (II) ، الكادميوم (II) ، الزئبق (II) واليورانييل (II) الجديدة مع الليكائنات المختلطة للبايريدائل وقاعدة شف الجديدة 2،3-ثنائي مثيل-1-فنيل-4-سالسيدين-3-بايروزولين-5-اون. تم عزل المعقدات وتشخيصها باستخدام طيف الأشعة تحت الحمراء ، طيف الأشعة فوق البنفسجية - المرئية و تقنية الامتصاص الذري اللهبي للعناصر فضلا عن التوصيلية الكهربائية . من النتائج المحصول عليها فان الشكل الفراغي المقترح هو هرم مربعي لمعقد الفنادايل (II) بينما اتخذت معقدات (الخارصين (II) ، الكادميوم (II) ، الزئبق (II)) و (الكروم (III) ، اليورانييل (II)) هيئة رباعي السطوح وثمان السطوح على التوالي .