

## Spectrophotometric Determination of Tetracycline by Coupling with Diazotised 4-Aminoantipyrine in Presence of Cetylpyridinium Chloride

Nabeel S. Othman

Rasha J. Al-Ashow

*Department of Chemistry*

*College of Science*

*University of Mosul*

(Received 26/ 6 / 2011 ; Accepted 31 / 10 / 2011 )

### ABSTRACT

A spectrophotometric method for the determination of microgram amounts of tetracycline -HCl(TCH) has been proposed. The method is based on coupling of tetracycline-HCl with diazotized 4-aminoantipyrine in presence of cetylpyridinium chloride in alkaline medium. The molar absorptivity of the formed dye reached to  $1.92 \times 10^4 \text{ l.mol}^{-1} \text{ cm}^{-1}$  at 537 nm and Beer's law is obeyed within the range of 10-300  $\mu\text{g}$  of tetracycline-HCl /10 ml (i.e.1.0 – 30 ppm). The colour reaction was highly stable, a relative error ranged from - 2.5 to + 0.21 % and a relative standard deviation of  $\pm 0.37$  to  $\pm 0.73$  %, depending on concentration level of tetracycline-HCl. The proposed method has been applied successfully to the determination of tetracycline-HCl in pharmaceutical preparations.

**Keywords:** Tetracycline-HCl; 4-aminoantipyrine; diazo-coupling; spectrophotometry.

-4

-4

$1.92 \times 10^4 \text{ l.mol}^{-1} \text{ cm}^{-1}$

300 10

537

( 30-1 ) 10 /

% 0.73 $\pm$  0.37 $\pm$

% 0.21 - 2.5

## INTRODUCTION

Tetracycline antibiotics are potent, broad spectrum antibacterial agents effective against a host of Gram positive and Gram negative aerobic and anaerobic bacteria. As a result, the tetracyclines are drugs of choice, or well accepted alternatives for a variety of infectious diseases (Albert, 1953). Assay of tetracycline has previously been achieved by several analytical techniques such as high performance liquid chromatography (Shalaby *et al.*, 2011), RPHPLC (Johnathan and Yugang, 2007; Kazuo *et al.*, 1992), LC (Innocenzo and Fabio, 2009) LC-Mass (Pena *et al.*, 2007), fluorimetry (Regos *et al.*, 1978), adsorptive voltammetry (Parviz *et al.*, 2007), flow injection (Saisunee *et al.*, 2006; Jose *et al.*, 2009; Prinya *et al.* 2008). However, many of these methods require expensive equipments and skilled operation. The most widely methods used for the determination of tetracycline are spectrophotometric methods such as: complexation with iron (Ronald, 2007), oxidative coupling (Ayad *et al.*, 1986), diazo-coupling (Al-Abbasi, 2009), charge transfer complex (Khairi, 2008), kinetic spectrophotometric method (Yongnian *et al.*, 2010). However, some of these methods suffer from one or more disadvantage such as insufficient sensitivity, poor selectivity or using of solvent extraction procedures. Therefore, a simple and accurate method for assaying of tetracycline in different pharmaceutical preparations is necessary for routine analysis.

## EXPERIMENTAL

### Apparatus

All absorption measurements were obtained by using a double beam CECIL CE7200 UV-Visible Recording Spectrophotometer with 1.0 cm matched quartz cells and the pH measurements were performed by using HANNA Instrument pH 211 microprocessor pH meter.

### Reagent

All Chemicals used are of analytical-reagent grade.

A pure tetracycline - HCl (TCH) was obtained from the State Company for Drug Industries and Medical Appliances (SDI), Sammara, Iraq. A solution of  $100 \mu\text{g ml}^{-1}$  was prepared by dissolving 0.01g of tetracycline -HCl in 100-ml distilled water. Sodium carbonate (1M) and different interferences solution ( $1000 \mu\text{g ml}^{-1}$ ) were prepared by dissolving the proper amounts in distilled water. The diazotised 4-aminoantipyrine (2.5 mM) solution was prepared by dissolving 0.05 g of 4-aminoantipyrine (Fluka) in 60 ml distilled water then 3ml of concentrated hydrochloric acid was added followed by heating. Finally the mixture was transferred to a 100-ml volumetric flask and cooled at  $0-5^{\circ}\text{C}$  in an ice-bath. A 0.017 g sodium nitrite was added and the mixture was stirred vigorously. After 5 minutes, the solution was made up to 100 ml with cold distilled water and was kept in brown bottle in a refrigerator for 1 hour before using, and it is stable for at least five days.

### General procedure and calibration graph

The aqueous solution (0.1 – 3ml) containing tetracycline-HCl ( $100 \mu\text{g ml}^{-1}$ ) was transferred into a series of 10-ml calibrated flasks. To each flask 1 ml of diazotised 4-aminoantipyrine solution (2.5 mM), 2 ml of sodium carbonate solution (1M) and 4 ml of CPC ( $1 \times 10^{-3}\text{M}$ ) were added then the volume was completed to the mark with distilled

water. The absorbance was measured at 537 nm against a reagent blank which was prepared in a similar way but without the addition of tetracycline-HCl. The calibration graph as shown in Fig. ( 1) was linear over the range of 10-300  $\mu\text{g}$  of tetracycline-HCl /10 ml (i.e 1– 30 ppm).The apparent molar absorptivity referred to tetracycline – HCl has been found to be  $1.92 \times 10^4 \text{ l.mol}^{-1}.\text{cm}^{-1}$ .




Fig. 1: Calibration graph for tetracycline – HCl determination using the proposed method.

#### **Determination of tetracycline in capsule**

Weight and mix the contents of five capsule (each one contains 250 mg tetracycline - HCl), an accurately weighed amount of powder equivalent to 0.01g tetracycline - HCl was dissolved in 100 ml distilled water in a volumetric flask. A suitable aliquot of solution was taken and the recommended procedure was followed for analysis of the drug .

#### **Determination of tetracycline in ointment**

Mixed well the contents of four containers of tetracycline ointment , an accurately weighed amount equivalent to 0.01g tetracycline - HCl was dissolved in a mixture of 3 ml ethanol with 50 ml distilled water, then the mixture was warmed and filtered into a 100- ml volumetric flask and the volume was completed to the mark with distilled water. A suitable aliquot of solution was taken and the recommended procedure was followed for analysis of the drug.


### **RESULTS AND DISCUSSION**

The effect of various variables on the colour development was tested to establish the optimum conditions for the determination of TCH by coupling with diazotized 4-aminoantipyrine reagent.

### Principle of the method

The method included the following steps:


- Preparation of diazotised 4-aminoantipyrene



4-aminoantipyrene

Diazotised 4-aminoantipyrene

- Coupling of tetracycline with diazotised 4-aminoantipyrene



Tetracycline

Diazotised 4-aminoantipyrene

Red azo dye

### Choice of diazotised agent and amount

Several aromatic diazotised agents have been tested for optimum conditions. The results in Table 1 show that p-phenyldiamine give the highest intensity but with low value of colour contrast, while 4-aminoantipyrene gives good sensitive reaction ( $\epsilon = 1.85 \times 10^4 \text{ l.mol}^{-1}.\text{cm}^{-1}$ ) with high value of colour contrast (165 nm) in alkaline medium. Therefore, it has been selected for the subsequent experiments.

Table 1: The selection of diazotised agent.

Reagent( 2.5 mM)	Absorbance	$\lambda_{\text{max}}(\text{nm})$	$\Delta\lambda_{\text{max}}(\text{nm})$	$\epsilon (\text{l.mol}^{-1}.\text{cm}^{-1})$
m-Nitroaniline	0.261	352	99	$1.26 \times 10^4$
p-Phenyldiamine	0.404	352	42	$1.95 \times 10^4$
Sulphanilic acid	0.114	352.5	102.5	$0.55 \times 10^4$
4-aminoantipyrene	0.384	537	165	$1.85 \times 10^4$

\* $\Delta\lambda_{\text{max}} = \text{Colour contrast} = \lambda_{\text{max}}\text{S} - \lambda_{\text{max}}\text{B}$  where S=The dye, B=Blank.

The effect of different amounts (0.5 – 2 ml) of diazotised 4-aminoantipyrene reagent on the absorbance of solutions containing different amounts of tetracycline-HCl (50-250  $\mu\text{g}/10\text{ml}$ ) have been studied. The obtained results indicated that the absorbance increases

with increasing the reagent concentration (diazotised 4-aminoantipyrine) and reached a maximum on using a volume of 1.0 ml of 2.5 mM diazotised 4-aminantipyrine, which also gives the highest value of determination coefficient ( $r^2 = 0.9938$ ). Therefore, 1.0 ml reagent was recommended for the subsequent experiments.

### Effect of base

Preliminary experiments have been shown that TCH gives coloured dye with diazotised 4-aminoantipyrine only in basic medium. Various bases (strong and weak) have been used and their results in Table 2 indicate that 2ml of  $\text{Na}_2\text{CO}_3$  (1M) give the highest intensity of the coloured dye, therefore this volume has been recommended for the subsequent experiments.

Table 2: The effect of base on absorbance .

Amount of base used (ml)	Absorbance/base used (1M)			
	NaOH	KOH	$\text{Na}_2\text{CO}_3$	$\text{NaHCO}_3$
0.25	0.015	0.018	0.242	0.009
0.5	0.103	0.019	0.379	0.194
1.0	0.061	0.053	0.386	0.350
2.0	0.057	0.045	0.413	0.354
3.0	0.050	0.032	0.413	0.388
pH range	(2.17- 13.16)	(1.98-13.20)	(6.68-10.73)	(3.75-9.73)
* $\Delta \lambda_{\text{max}}$	135	137	165	165

\* $\Delta \lambda_{\text{max}}$  = Colour contrast  $\lambda_{\text{max}}\text{S} - \lambda_{\text{max}}\text{B}$

where S=The dye, B=Blank

### Effect of surfactant

The effect of several types of surfactants on colour intensity of the dye has been investigated. (Table 3 and Fig. 2).

Table 3: The effect of surfactant on dye absorbance.

* Order	Absorbance / ml of CPC ( $1 \times 10^{-3}\text{M}$ )			Absorbance/ml of SDS ( $1 \times 10^{-3}\text{M}$ )			Absorbance/ml of Triton-X100 (1%)		
	1 ml	3 ml	4 ml	1 ml	3 ml	4 ml	1 ml	3 ml	4 ml
I	0.282	0.263	0.242	0.409	0.407	0.380	0.409	0.407	0.392
II	0.283	0.275	0.244	0.405	0.398	0.386	0.389	0.405	0.420
III	0.429	0.445	0.458	0.401	0.386	0.374	0.397	0.401	0.415

Note Absorbance without surfactant = 0.412

I\* = Sample (S) + Surfactant( Sur.) + diazotised (R) + Base (B)

II = S+R+Sur.+B

III = S+R+B+Sur.




Fig. 2: The effect of CPC on absorbance (order III)  
A- Sample with CPC B- Sample without CPC

The results indicate that the addition of CPC in the order (III) increases the intensity of the formed dye, therefore it has been recommended to use 4 ml of CPC ( $1 \times 10^{-3}$  M) in this order in subsequent experiments.

#### Effect of time and amount of TCH on absorbance

The effect of time on the development and stability period of the formed coloured dye was investigated under optimum experimental conditions described before. The formation of coloured dye being complete after mixing the components of reaction and the absorbance of the coloured species remained constant for at least 60 minutes. (Table 4).

Table 4: The effect of time on absorbance.

Time/min.	Absorbance/ $\mu\text{g}$ of tetracycline-HCl		
	100	200	250
0	0.454	0.776	1.006
5	0.458	0.778	1.004
10	0.456	0.774	0.998
20	0.457	0.775	0.995
30	0.454	0.776	0.993
40	0.456	0.774	0.990
50	0.455	0.775	0.989
60	0.450	0.774	0.987

### Final Absorption Spectra

An absorption spectra of the formed coloured dye by coupling of TCH with diazotised 4-aminoantipyrine in the presence of CPC and in basic medium, against its corresponding reagent blank shows a maximum absorbance at 537 nm in contrast to the reagent blank which shows maximum absorbance at 425 nm ( Fig 3).




Fig. 3: Absorption spectra of 100µg TCH treated according to the recommended procedure and measured against (A) blank (B) distilled water and (C) blank measured against distilled water

### INTERFERENCE

The criterion of interference was an error of not more than  $\pm 5.0\%$  in the absorbance. To test the efficiency and selectivity of the proposed analytical method, a systematic study of excipients (e.g. , glucose , lactose , gum Arabic and starch) at various levels, that usually present in dosage forms. Experimental results showed that there was no interference from additives or excipients up to 1500 µg in the present method as shown in Table 5.

Table 5: Effect of foreign compounds for assay of tetracycline-HCl.

Foreign compound	Recovery (%) of 100 µg tetracycline-HCl per µg foreign compound added		
	500	1000	1500
Glucose	99.3	100.0	98.9
Gum Arabic	100.0	99.7	99.3
Lactose	100.2	100.6	100.4
Starch	100.2	99.5	100.6



### The effect of organic solvents

Different organic solvents have been examined to evaluate their effects on the spectrum of the resulting azo dye, the results are given in Table 7 and Fig. 4

Table 7: The effect of organic solvents compared with water.

Solvent	$\lambda_{\max}$ , nm	Absorbance	$\epsilon$ l.mol <sup>-1</sup> .cm <sup>-1</sup>
Ethanol	554	0.396	$1.91 \times 10^4$
Formic acid	561	0.233	$1.12 \times 10^4$
Methanol	606	0.396	$1.91 \times 10^4$
Acetic acid	612	0.051	$0.25 \times 10^4$
Water	537	0.451	$2.178 \times 10^4$




Fig. 4: The effect of solvents

Although formic acid and acetic acid caused red shift, water still used in the subsequent experiments according to the highest sensitivity.

### Application of the method

The proposed method was applied to determine tetracycline- HCl in its pharmaceutical preparations(capsules and skin ointment). The results which are shown in Table 8 indicate that a good recovery was obtained.

Table 8: Analytical applications of the proposed method.

Pharmaceutical preparation	Amount taken ( $\mu\text{g}$ )	Amount measured	Recovery
Samacycline 250mg/Capsules (S.D.I Iraq)	50	49.7	99.4
	100	100.2	100.2
Apecycline 250mg/Capsules (PVT.SECTOR India)	50	49.7	99.4
	100	99.9	99.9
Samacycline 3% Skin ointment ( S.D.I Iraq)	50	49.7	99.4
	100	100	100

#### Average of four determinations

The results obtained are in the agreement with certified values compared with standard addition method (Fig. 5 and Table 9).





Fig. 5 : Standard addition plot for the recovery of 50 and 100 mg of TCH in S.D.I Iraq capsule and PVT.SECTOR India capsule.

Table 9: The results of standard addition method.

Pharmaceutical preparation	Amount taken (mg)	Amount measured(mg)	Recovery, %
Samacycline 250mg /Capsules S.D.I Iraq	50	50	100
	100	98.5	98.5
Apecycline 250mg/Capsules PVT. SECTOR. India	50	50	100
	100	101	101

The performance of the proposed method for determination of tetracycline in ointment was assessed by calculation of the t-test compared with the standard method (British Pharmacopeia, 2000) (spectrophotometric method ) for 95% confidence level with eight degrees of freedom. The results (Table 10) showed that the t-value was less than the critical value, indicating no significant difference between the proposed and standard method for the determination of tetracycline in ointment.

Table 10: Analytical applications of the proposed method and t-values experimental.

Drug	Recovery, % *		t. exp
	Present method	British pharmacopeia method <sup>(14)</sup>	
Samacycline Skin ointment S.D.I (Iraq)	99.70	99.55	0.459

\*Average of five determinations of 100 µg TCH.

### Comparison of the methods

Table(11) shows the comparison between some of analytical variables for the present method with that of another literature spectrophotometric methods.

Table 11: Comparison of the methods.

Technique	Reaction	Reagent	$\lambda_{\max}$ (nm)	Berr's law	$\epsilon$ (l.mol <sup>-1</sup> .cm <sup>-1</sup> )	Reference
FI**/Spec.*	Complexation	Aluminum chloride	376		.....	Saisunee, <i>et al.</i> , 2006
FI/Spec.	Oxidation	Chloramine -T	535	$6.62 \times 10^{-5}$ -. $72 \times 10^{-4}$	.....	Jose <i>et al.</i> , 2009
Spec.	Oxidative coupling	4-Aminoantipyrine	520	0.04-12 mg/ml	.....	Ayad <i>et al.</i> , 1986
Spec.	Complexation	Fe <sup>+3</sup>	435	$2.81 \times 10^{-3}$ -. $50 \times 10^{-4}$	.....	Ronalad, 2007
Spec.	Charge transfer complexation	Chloranilic acid	540	2.5-30.0 mg/ml	.....	Khairi, 2008
Spec.	Diazo coupling	Diazotised p-nitro aniline	569	2-400 $\mu$ g/25ml	$7.8 \times 10^4$	Al-Abbasi, 2009
FI/ Spec.		Uranyl acetate	410	1.0-3.0 $\mu$ g/ml	.....	Prinya <i>et al.</i> , 2008
Spec.	Diazo coupling	Diazotised 4-aminoantipyrine	537	1.0-3.0 $\mu$ g/ml	$1.923 \times 10^4$	Present method

\*Spec.= Spectrophotometric method

\*\* FI= Flow injection method

The results indicate that the proposed method has a good sensitivity and has a wide application part in determination of drug under investigation in its pharmaceutical preparations.

### CONCLUSION

The proposed method for the determination of tetracycline -HCl is simple and has a good sensitivity. The proposed method has advantageous over some of the reported visible spectrophotometric methods with respect to reproducibility, precision, accuracy and stability of the coloured species. The proposed method is suitable for the determination of tetracycline -HCl in pure form and in its pharmaceutical preparations.

### REFERENCES

- Al-Abbasi, K. (2009). Spectrophotometric determination of tetracycline by azo dye formation with diazotised p-nitroaniline –application to pharmaceutical preparation and biological fluids. *J. Raf. Sci.*, **20**, 61-74.
- Albert, A. (1953). Avidity of tetracycline and aureomycin for metallic cations. *Nature* . **172**, 201-211.

- Ayad, M.; El-Sadik, M. ; Mostaffa, S. (1986). 4-Aminoantipyrine as an analytical reagent for the colorimetric determination of tetracycline and oxytetracycline. *Anal. Lett.*, **19**, 2169-2181.
- " British Pharmacopeia on CD- ROM ", (2000). 3rd edn., System Simulation Ltd , the stationary office, London ,
- Hargis, L.G. (1988). "Analytical Chemistry Principles and Techniques". Prentice-Hall Inc, New Jersey, 424-427 p.
- Innocenzo, G.; Fabio, P. (2009). Determination of tetracycline residues by liquid chromatography coupled with electrochemical detection and solid phase extraction. *J. Agric. Food Chem.*, **57**(19), 8735-8741.
- Johnathan, W.; Yugang, Z. (2007). Simultaneous determination of tetracycline, oxytetracycline and 4-epitetracycline in milk by high-performance liquid chromatography . *Food Chem.* , **103**, 1297-13 .
- Jose, L.; Patricia, L.; Helena, R.; Pezza, L. (2009). Flow injection spectrophotometric determination of tetracycline and doxycycline in pharmaceutical formulations using chloramines-T as oxidizing agent. *Quim Nova*, **32**(7), 1764-1769.
- Kazo, I.; Norio, O.; Mitsuru, Y. (1992). Determination of tetracycline antibiotics by reversed phase high-performance liquid chromatography with fluorescence detection. *J. Chromatog. A*, **623**, 153-158.
- Khairi, F. (2008). Analysis of tetracycline and oxytetracycline through charge transfer complexation . *Amer. J.*, **3** (3), 212-218.
- Parviz, N.; Reza, M.; Parandis, D. (2007). " FFT –Adsorptive voltametric technique for pico-level determination of tetracycline in capsules at an micro electrode in flowing solutions" . *Turk. J. Chem.* , **31**, 279-291.
- Pena, A.; Lino, M.; Alonso, D.; Barcelo, D. (2007). Determination of tetracycline antibiotic residues in edible swine tissues by liquid chromatography with spectrofluorometric detection and confirmation by mass spectrometry, *J. Agriculture and Food Chem.* , **55**, 4973-4979
- Prinya, M.; Saisunee, L.; Suphachock, U. (2008). Spectrophotometric flow-injection analysis assay of tetracycline antibiotics using a dual light –emitting diode based detector . *Mj. Int. J. Sci.Tech.* , **2** (1), 201-209
- Regos, A. ; Zuk, G. ; Lukasiak, J. D. (1978). "Flourimetric determination of tetracycline hydrochloride". *Sci. Pharm.*, **46**,249-254; *Anal. Abst.* (1979), **37**, 1E37.
- Ronald, B. (2007). Determining trace levels of tetracycline in solid and liquid samples utilizing complexation with transition metal cation. *Anal. Chem.*, **3**, 4-6.
- Saisunee, L.; Boonsom L.; Surasat W.; Wirt R. (2006). Flow injection spectrophotometric determination of tetracycline in a pharmaceutical preparation by complexation with aluminium III. *Anal .Sci.***22**,15-18.
- Shalaby, A. ; Ndia, A.; Abou-Raya, S.; Wafaa, H. ; Mehaya, F . (2011). Validation HPLC method for determination of tetracycline residue in chicken meat and liver. *Food Chem.* , **124**(4), 1660-1666.
- Yongnian, Ni .; Dong, Na. ; Serge, K. (2010). " A simple kinetic spectrophotometric method for simultaneous determination of tetracycline by use of chemometrics". *Anal. Methods* , **2** ,1302 -1309.