

**The effect of varicocele treatment on human sperm plasma membrane integrity and pregnancy rate in infertile men after varicelotomy**

**Bassim KH. Kouti      \*Mohammed – Baqer M.Fakur Al deen**

**College of Sciences - university of Thi-Qar**

**\*IVF institutie of embryo Research and infertility treatment Al-**

**Nahrain university-applied embryology department**

**Abstract**

This study was designed to evaluate the results of semen parameters and hypo-osmotic swelling (HOS) test for semen samples of infertile patients with and without varicocele undergoing intra-uterine insemination (IUI) techniques. Twenty-four semen sample from infertile patients with varicocele and fifty-four without varicocele were collected by masturbation and evaluated after and before *in vitro* sperm activation.

Spermatozoa prepared by conventional layering technique by using 1ml of the liquefied semen were layered beneath a culture medium (IVF medium). The HOS test was performed after and before *in vitro* sperm preparation by mixing 0.1 ml of semen with 1.0 ml of a 150 mOsm/ L NaCl as a hypo-osmotic solution. Sperm function tests (SFT) including sperm concentration, sperm motility (%), progressive sperm motility (%), and normal sperm morphology (%) were evaluated according to standard World Health Organization (WHO) criteria and subjected to HOS test. For IUI, the sperm prepared and incubated for 30 minute in 5% CO<sub>2</sub> at 37°C after *in vitro* sperm preparation.

The results of the present study reported that successful pregnancy rates was more chance for semen sample with normal HOS test scores than abnormal HOS test. Eighteen clinical pregnancy rates were achieved for non-varicocelic infertile patients with normal HOS test scores and have SFT recovered than *in vitro* sperm activation as compared to varicocelic infertile patients with abnormal HOS test scores. Therefore, it was concluded that the use of HOS-test as an easy and reliable test to identify successful pregnancy after IUI technique. Further studies are recommended to assess the presence of varicocele on the percentage of hamster ova penetration assay after IVF-ET.

## **Introduction**

The presence of varicocele in the general male population is 15-20% (1). Also, varicocele causes infertility in 30-50% of all marriages (2). An abnormal spermogram is more frequent in men with varicocele than in those with normal parameter findings (3). According to WHO (4), the infertility of a great number of couples is caused by varicocele, which is the main cause of male infertility (30-50%). Moreover, Varicocele is associated with a decrease in fertility and testicular functions. If varicocele is a consequence of a congenital lack in the vein drainage of the testicular groove, it can appear between the ages of 13 and 15 (5) and its occurrence is caused by the presence of vein reflux (6). In general, damage to spermatogenesis caused by the negative influence of varicocele occurs as the result of the integrated effect of increased testicular temperature and the slower flow and stagnation of blood in the pampiniform plexus due to the effect of prostaglandins and serotonin. In the pathogenesis of varicocele, gradual damage is done to the germ cell epithelia and Leydig cells (7).

The sperm functions tests are the best way to investigate male fertility and considered the most important factors in male reproduction potential (8). In general, semen analysis is the cornerstone and primary laboratory test for the evaluation of male fecundity (9), and useful in predicting the results of assisted reproductive technology (10). Sometimes, fertilization occurs despite an abnormal semen analysis, or it fails to occur when analysis values are normal. Moreover, sperm functions tests are based on the detection of sperm count, percentage of sperm motility, progressive sperm motility (%), percentage of normal sperm morphology (%) and sperm viability (%) (11, 12).

The Hypo-osmotic swelling test is a simplest and less expensive test for assessing the functional integrity of sperm plasma membrane (13). The HOS test introduced by

Jeyendran *et al.* (1984) has added a new parameter for the evaluation of human spermatozoa. The functional integrity of sperm plasma membrane is an important factor in sperm metabolism, capacitation, acrosome reaction, and binding of spermatozoa to the egg surface (14). Nevertheless, in the presence of normal semen parameters, an HOS test less than 50% is rarely associated with pregnancy *in vivo* (15). The HOS test was demonstrated as an important test and indicator for male factor infertility and success of intra-uterine insemination. However, HOS test predict pregnancy rate and outcome in couples undergoing IVF and IUI procedure (16, 17). Therefore, in the current study, sperm function test (SFT), sperm HOS test, and IUI were applied to observe the diagnostic tools of HOS test in predicting pregnancy rates and outcome in couples undergoing IUI performance.

## **Materials and Methods**

### **1. Subjects**

Twenty-four infertile patients with varicocele and fifty-four infertile patients without varicocele were obtained from IVF Institute of Embryo Research and Infertility Treatment/Al-Nahrain University. The study extended from September, 2006 to March, 2007. The selection of varicocelic infertile patients was based on the physical examination performed with the patients in both recumbent and upright position. The varicocele was confirmed by palpable on physical examination of the scrotum by the clinical evaluation in order to measure the severity of varicocele. Therefore, ancillary diagnostic measures, such as scrotal ultrasonography and Doppler examination were done.

### **2. Laboratory assessments**

#### **1. Semen analysis and collection**

The sample of seminal fluid was collected after 3-5 days of abstinence directly in a clean, dry and sterile disposable Petri-dish by

masturbation in a private and quite room adjacent to the semen analysis laboratory. The container labeled with the following information, name, age, abstinence period and time of sample collection. The specimens were placed in an incubator at 37°C for 30 minutes to allow the semen liquefaction (18). The liquefied semen is then carefully mixed for few seconds, and then specimen was examined in details by microscopic and macroscopic examinations including semen volume, semen liquefaction time, semen viscosity, semen pH, sperm concentration, sperm motility (%), progressive sperm motility (%), normal sperm morphology (%). The WHO criteria for normal semen values were applied (19).

## 2. *In vitro* sperm activation and preparation

Sperm processing for intra-uterine insemination were done using a conventional layering technique. For this, 1ml of the liquefied semen was layered beneath a culture medium (IVF medium, Medi-cult, Jyllinge, Denmark), after incubation for 30 minute in 5% CO<sub>2</sub> at 37°C. The supernatant was removed and was used for treatment. Intra-uterine insemination was performed using a Portex catheter. The cervix was exposed and the catheter was passed into the uterus to about 0.5 cm from the top of the uterine cavity. The sperm were then expelled. However, the sperm parameters were examined after 10 minutes of incubation and a one drop (10µl.) of the mixture was added to a clean class and examined at light microscope under 400X objective for assessment of HOS test score for both groups of infertile patients.

## 3. Hypo-Osmotic Swelling (HOS) Test

The HOS test was performed before and after *in vitro* sperm preparation technique and examination of standard semen parameters by mixing 0.1 ml of semen with 1.0 ml of a 150 mOsm/ L NaCl as a hypo-osmotic solution (20). The mixture was incubated for 30 minute at 37 °C in 5% CO<sub>2</sub> in accordance with a previously described technique (21). One

hundred spermatozoa were examined, and the morphological changes in sperm tail were classified according to the types described by Jeyendran *et al.* (22). The overall rate of sperm swelling was calculated. Finding at least 50% swollen spermatozoa was considered normal.

## 4. Statistical analysis of the data

Statistical analysis was performed with the SPSS version 12.00 by the Statistical Package for Social Sciences Software. The data analysis was done using paired sample t-test to assess the statistical differences in the results of SFTs and sperm HOS test for both varicocele and non-varicocele infertile patients. Mean and standard error of mean (S.E.M) obtained from crude data to compare between Pre-and Post-activation for semen parameters. P-value < 0.05 was used as a level of statistically significant.

## 5. Results

Twenty-four varicocele infertile patients with mean age (30.83 ± 1.33 years) and duration of infertility (5.66 ± 0.33 years) and fifty-four infertile patients without varicocele with mean age (31.24 ± 0.89 years) and duration of infertility (3.16 ± 0.21 years) were involved in the present study. The results of the present study showed that varicocele infertile patients have least value for sperm functions and sperm HOS test (36.40 %) as compared to non varicocele infertile patients (51.29 %) before *in vitro* sperm preparation. However, it was noticed a highly significant (P<0.001) differences in sperm functions were assessed post *in vitro* sperm activation for both groups as compared to pre-activation. Meanwhile, significant (P<0.001) and markedly reduction in sperm concentration were observed for *in vitro* post-activation. In contrast, a highly significant (P<0.001) differences in the percentage of sperm motility (%), progressive sperm motility (%), normal sperm morphology (%), and HOS test were noticed after sperm activation and processing. In general, in the present study, the best results for clinical

pregnancy rates were observed in the infertile patients without varicocele (18%) after IUI techniques. It was advisable, clinical pregnancies were achieved for infertile couples where their male have normal percentages of

the sperm HOS test scores as compared to male partners have abnormal HOS-test scores with varicocele, who have no clinical pregnancy rates were achieved in the present study.

**Table (1): Sperm functions tests and sperm hypo-osmotic swelling test pre-and post-activation *in vitro* for human spermatozoa for non-varicocele infertile patients\* undergoing IUI technique**

Parameters	Conventional layering technique	
	Pre-activation <i>in vitro</i>	Post-activation <i>in vitro</i>
Sperm Concentration ( $\times 10^6$ sperm/ml)	59.18 $\pm$ 2.37	32.64 $\pm$ 1.48 a
Sperm Motility (%)	63.05 $\pm$ 0.96	85.92 $\pm$ 0.89 a
Progressive sperm Motility (%)	39.64 $\pm$ 0.99	60.38 $\pm$ 1.24 a
Normal Sperm Morphology (%)	54.25 $\pm$ 1.46	86.20 $\pm$ 0.85 a
Sperm HOS test (%)	51.29 $\pm$ 1.01	66.05 $\pm$ 1.07 a

Number of infertile patients=54  
Values are Mean  $\pm$  S.E.M


a: Means a highly significant ( $P < 0.001$ ) difference between pre-and post-activation *in vitro* for human spermatozoa.

**Table (2): Sperm functions tests and sperm hypo-osmotic swelling test pre-and post-activation *in vitro* for human spermatozoa for varicocele infertile patients\* undergoing IUI technique**

Parameters	Conventional layering technique	
	Pre-activation <i>in vitro</i>	Post-activation <i>in vitro</i>
Sperm Concentration ( $\times 10^6$ sperm/ml)	18.20 $\pm$ 2.94	9.75 $\pm$ 1.04 a
Sperm Motility (%)	43.54 $\pm$ 1.54	71.50 $\pm$ 1.82 a
Progressive sperm Motility (%)	26.08 $\pm$ 1.48	51.87 $\pm$ 1.83 a
Normal Sperm Morphology (%)	43.64 $\pm$ 1.70	83.54 $\pm$ 1.14 a
Sperm HOS test (%)	36.40 $\pm$ 1.24	45.58 $\pm$ 1.04 a

Number of infertile patients=24  
Values are Mean  $\pm$  S.E.M

a: Means a highly significant ( $P < 0.001$ ) difference between pre-and post-activation *in vitro* for human spermatozoa.



**Figure (1): Outcomes of intra-uterine insemination for varicocele and non-varicocele infertile patient enrolled in this study**

## **Discussion**

In the present study, sperm functions tests, hypo-osmotic swelling (HOS) test, and IUI were evaluated to scrutinize the competence of HOS test in predicting pregnancy rate and outcome in varicocele and non-varicocele infertile patients undergoing intra-uterine insemination. A highly significant ( $P < 0.001$ ) differences in sperm concentration, percentage of sperm motility (%), progressive sperm motility (%), normal sperm morphology (%) were observed as compared to pre-activation for both study groups. However, only active motile mature normal sperm were able to swim to upper layer of the culture medium. A highly reduction in sperm concentration post *in vitro* sperm activation as a result of sperm preparation techniques and selection of highly recovery of spermatozoa.

The percentage of HOS-test statistically significant ( $P < 0.001$ ) differences between pre-activation and post-activation of human spermatozoa for varicocele and non-

varicocele infertile patients (Table 1, 2), but the best results and response were reported in non-varicocele infertile patients. The peroxidative damage of unsaturated fatty acids in sperm plasma membrane induced by reactive oxygen species (ROS) lead to low scores of HOS test and defective sperm function in patients with varicocele as compared with non-varicocele infertile patients (23). However, ROS is associated with varicocele and correlate with fertility potential. A significantly elevated ROS levels and depressed total antioxidant capacity levels for infertile patients with varicocele related infertility as compared to those without varicocele (24). The markedly increased ROS generation in infertile patients with varicocele compared with without varicocele who suggesting a possible association of ROS with fertility (25).

The percentage of swollen spermatozoa was lower in the infertile patients with varicocele than without varicocele (26). Varicocele may cause injury to the sperm plasma membrane

integrity and lead to low scores of HOS test. Also, may cause deleterious effects by affecting sperm penetration and motility in the female genital tract which in turn may affect gamete interaction (27). In addition, improvement in sperm membrane swelling was detected in a group of non-varicocele infertile patients who achieved pregnancies as compared with those who failed to achieve pregnancies (28). In contrast, significantly better semen characteristics (concentration, motility, and normal morphology) as well as enhanced acrosin activity for non-varicocele infertile patients (29). Varicocele impairs and alters the mechanism of sperm binding to zona pellucida (ZP) and gamete membrane fusion, resulting in impaired potential fertilization (30). Additionally, the numbers of sperm cells bound to the zona pellucida (ZP) are significantly lower as compared with non-varicocele infertile patients (31). The semen parameters for non-varicocele infertile patients have SFTs and HOST better than post-activation *in vitro* of varicocele infertile patients (Table 1).

### References

- Greenberg Sh. (1977): Varicocele and male fertility. *Fertil. Steril.* 28: 699-706.
- Gorelick J. and Goldstein M. (1993): Loss of fertility in men with varicocele. *Fertil. Steril.* 59: 6-3.
- Hoekstra T. and Witt M. (1995): The correlation of internal spermatic vein palpability with ultrasonographic diameter and reversal of venous flow. *J. Uro.* 153: 82-84.
- World Health Organization (WHO). WHO manual for the standardized investigation, diagnosis and management of infertile male. Cambridge University Press, UK. 2000.
- Hargreave TB. (1994): Varicocele in male infertility. Springer Verlag, London, Berlin.
- Akbay E., Cayan S., Doruk E., Duce MN. and Bozlu M. (2000): The prevalence of varicocele and varicocele-related testicular atrophy in Turkish children and adolescents. *B.J.U.* 86: 490.
- Ombelet W., Pollet H., Bosmans E. and Vereecken A. (1997): Results of a questionnaire on sperm morphology assessment. *Hum. Reprod.* 12: 1015-20.
- Liu DY., Du Plessis YP., Nayudu PL., Johnston WI. and Baker HW. (1988): The use of *in vitro* fertilization to evaluate putative tests of human sperm function. *Fertil. Steril.* 49:272-7.
- Talbert LM., Hammond MG., Halme JO., Rand M., Fryer JG. and Ekstrom RD. (1987): Semen parameters and fertilization of human oocyte *in vitro*: a multivariable analysis. *Fertil. Steril.* 48:270-277.
- Sakkas D., Urner F. and Bianchi PG. (1996): Sperm chromatin anomalies can influence decondensation after intracytoplasmic sperm injection. *Hum. Reprod.* 11:837-43.
- Bostofte E., Serup J. and Robbe H. (1984): Interrelation among the characteristic of human semen, and a new system for classification of male infertility. *Fertil. Steril.* 41:95-102.
- Cepni I., Idil MH., Ocal P., Budak E., Elibol F., Sonol H., Duran A., Yaldyr F., Irez T. and Aksu MF.(1997): Sperm parameters and their effects on fertilization rate in conventional IVF-ET. *Mid. East. Fert. Soc. J.* 2: 155-161.
- Kiefer D., Check JH. and Katsoff D. (1996): The value of motile density, strict morphology, and the hypo-osmotic swelling test in *in vitro* fertilization with embryo transfer. *Arch. Androl.*37:57-60.
- Check JH., Baker A., Benfer K., Lurie D. and Katsoff D.(1996): Transfer of cryopreserved embryos improved pregnancy rates in patients with damage to the functional integrity of the sperm membrane as measured by the hypo-osmotic swelling test. *Fertil. Steril.* 65: 1241- 1244.
- Check JH., Epstein R., Nowroozi K., Shanis BS., Wu CH. and Bollendorf A. (1989): The hypo-osmotic swelling test as a useful adjunct to the semen analysis to predict fertility potential. *Fertil. Steril.* 52:159-16.
- Silverberg KM. and Turner T. (2001): Evaluation of sperm. In: Textbook of Reproductive Techniques, Laboratory and

- Clinical Perspectives. Gardner, D.K., Weissman, A., Howles, C.M., and Shoham, Z. (eds.), Martin Dunitz Ltd.; London. Pp.61-74.
17. Check JH., Katsoff D., Check ML., Choe JK. and Swenson K. (2001b): *In vitro* fertilization with intra cytoplasmic sperm injection is an effective therapy for male factor infertility related to subnormal hypo-osmotic swelling test scores. *J. Androl.* 22:261-265.
  18. Tartagni M., Schonauer MM. and Cicinelli E. (2002): Usefulness of the hypoosmotic swelling test in predicting pregnancy rate and outcome in couples undergoing intrauterine insemination. *J. Androl.* 23:498-502.
  19. World Health Organization (WHO). Laboratory Manual for the Examination of Human Semen and Semen-Cervical Mucus Interaction, 4th ed. Cambridge, Cambridge University Press UK. 1999; Pp.8-11.
  20. Liu J., Tasi Y., Katz E., Compton G., Garcia JE. and Baramki TA. (1997): High fertilization rates obtained after Intracytoplasmic sperm injection with 100% non-motile spermatozoa selected by using a simple modified hypo-osmotic swelling test. *Fertil. Steril.* 66:373-5.
  21. Jeyendran RS., Van Der Ven HH., Perez-Pelaez M., Crabo BG. and Zaneveld LJD. (1984): Development of an assay to assess the functional integrity of the human sperm membrane and its relationship to other semen characteristics. *J. Reprod. Fertil.* 70: 219-228.
  22. Jeyendran RS., Van der Ven HH. and Zaneveld LJD. (1992): The hypo-osmotic swelling test: an update. *Arch. Androl.* 29:105-106.
  23. Benoff S., Hurley IR. and Barcia M. (1997): A potential role for cadmium in the etiology of varicocele-associated infertility. *Fertil. Steril.* 67: 336-347.
  24. Hendin EN., Kolettis PN. and Sharma RK. (1999): Varicocele is associated with elevated spermatozoal reactive oxygen species production and diminished seminal plasma antioxidant capacity. *J. Urol.* 161: 1831-1834.
  25. Weese DI., Peaster ML. and Himsl KK. (1999): Stimulated reactive oxygen species generation in the spermatozoa of infertile men. *J. Urol.* 149: 64-67.
  26. Fuse H., Kazama T. and Katayama T. (1999): Hypo osmotic swelling test in patients with varicocele. *Arch. Androl.* 27: 149-154.
  27. Villanueva-Diaz CA., Vega-Hernandez EA. and Diaz-Perez MA. (1999): Sperm dysfunction in subfertile patients with varicocele and marginal semen analysis. *Andrologia.* 31: 263-267.
  28. Fuse H., Akashi T. and Fujishiro Y. (1995): Effect of varicocele on fertility potential: comparison between impregnating and nonimpregnating groups. *Arch. Androl.* 35: 143-148.
  29. El-Mulla KF., Kohn FM. and El-Beheiry AH. (1995): The effect of smoking and varicocele on human sperm acrosin activity and acrosome reaction. *Hum. Reprod.* 10: 3190-3194.
  30. Vigil P., Wohler C. and Bustos-Obergon E. (1995): Assessment of sperm function in fertile and infertile men. *Andrologia.* 26: 55-60.
  31. Yamamoto M., Hibi H., Tsuji Y. and Miyake K. (1994): The effect of varicocele ligation on oocyte fertilisation and pregnancy after failure of fertilization in *in vitro* fertilization-embryo transfer. *Hinyokika-Kyo.* 40: 683-687.

## تأثير علاج دوالي الخصية على كفاءة الغشاء البلازمي للنطفة البشرية ومعدلات الحمل لمرضى العقم بعد اجراء التدخل الجراحي

\*محمد باقر محمد رشاد فخر الدين  
\*معهد ابحاث الاجنة وعلاج العقم- جامعة النهرين  
قسم علم الاجنة التطبيقي

باسم خميس كوتي الركابي  
كلية العلوم- جامعة ذي قار  
قسم علوم الحياة

### الخلاصة

هدف هذه الدراسة الى تقييم معايير السائل المنوي و فحص كفاءة غشاء النطفة البلازمي لمرضى المصابين وغير المصابين بدوالي الخصية الذين تم ادخالهم في برنامج الاستمناء داخل الرحم. تضمنت هذه الدراسة (٢٤) مريضاً مصاباً بدوالي الخصية و(٥٤) مريضاً غير مصاب وجمعت العينات بطريقة الاستمناء وتم فحصها قبل وبعد اجراء عملية التنشيط خارج الجسم. عينات السائل المنوي تم تحضيرها و تنشيطها بوساطة التقنية الطباقية التقليدية. تم اجراء فحص كفاءة غشاء النطفة البلازمي قبل وبعد التنشيط. اختبارات وظائف النطف و المتضمنة تركيز النطف, النسبة المئوية لحركة للنطف, النسبة المئوية للتقدمية للنطف, والنسبة المئوية للسوية شكلياً تم تقييمها وفقاً لمقررات منظمة الصحة العالمية. سُجلت افضل النتائج الخاصة بالنسب المئوية للحمل للعينات ذات فحص كفاءة غشاء النطفة البلازمي الطبيعي مقارنة بغير الطبيعي. ثمانى عشرة حالة حمل تم الحصول عليها بالنسبة للمرضى غير المصابين بدوالي الخصية و ذات فحص كفاءة غشاء النطفة البلازمي طبيعي مقارنة بالمرضى المصابين بدوالي الخصية و ذات فحص كفاءة غشاء النطفة البلازمي غير طبيعي. لذلك نستنتج انه بالامكان استعمال فحص كفاءة غشاء النطفة البلازمي كاختبار ذات كفاءة عالية لمعرفة معدلات الحمل بعد اجراء الاخصاب داخل الرحم.