

# Treatment of Bilateral Vocal Cord Paralysis by ((810 Nanometer Wavelength)) Diode Laser

**Ammar Hadi Khammas**  
FICMS, DLO

**Abstract:**

**Purpose:** This is prospective study began in Jan. 2003 and concluded in April 2004, was undertaken to examine the benefits of 810 nanometer diode laser in treatment of four patient with bilateral vocal cord paralysis also to compare the results with conventional treatment

**Patients & Methods:** 810 nm diode lasers 15 watt was used in these cases under general anesthesia, and induction of anesthetic drug done through tracheostomy tube in all patients.

**Results:** All patients were decanulated “Tracheostomy tube removed”, the voice of all preserved within normal.

**Conclusion:** Laser surgery in this case has more benefit and advantage than conventional methods even if the patient need more than on session of laser operation because of high success rate, less complication and easy technique.

**Key words:** vocal cord paralysis, CO<sub>2</sub> laser, diode laser

**Introduction:**

The treatment of vocal cord paralysis remains one of the difficult problems in laryngology. This condition most commonly results from injury to both recurrent laryngeal nerves during thyroid surgery; less frequently it is caused by some other external injury to the larynx or neck, or may be due to neurological disorder. Initially these patients have a fairly adequate laryngeal airway with a breathy voice. After varying periods of time, while the voice improves, the airway becomes inadequate, leading to dyspnoea and stridor.

They invariably end up with a tracheostomy. Tracheostomy was required in about 51 cent of cases of bilateral paralysis, [1]. Various surgical methods

have been used from time to time Table 1. The main aim of the treatment has been to provide an adequate laryngeal airway so as to get rid of tracheostomy tube. Although woodman’s arytenoidectomy [2] and nerve-muscle pedicle reinnervation [3] have been reported to yield better results, the failure rates for both procedures have remained significantly high, 20 to 40 per cent for the former and 18 per cent for the latter [4].

The purpose of this paper is to present a preliminary report on a prospective study on the use of 810nm diode surgical laser in the management of bilateral vocal cord paralysis. Besides being an uncomplicated one, the procedure has so far given high percent success.

**Table 1 Various surgical methods have been used for treatment of Bilateral Vocal Cord Paralysis**

<p><b>Surgical procedure</b></p> <p><b>Hoover (1932):</b></p> <p><b>Lore (1936):</b></p> <p><b>Kelly (1941):</b></p> <p><b>Orton (1944):</b></p> <p><b>Woodman (1946)</b></p> <p><b>Thornell (1957)</b></p> <p><b>Miehleke (1974)</b></p> <p><b>Doyle et al. (1967):</b></p> <p><b>Tuker (1976)</b></p>	<p><b>Historical Review</b></p> <p><b>Submucous resection (SMR) of the vocal cord (VC)</b></p> <p><b>SMR of V.C. + Arytenoidectomy</b></p> <p><b>Extralaryngeal Arytenoidectomy</b></p> <p><b>Modified Arytenoidectomy</b></p> <p><b>Modified Arytenoidectomy +Cord Lateralization</b></p> <p><b>Intralaryngeal Arytenoidectomy +Cord Lateralization</b></p> <p><b>Anastomosis of the severed recurrent laryngeal nerve</b></p> <p><b>Nerve-Muscle Pedicle Reinnervation</b></p>
---	--

**Patients & Methods:**

Four patients with bilateral vocal cord paralysis on whom tracheostomy had been done, were managed at Al-Yarmouk teaching hospital during a period from January 2003 till April 2004, two of them due to thyroid surgery, the third one is due to head injury and the fourth due to idiopathic cause neurological, i.e intracranial lesion.

Endoscopic laryngeal laser surgery was done under general anesthesia. The type of laser used in this study and instrumentation, dose parameter, safety rules, procedure were applied as follow:

**Surgical Laser system:**

The Dio-med surgical laser delivering 15 watts model of continuous wave laser radiation, the laser energy is delivered to the treatment site by means of a flexible optical fiber coupled to the laser aperture.

The Dio-med surgical laser incorporate class IV GA AIAs (Gallium Aluminum Arsenide) diode laser with wavelength in the range 790-830 nm and visible ((aiming beam)) 4mw class IIIa diode laser with wave length in the range 630-660 nm.

**Procedure:**

Two techniques were used in management of BVCP, first technique I apply endoscopic resection of vocal process of arytenoid and resection of posterior third of ipsilateral vocal cord. This was done in two cases.

The other technique I apply endoscopic Arytenoidectomy with resection of posterior third ipsilateral vocal cord, this was done in other two cases.

**Instrumentation and Accessories:**

To expose the larynx, laryngoscope used which connects with light source for illumination; suspensor

use for fixation of laryngoscope in certain position these allow getting two hands free. Microscope 400 lens for magnification was used.

Bare fiber of 600 μm was used with specific laryngeal hand pieces with channel for suction clearance.

**Dose Parameter:**

The power was 5-10 watts, 5 watts were used in resection of vocal process of arytenoid, and 10 watts was used in arytenoidectomy. The laser mode is continuous, controlled by foot piece.

**Safety:**

Safety to the surgeon and personnel is mainly for eye protection by wearing goggle of WL 780-830 nm with optic density > 5.

Safety to the patient aiming of protection of eyes and also airway protection eyes covered with wet drapes. Protection of airway either by using anaesthetic tube specialized of laser operation; metallic spiral type. Or wrapping the proper tube with foil Aluminum paper to prevent ignition. In my study induction of anesthesia was done through tracheostomy tube in all patients.

**Evaluation of the Result postoperatively:**

Post operatively evaluation depends on successfully closing the tracheostomy which was present preoperatively. Or relieve respiratory distress if there is no tracheostomy. Also keep the speech acceptable.

I used other parameter for assessment in two patients who test the pulmonary function preoperatively and postoperatively. I found there was improvement in figure preoperatively as it was low to almost near normal postoperatively (normal figure 70-80%).

$$\frac{FEV_1}{FVC} = \frac{\text{Force expiratory volume in one sec}}{\text{Force Vital Capacity}}$$

**Results:**

Four patients, two of them thyroid surgery was cause of injury to both recurrent laryngeal nerves while in one case it was due head injury and other was due neurological disorder.

All tracheostomised immediately after injury or damaged to the nerve except case No. 1 by which tracheostomy was done two years after injury to nerve table No.2. All patients presented immediately after injury to the nerve with stridor except the case No. 1 as in table No.3 .All decanulated after diode laser

surgery except one case No. 4 decanulation was few days after surgery in cases 1,3 in which technique excision of vocal process of arytenoid cartilage and posterior third part of vocal cord was done. While decanulation was late in

Case (No. 2) and (No. 4) in which technique excision of arytenoid and posterior third part of vocal

cord was done table No.4. Evaluation of patient postoperatively can be considered

By pitch of voice, decanulation and pulmonary function test which was done in two cases (No. 1, 3) as in table No. 5.

**Table No. 2 Patient Summary**

<b>Patient</b>	<b>Age</b>	<b>Sex</b>	<b>Duration of BVCP</b>	<b>Tracheostomy</b>	<b>Cause of BVCP</b>
<b>1- BH</b>	24 y	♀	2 y	+ ve	<b>Thyroidectomy</b>
<b>2. AA</b>	50 y	♂	6 m	+ ve	<b>Neurological</b>
<b>3. Km</b>	46 y	♂	7 m	+ ve	<b>Thyroidectomy</b>
<b>4. SA</b>	15 y	♂	6 m	+ ve	<b>Head Injury</b>

**Table No. 3 Presentation of Patients Pre-operatively**

<b>Case No.</b>	<b>Speech</b>	<b>Breath</b>
1	<b>Good</b>	<b>Respiratory distress</b>
2	<b>Good</b>	<b>Stridor</b>
3	<b>Good</b>	<b>Stridor</b>
4	<b>Good</b>	<b>Stridor</b>

Table No. 4 Laser Techniques used in Management of BVCP

Case No.	Laser Technique	Follow Up	Time of Closure
1	Excision of vocal process of arytenoid with posterior part of V.C	Decanulated	After few days
2	Excision of arytenoid and posterior part of vocal cord	Decanulated	After two months
3	Excision of vocal process of arytenoids and posterior part V.C	Decanulated	After few days
4	Excision of Arytenoid and posterior part of vocal cord	Decanulated	After 2 months

Table No. 5 Criteria for evaluation of patients BVCP managed by diode laser post-operatively

Case No.	Speech	Breath	Pulmonary Function test	
			Pre-operative	Post-operative
1	Good	Closure	40% before tracheostomy 10% (tracheostomised)	70%
2	Good	Closure		
3	Good	Closure	10% (tracheostomised)	70%
4	Good	Closure		

**Discussion:**

BVCP causes sever airway compromise with good voice. The principle goal of the therapy is to improve the airway and at the same time maintain good voice quality traditionally; surgical options have included permanent tracheostomy with speaking valve, cordectomy, vocal cord lateralization and vocal cord reinnervation procedure. These performed endoscopically or through external approach.

At present time most commonly practiced procedure are arytenoidectomy either via external

approach (Woodman technique) or via endoscopic technique.

But now with development of CO<sub>2</sub> laser, the procedure of choice in treatment of BVCP is endoscopic arytenoidectomy with or without complete or partial resection of ipsilateral vocal cord with CO<sub>2</sub> laser [5]. With precision, great ease and in blood less field.

The CO<sub>2</sub> laser has been the most frequently used laser in the larynx. In this study I used diode 15 lasers in management of BVCP and I used two techniques, the first technique applied on case No. 1, 3 in which excision of vocal process of arytenoid

and posterior part of vocal cord was done and patients get benefit from operation as they were decanulated few days postoperatively with good voice. So diode laser by this technique provides a great degree of precision, an almost completely bloodless field, minimal postoperative oedema, and rapid postoperative healing without the formation of scar tissue and in comparison with CO<sub>2</sub> laser the advantage is the same except that the use of CO<sub>2</sub> laser allows the otolaryngologist surgeon to perform a precise operation through the relatively narrow field of the laryngoscope without the need for tissue manipulation (hands-off surgery).

Other technique used in cases 2, 4 in which excision of the arytenoid and posterior part of ipsilateral vocal cord was done. I faced difficulty peroperatively like time consuming on cutting the arytenoid with more bleeding, also postoperatively decanulation was late as in 2<sup>nd</sup> case and 4<sup>th</sup> cases. The time measured from starting cutting by laser till the end of operation by the first technique, it was a few minutes, while the second technique it was about one hour duration. Delayed decanulation in the second technique was due to odema of posterior laryngeal region. There are, however, other different techniques to perform laser endoscopic operation one of these techniques was first introduced by Strong et al in 1976 and Groft in 1984 in which wide resection of the true vocal cord with good result in a short series [6].

Shaheen in 1988 preferred to do posterior wedge excision of one vocal cord in conjunction with laser ablation of homolateral arytenoids [7]. Ossof et al. al in 1984 reported success in 10 of 11 patients by laser arytenoidectomy in which the cartilage was exposed and vaporized with laser [8]. Prasad (1985) also reported successful outcome in short series of 6 patients who had been subjected to CO<sub>2</sub> laser vaporization of the part of the cord and vocal process of arytenoid rather than whole arytenoid cartilage [9]. Dennes et al. al 1989 reported a successful and good result in these series of 6 patients who had been subjected to posterior partial cordectomy. Szmaja-Z 1993 used a technique of arytenoidectomy and ipsilateral posterior part cordectomy by CO<sub>2</sub> Laser [10]. The use of this method gave good result in preserving breathing and phonation; it was also emphasized that the fast process of healing and lack of granulation in operated field allow for the quick recovering patient. Remacle-M 1996 reported subtotal CO<sub>2</sub> laser arytenoidctomy, the principle modification involve preservation of thin posterior

shell providing good postoperative fixation of arytenoid region, the risk of aspiration is thus averted [11]. Also Izquierdo-Luzon-J 2001 reported the same procedure subtotal arytenoidectomy and posterior cordectomy with the CO<sub>2</sub> laser in treatment of bilateral vocal cord paralysis [12].

#### Conclusion:

Four patients with bilateral vocal cord paralysis who had worn tracheotomy tube for varying periods of time were treated by diode laser by using the technique excision of vocal process of arytenoid and posterior part of ipsilateral vocal cord give the same advantage of the CO<sub>2</sub> laser as bloodless field and ease procedure, the two patients operated on by this technique have been successfully decanulated.

In spite of problem of the other technique (excision of arytenoid and posterior part of ipsilateral vocal cord), it has an advantage over traditional surgical methods, as less instrumentation, less manipulation, less time consuming and easy technique.

Further studies are needed because the number of cases is low.

#### References:

- 1-Holinger, L. D., Holinger, P. C. and Holinger, P. H. (1976) Etiology of bilateral abductor vocal cord paralysis. *Annals of Otolaryngology, Rhinology and Laryngology*, 85:828-436.
- 2-Woodman, D. G. (1953) bilateral abductor paralysis. A survey of 521 cases of arytenoidectomy via the open approach as reported by ninety surgeons. *Archives of Otolaryngology*, 58:150-153.
- 3-Tucker, H. M. (1978) Human laryngeal reinnervation: Long-term experience with the nerve-muscle pedicle technique. *Laryngoscope*, 88:598
- 4-Tucker, H. M. (1983) Complications after surgical management of the paralysed larynx. *Laryngoscope*, 93:295-298.
- 5-Carruth, J.A.S. The principles of laser surgery. Scott-Brown's otolaryngology. 5th edition Basic Sci. ch. 23, 1987.
- 6-Carruth, J.A.S. (1985). Laryngeal microsurgery: instrumentation and technique. *J. laryngology and otology* 99:573.
- 7-Shaheen, O.H. 1988. Laser surgery recent advances in otolaryngology.

8-Ossoff, R.H. Sission, G.A. & Duncavage, J.A. 1984. Endoscopic arytenoidectomy for bilateral vocal cord paralysis laryngoscope 94:1293-1297.

9-Prasad, U (1985). CO<sub>2</sub> surgical laser in the management of bilateral vocal core paralysis. J. Laryngology & Otology 99:891-894

10-Szmeja-A; Wojtowicz-JG otolaryngal-pol 1993 47(3):221-6.

11-Remacle-M; Lawson; Mayne A; jamart-J Ann-otol-Rhinol-Laryngol. 1996 Jun. 105 (6): 438-45.

12-Izquierdo-Luzon,-J; Campos-catala-Armengo-Carceller-M; Basterra-Alegria- J.An.otorrinolaringol-Ibero-Am. 2001; 28(4)341-9.

---

From Surg Dept, Coll. of Med., Al- Mustansyrya Univ.